

**Robert Fogel**

Forest Science Department  
Oregon State University  
Corvallis, Oregon 97331

and

**James M. Trappe**

Pacific Northwest Forest and Range Experiment Station  
Forestry Sciences Laboratory  
Corvallis, Oregon 97330

## **Fungus Consumption (Mycophagy) by Small Animals**

### **Abstract**

A review of the natural history literature of small mammal feeding habits, based on many fortuitous field observations, some analysis of stomach contents, and a few feeding experiments, shows that diverse animals feed on similarly diverse fungi. The interdependence of animals and fungi has evolved to a high degree in some cases, e.g., the loss of alternative spore dispersal mechanisms by some fungi and the strong reliance of some mammals on fungi as a primary food. Consequently, adaptation for mycophagy and the effects on mycophagy of habitat, fungal toxicity, and the food value of fungi have implications in the interpretation of ecosystem structure and function.

### **Introduction**

Truffles and other hypogeous fungi occupy a highly specialized, protected niche as subterranean-fruited symbionts with roots of higher plants. Such fungi depend on mammal and invertebrate mycophagy for spore dispersal. These and other fungi, in turn, are important in the diet of many animals and in some cases comprise the major food source.

These interrelationships between fungi and animals engender many questions relevant to the life histories and ecosystematic roles of both. For example, which animals depend largely on fungi for food, for water, or for vitamins? What are the behavioral implications of seeking, drying, and caching fungi? What food values do mycophagists derive from fungi? Can animals differentiate between poisonous and nonpoisonous fungi? What is the phylogenetic significance of the various fungal mechanisms evolved for attracting mycophagists? What role does mycophagy play in fungus dispersal? What impact does mycophagy have on food chain and ecosystem dynamics, and where need it be recognized in ecosystem modeling?

The following literature review is intended to aid and stimulate a search for answers to such questions with emphasis on mammals, although similar questions can be asked about other animals as well. Little comprehensive research has been done on animal mycophagy. To interpret scattered fragments of published information, we have filled some gaps with speculation in hope of fostering further research in this fascinating field. Our literature search led to compilation of the reciprocal lists of mammal mycophagists and fungi eaten, presented later in this paper. Authorities for all species mentioned in the text are given in these lists.

### *The Small Mammal as Mycophagist*

Many fortuitous field observations, some analyses of mammal stomach contents, and

a few feeding experiments have revealed that diverse mammals feed on similarly diverse fungi. The mammals include insectivores, omnivores, herbivores, and carnivores. The fungi include saprophytes, symbionts, and parasites. The interdependence of animals and fungi has evolved to a high degree in some cases, e.g., the loss of alternative spore dispersal mechanisms by the phylogenetically advanced truffles and other subterranean-fruited fungi (Bessey, 1935; Burnett, 1968; Christenson, 1951; Kavalier, 1965; Gäumann and Dodge, 1928; Ingold, 1953, 1961, 1966; Korf, 1973; Lilly and Burnett, 1951; Rolfe and Rolfe, 1925; Smith, 1938; Trappe, 1971) and the strong reliance of some mammals on fungi as a primary food.

The Sciuridae (chipmunks and squirrels) have been the most observed mammalian mycophagists; consequently, they are also reported to eat the greatest variety of fungi. *Tamiasciurus hudsonicus* (red squirrel) has been credited with eating 89 species of fungi, more than any other mycophagist. Other relatively small mycophagists include members of the Cricetidae (mice, rats, lemmings, voles), Zapodidae (jumping mice), Ochotonidae (pikas), Soricidae (shrews), Didelphidae (opossums), Peramelidae (bandicoots), Phascolomidae (wombats), Macropodidae (rat-kangaroos), Cynopithecinae (baboons), Dasypodidae (armadillos), Leporidae (rabbits and hares), Castoridae (beavers), and Mustelidae (weasels). The reports of fungi in stomachs of shrews may at least in part reflect predation on fungus-feeding invertebrates or other animals rather than mycophagy by the shrews themselves.

The relative importance of fungi in small mammal diets has been estimated as percent of yearly dietary volume for several species (Table 1).

TABLE 1. Yearly dietary volume of fungi (%) for nine small mammal species.

Species	Volume (%)	Source
<i>Eutamias townsendii</i>	72	Tevis, 1953
<i>Eutamias quadrimaculatus</i>	66	Tevis, 1953
<i>Spermophilus lateralis</i>	61	Tevis, 1953
<i>Tamiasciurus douglasii</i>	56	McKeever, 1964
<i>Sciurus griseus</i>	52	Steinecker and Browning, 1970
<i>Eutamias speciosus</i>	32	Tevis, 1953
<i>Eutamias amoenus</i>	27	Tevis, 1953
<i>Clethrionomys glareolus</i>	7	Drożdż, 1966
<i>Apodemus flavicollis</i>	1	Drożdż, 1966

Tevis (1953) suggested a direct correlation between mammal body weight and importance of fungi in the diet among the *Spermophilus* spp. and *Eutamias* spp. that he studied. Other factors of habit and habitat are undoubtedly as much or more causally related to mycophagy. However, a relationship between mammal size and size of fungus eaten does seem likely in the sense that very small fungal species appear to be eaten only by smaller mammals. For instance, sporocarps of many species of the phycomycetous family, Endogonaceae, are often less than a millimeter in diameter and rarely exceed a few millimeters. These species have been reported only in stomach contents of small mammals such as shrews, voles, mice, jumping mice, pikas, etc., but they eat larger fungi as well (Bakerspiegel, 1958; Calhoun, 1941; Hamilton, 1941a; Jameson, 1952; Whitaker, 1962, 1963a; Whitaker and Ferraro, 1963; Whitaker and Maser, 1976).

### *Adaptations for Mycophagy*

Morphological adaptations of small mammals are probably directly related to mycophagy. Animals adapted for digging might be expected to have an advantage in the search for truffles and other hypogeous fungi. Species with weak teeth might be more impelled to eat fungi when available. Unfortunately, the data presently available are too scanty to test these hypotheses.

Mammalian food-gathering behavior has evolved for locating fungi. Tevis (1953) observed innumerable small pits left in the soil from extraction of hypogeous fungi by animals; indeed, mycologists have long used this sign to indicate when and where to seek such fungi. Small mammal mycophagists probably locate hypogeous fungi by odor. Most hypogeous fungi have a noticeable odor at maturity which is almost overwhelming to many humans (Parks, 1919). Often the presence of these fungi is not visible on the soil surface. The ability of squirrels to detect mast by odor has been reported many times (Allen, 1943; Cahalane, 1942; Cram, 1924; Dice, 1927; Ingles, 1947; Ognev, 1940). We have observed over years of extensive collecting of hypogeous fungi that sporocarp odor is not detectable on immature specimens. As some spores reach maturity, a light odor often can be noticed; as the proportion of mature spores increases, so does the intensity of odor. Thus the fungus may remain undetected at early stages of development but emits increasingly strong odors as it matures.

Bright color as well as odor may attract small mammals to epigeous mushrooms and puffballs (Ingles, 1947). Mushrooms are sometimes nibbled *in situ* but are also harvested whole. Numerous observers have reported mushrooms hung in tree branch forks by squirrels, at times the tree "appearing bedecked for Christmas," (Cram, 1924; Hatt, 1929; Krieger, 1967; Marie, 1927; Odell, 1925, 1926; Ognev, 1940; Stakhrovskii, 1932). Since dried fungi preserve indefinitely, the intent of the squirrels is presumably to dry them for caching. Hardy (1949) found a cache of sporocarps left by a *Tamiasciurus hudsonicus* in a hollow tree trunk.

### *Mycophagy and Habitat*

The characteristics of plant communities to which small mammals have become adapted are decidedly related to mycophagy. Forests of ectomycorrhizal trees such as the Pinaceae, Fagaceae, or Betulaceae produce a greater abundance of relatively large, fleshy mushrooms and hypogeous fungi, the typical fruiting forms of ectomycorrhizal fungi (Trappe, 1962). Plant communities of vesicular-arbuscular (VA) mycorrhizal hosts such as the Cupressaceae, Aceraceae, or herbaceous plants of fields, meadows, and bogs form mycorrhizae with certain genera of the Endogonaceae, which fruit hypogeously as single spores or very small, compact sporocarps (Gerdemann and Trappe, 1974). Thus, adjacent habitat types may differ strikingly in form and biomass of sporocarps produced. The small mammals that inhabit ectomycorrhizal forests might be expected to feed more on macrofungi than inhabitants of VA-mycorrhizal meadows. Although adequate data are not available, it can be hypothesized that mycophagy in VA-mycorrhizal redwood or maple forests resembles that of meadows more closely than that of ectomycorrhizal pine or oak forests. If this hypothesis is true, the qualitative and quantitative relationships of small mammal populations between these various kinds of habitats might deserve a new look.

Estimated annual production of epigeous mushrooms is summarized for several forest types in Table 2. The data of Richardson (1970) provide the best of the estimates presented, because his estimates are based on nearly weekly visits to the study site over a period of five years. The other estimates are based on relatively scanty data.

TABLE 2. Estimated production of epigeous mushrooms per ha yr<sup>-1</sup>.

Forest Type	Kg Dry Wt	Kilocalories	Source
Oak and Beech, Hungary	7-160	30,000-658,000*	Hering, 1966
Beech, near Vienna	18-170	77,400-731,000*	Hering, 1966
Beech, Poland	4-5	17,200-21,500	Drożdż, 1965
Oak on slate soil, England	12-95	53,600-407,600	Hering, 1966
Oak on limestone soil, England	3-37	12,900-158,700	Hering, 1966
Conifer, Scotland	16-30	68,600-128,700	Richardson, 1970
Conifer, xeric inland forest, Pacific Northwest, U.S.A.	3	11,200	Smith, C., 1965
Conifer, xeric inland forest, Pacific Northwest, U.S.A.	10-17*	41,200-115,800*	Cooke, 1955
Conifer, Finland	5-19	22,300-82,800	Richardson, 1970
Conifer, Sweden	180	772,200	Richardson, 1970

\* Our calculation, assuming 6.4% dry wt and 4300 calories g<sup>-1</sup> dry wt (cf. Table 3).

Many variables interact to account for the wide range of production estimates; two most obvious are temperature and moisture and their interaction (Smith, 1949). Consequently, each climatic region has a characteristic fruiting pattern (Bohus, 1973; Ceruti, Montacchini and Duployez, 1967; Endo, 1973; Fogel, 1976; Grainger, 1946; Wilkins and Harris, 1946). Production will vary from year to year within a region, depending on weather. Habitats within a region will vary one from another in production, depending on soils, microclimate, and vegetation.

No biomass production studies of VA-mycorrhizal fungi have been reported. However, Kessler and Blank (1972) estimated that nearly 7 million minute sporocarps of Endogonaceae occurred per ha in soil of a Michigan forest dominated by the VA-mycorrhizal *Acer saccharum* Marsh. A substantial biomass may be inferred. Fogel (1976) estimated the production of hypogeous basidiomycete and ascomycete sporocarps in a Western Oregon Douglas-fir stand to range from 11,052 to 16,753 sporocarps ha<sup>-1</sup> and 2.3 to 5.4 kg dry weight ha<sup>-1</sup>.

#### *The Food Value of Fungi*

Fresh, fleshy fungi range from 70 to 94 percent water by weight; large volumes therefore must be eaten to provide adequate protein and phosphorus (Miller and Halls, 1969; Winton and Winton, 1935). The drying of mushrooms by squirrels concentrates their food value. At the same time, fresh fungi might be an important source of water for animals unable to exist on bound or metabolic water when free water is unavailable (Getz, 1968; C. Smith, 1965).

Some measures of the food value of fungi are compared with those of other food sources in Tables 3-5. The data should be interpreted only in a broad sense, since several sources and differing analytical methodologies are represented in the compilation. Fresh, fleshy fungi resemble fruits and vegetables in containing substantially fewer

calories  $\text{gm}^{-1}$  than fresh nut kernels, eggs, or meat (Table 3). On a dry-weight basis, fungi compare favorably with blueberries and mast but contain about a third fewer calories than conifer seed (Table 4). Compared to nut kernels, dried mushrooms are good sources of protein, carbohydrates, and minerals but not fats (Table 5).

The net caloric value of any food to a consumer is the total caloric content digested minus the calories expended in seeking, extracting, ingesting, digesting, and excreting

TABLE 3. Caloric value of fresh foods ( $\text{kcal. gm}^{-1}$  fresh wt.).

Food	Calories	Source
<b>FUNGI</b>		
<i>Agaricus bisporus</i>	0.3-0.5	Singer, 1951; Pilát and Ůsak, 1958
<i>Boletus</i> sp.	0.3	Singer, 1951
<i>Boletus edulis</i>	0.3-0.7	Pilát and Ůsak, 1958; Andreotti and Casoli, 1968
<i>Lentinus edodes</i>	0.3	Singer, 1951
<i>Lactarius deliciosus</i>	0.5	Singer, 1951; Pilát and Ůsak, 1958
<i>Tuber melanosporum</i>	0.4-0.6	Andreotti and Casoli, 1968
"mushrooms"	0.1-0.5	Singer, 1951; Hodgman <i>et al.</i> , 1959; Proudfit and Robinson, 1961
Mean	0.4	
<b>NUTS</b>		
pine kernels	6.3	Hodgman <i>et al.</i> , 1959
Beechnut kernels	6.3-7.0	Proudfit and Robinson, 1961
butternut kernels	7.1-7.4	Hodgman <i>et al.</i> , 1959; Proudfit and Robinson, 1961
chestnut kernels	7.5-7.6	Proudfit and Robinson, 1961
hazelnut kernels	6.7	Proudfit and Robinson, 1961
hickory nut kernels	1.9	Hodgman <i>et al.</i> , 1959; Proudfit and Robinson, 1961
pecan kernels	6.6	Hodgman <i>et al.</i> , 1959; Proudfit and Robinson, 1961
walnut kernels	6.7-7.0	Proudfit and Robinson, 1961
Mean	6.2	
<b>FRUITS</b>		
apples	0.6	Proudfit and Robinson, 1961
blackberries	0.6	Proudfit and Robinson, 1961
blueberries	0.7	Proudfit and Robinson, 1961
cherries	0.7	Proudfit and Robinson, 1961
currants	0.6	Proudfit and Robinson, 1961
plums	0.6	Proudfit and Robinson, 1961
strawberries	0.4	Hodgman <i>et al.</i> , 1959; Proudfit and Robinson, 1961
Mean	0.6	
<b>VEGETABLES</b>		
lettuce	0.2	Hodgman <i>et al.</i> , 1959; Proudfit and Robinson, 1961
onions	0.5	Proudfit and Robinson, 1961
water cress	0.2	Proudfit and Robinson, 1961
Mean	0.3	
<b>EGGS and MEAT</b>		
eggs	2.5	Proudfit and Robinson, 1961
beef	1.8-3.4	Hodgman <i>et al.</i> , 1959; Proudfit and Robinson, 1961
Mean	2.1	

the nondigested residual of that food. The relative value of fungal caloric sources for any mycophagist must be compared to other sources in terms of an energy budget. If less energy is expended per calorie gained in seeking and eating fungi than in seeds or insects, the relative value of fungi as an energy source is increased.

TABLE 4. Caloric value of dried plant parts (kcal. gm<sup>-1</sup> dry wt.).

Food	Calories	Source
FUNGI		
<i>Peridermium barknessii</i> in <i>P. contorta</i> bark	5.0	Smith, C. 1965
<i>Rhizopogon</i> sp.	4.9	Smith, C. 1965
<i>Russula decolorans</i>		
gills and spores	4.8	Smith, C. 1965
pileal context	4.0	
stipe	4.2	
<i>Suillus tomentosus</i>		Smith, C. 1965
tubes and spores	5.2	
pileal context	4.2	
stipe	4.1	
Mean	4.5	
CONIFER SEEDS		
<i>Abies amabilis</i>	6.8	Smith, C. 1965
<i>Abies lasiocarpa</i>	7.1	Smith, C. 1965
<i>Picea engelmannii</i>	7.1	Smith, C. 1965
<i>Picea excelsa</i>	6.7	Danilov, 1938
<i>Pinus contorta</i>	6.8	Smith, C. 1965
<i>Pinus silvestris</i>	6.2	Danilov, 1938
<i>Pinus monticola</i>	7.4	Smith, C. 1965
<i>Pinus ponderosa</i>	7.6	Smith, C. 1965
<i>Pinus pumila</i>	6.9	Danilov, 1938
<i>Pseudotsuga menziesii</i>	7.1	Smith, C. 1965
<i>Tsuga heterophylla</i>	7.1	Smith, C. 1965
Mean	7.0	
MAST		
<i>Acer saccharinum</i>	4.2	Smith, C., 1970
<i>Carya ovata</i>	7.2	Smith, C., 1970
<i>Juglans nigra</i>	6.3	Smith, C., 1970
<i>Quercus alba</i>	4.0	Smith, C., 1970
<i>Quercus macrocarpa</i>	4.2	Smith, C., 1970
<i>Quercus shumardii</i>	5.3	Smith, C., 1970
Mean	5.2	
FRUIT		
<i>Vaccinium deliciosum</i>	5.3	Smith, C. 1965
BUDS		
<i>Picea excelsa</i>	4.0	Danilov, 1938

Unfortunately, the relative digestibility of different food sources is not known for different mammals. The large gastric caecum of many rodents has been suggested as a site of enzyme systems capable of extracting energy from  $\beta$ -linked carbohydrates of fungal cell walls (C. Smith, 1965). Our impression from examining stomach and fecal contents of several hundred small mammal mycophagists is that fungal cell cytoplasm is readily digested, cell walls are sometimes digested, and spores are not digested at

TABLE 5. Chemical composition of dried plant parts and meat.

Plant Species	% Dry Weight			Ash	Source
	Protein	Fat	Carbo- hydrate		
<b>FUNGI</b>					
<i>Agaricus bisporus</i>	48-62	1-2	—	7	Singer, 1961
<i>Amillaria mellea</i>	16	—	76	8	Singer, 1961
<i>Boletus edulis</i>	32-35	5	58-59	6-8	Singer, 1961
<i>Clavaria flava</i>	19	2	47	5	Winton and Winton, 1935
<i>Hygrophorus, Amillaria, Russula, Clitocybe, Amanita, Tricholoma, Pholiota</i> sp.	12-27	—	—	—	Miller and Halls, 1969
<i>Hygrophorus, Amanita</i> sp.	23	—	—	—	Miller and Halls, 1969
<i>Lactarius deliciosus</i>	27	7	28	6	Singer, 1961
<i>Lentinus edodes</i>	19	5	54	3	Singer, 1961
<i>Marasmius oreades</i>	35-43	2-4	34	10	Singer, 1961
<i>Morchella esculenta</i>	34-35	2	46-47	9-11	Singer, 1961
<i>Morchella</i> sp.	30	—	—	—	Mendel, 1898
<i>Morchella, Rhizopogon, Cortinarius, Lactarius, Amanita, Clitocybe</i> sp.	30	—	—	—	Miller and Halls, 1969
<i>Pleurotus cretaciuss</i>	21	—	—	—	Winton and Winton, 1935
<i>Saccharomyces cerevisiae</i>	12	<1	21	—	Hodgman, <i>et al.</i> , 1963
<i>Suillus granulatus</i>	14	2	70	6	Singer, 1961
<i>Suillus grevillei</i>	21	2	64	6	Singer, 1961
<i>Suillus luteus</i>	20	4	53	6	Singer, 1961
<i>Tricholoma favovirens</i>	15-18	—	71-78	7-11	Singer, 1961
<i>Tuber melanosporum</i>	25-26	2	39-46	8	Andreotti and Casoli, 1968
<b>NUTS</b>					
Chestnut kernels	11	7	72	2	Hodgman, <i>et al.</i> , 1963 Chapman and Baumgartner, 1939
Butternut kernels	28	61	3	3	Winton and Winton, 1932
Black walnut kernels	28-30	56-58	6	2	Winton and Winton, 1932 Chapman and Baumgartner, 1939
Pecan kernels	10-11	70-75	10	2	Winton and Winton, 1932
Hickory kernels	13-20	65-70	6-9	2	Winton and Winton, 1932
Filbert kernels	16-17	63-65	10-13	2	Winton and Winton, 1932
Beechnut kernels	22	42-57	19	3-4	Winton and Winton, 1932 Chapman and Baumgartner, 1939
<b>MEAT</b>					
Chipped beef	30	6	<1	—	Hodgman, <i>et al.</i> , 1963

all. Spores presumably contain concentrated energy sources, as indicated by the relatively high caloric values of gills or tubes of mushrooms as compared to the somatic parts (Table 4) and by the high lipid content seen in spores of some fungi (Gerdemann and Trappe, 1974). However, reports of squirrels eating caps and gills of mushrooms while leaving the stems (Hatt, 1929; Ognev, 1940; C. Smith, 1965) probably reflect palatability rather than selection for food value.

Aside from their caloric value, some of the diverse organic compounds of fungi may be important in small mammal physiology. For example, ergosterol, a possibly important source of mammalian hormone precursors, occurs widely in the higher fungi

and ranges from about 0.2 to 0.5 percent of the dry weight of fleshy fungi (Milazzo, 1965; Shivrina *et al.*, 1968; Catalfomo and Trappe, 1970). Fungi commonly contain other steroids, triterpenes, amines, indoles, and phenols (Catalfomo and Trappe, 1970) of unknown but potential use to mammals. Vitamins such as biotin, niacin, pantothenic acid, and riboflavin are produced in significant amounts by some fleshy fungi (Shemakhanova, 1967). On a fresh weight basis, mushrooms have higher concentrations of niacin and riboflavin than most nuts, fruits, and leaf vegetables and equal or excel most meats in this regard (Proudfit and Robinson, 1950).

Fungal tissues effectively accumulate nonmetallic and metallic elements. Stark (1972) found that sporocarps of five fungal species contained substantially higher concentrations of Cu, N, P, and Zn than did pine needles (dry wt. basis). At least some of the five also contained higher concentrations of Ca, Fe, K, Mg, and Na. Fungal rhizomorphs were also high in concentrations of these elements, e.g., 180 times as much Fe as pine needles.

The overall quality of fungi as a food source for small mammals has not been experimentally assessed. Tevis (1952) reported that *Eutamias* stomachs containing only fungi were always heavier than those of animals feeding on non-fungal material. He also found that when chipmunks began to acquire hibernation fat, those eating fungi became fat sooner. But Naumov's report of squirrels starving to death while gorged with *Polyporus betulinus* Fr. suggests that some fungi may have very little food value (Ognev, 1940), at least for some mammals.

Cyclic over-population of mammals and poor seed crops probably periodically enhance the importance of mycophagy in reducing food stress (Lampio, 1967; Ognev, 1940; Rajala and Lampio, 1963). The effect of poor mast crops, traditionally considered a key factor in regulating squirrel populations, might be moderated in good mushroom years (Baumgartner, 1939; Brown and Yeager, 1945; Stienecker and Browning, 1970; Uhlig, 1955; and others). Seasonal buildup of rodent populations, together with the seasonality of fungal fruiting, might explain the increased seasonal use of *Endogone* by *Peromyscus maniculatus*, *Microtus longicaudus*, *Clethrionomys gapperi*, *Phenacomys intermedius*, and *Zapus princeps* (Williams and Finney, 1964).

#### *Fungal Toxins and Mycophagy*

Feeding trials and field observations of *Amanita muscaria*, *Amanita phalloides*, *Lactarius rufus*, and *Lactarius torminosus* consumption by squirrels indicate that they and presumably other rodents can safely eat mushrooms considered poisonous to man (Ballou, 1927; Cram, 1924; Hastings and Mottram, 1916; Hatt, 1929; Klugh, 1927; Metcalf, 1925). With this evidence in mind, Walton's (1898) observation, "For thirteen years I have made use of the varieties of mushrooms selected by squirrels and wood mice as edible. . .," should be taken skeptically. He was lucky to have escaped poisoning.

We have found no reports on poisoning of small mammals by fungi in the wild. One prime reason may be that no one has seriously studied the matter. Several hypotheses might be proposed, among them that (A) no fungi are poisonous to small mammals; (B) mammals can detect toxins by taste, smell, or in some other way; or (C) mammals are poisoned by some fungi, but when poisoning occurs, they seclude themselves before symptoms become evident to observers.



Hypothesis A was supported by Ford (1908, 1910), who suggested that herbivore alimentary canals can neutralize poisons which are highly toxic if administered by subcutaneous or interperitoneal inoculation. However, chickens and a cow are reported to have died after eating sporocarps of *Amanita* sp. (Ballou, 1927). Recently it has been demonstrated that a number of toxins each with different properties are responsible for mushroom poisoning symptoms (Litten, 1975). The *Amanita* toxin phalloidin, for instance, is not toxic when given orally, but mice given *Amanita* whole toxic extract containing myriaphalloisins die in a few hours to four days.

Hypothesis B seems a reasonable possibility, considering the selective pressure against genetic lines given to feasting on poisonous fungi. Hypothesis C also seems reasonable, especially since most mycotoxins require some incubation time before symptoms are expressed: mushroom poisoning of humans often does not arouse symptoms for several hours after ingestion. Hypothesis A has been proven wrong for a limited number of mammals; B and C remain to be tested.

#### *Mycophagists as Disseminators of Fungi*

Mycophagy serves in fungal dissemination by physical transport of spores and hypothetically, at least in some cases, by breaking spore dormancy. Most epigeous fungi, even the coprophilous species, are adapted for aerial or water transport of spores; mycophagy would seem incidental to primary dispersal mechanisms (Ingold, 1953). Of course, mushrooms hung in trees by squirrels are exposed to air currents that could carry spores farther than those at the ground surface. Mycorrhizal species or root pathogens may gain some strategic advantage over aerial dissemination if spores are eaten and then excreted near susceptible host roots by burrowing mycophagists.

Hypogeous fungi, on the other hand, are clearly dependent on mycophagy for physical transport. They fruit underground, their spore-bearing tissues are typically enclosed in a persistent peridium, and for the most part they do not forcibly discharge their spores. Even those which can forcibly discharge spores, such as the truffle relative, *Geopora cooperi* Harkn. (Burdall, 1965), must be dug up and opened above ground by an animal to release spores to air currents (Korf, 1973).

The phycomycetous fungi of the family Endogonaceae deserve special mention because they have been found so often in small rodent stomachs. The various species may be epigeous or hypogeous and may form sporocarps or be borne as individual spores in soil. At present, only the sporocarp-forming species have been found in rodent stomachs. Whether epigeous or hypogeous, the sporocarps are typically compact clusters of very large spores devoid of discharge mechanisms. Presumably, these sporocarps are strongly dependent on mycophagy for transport. Spores borne singly in soil, on the other hand, may be eaten by insects or arthropods but also can be transported by any of the forces that move soil (Gerdemann and Trappe, 1974).

The effects of passage through a digestive tract on spore germination are unknown for most fungi. In general, spores of coprophilous and xylophilous fungi germinate poorly or not at all under any circumstances (Fries, 1943; Benedict *et al.*, 1967). The relatively few reported successes in inducing germination of spores of ectomycorrhizal fungi entailed treatments such as low temperature, alternate wetting and drying, soil extracts, co-culture with other organisms, slug digestive enzymes, or exposure to living roots (Sussman and Halversen, 1966; Bowen and Theodorou, 1974; Lamb and Richards,

1974). Of the Endogonaceae, chlamydosporic species which form vesicular-arbuscular mycorrhizae germinate readily without special treatment; one species has been shown to be viable after passage through a *Microtus* digestive tract (Trappe and Maser, 1976). No one has yet succeeded in germinating spores of those that occur only in sporocarps, at least some of which are ectomycorrhizal species (Gerdemann and Trappe, 1974).

Mycophagy would subject ingested fungal spores to body heat, enzymatic action, and co-culture with the myriad microorganisms of digestive tracts and feces. The process has not been demonstrated to affect spore dormancy as yet. Indeed root exudates appear to induce germination of some hypogeous mycorrhizal fungi without mycophagy (Palenzona, 1969; Bowen and Theodorou, 1974). Since essentially no research has been reported on the physiological effects of mycophagy on spores, the question remains moot.

#### *Mycophagy in Ecosystem Analysis*


Invertebrates and vertebrates that feed wholly or in part on fungi have been arbitrarily categorized either as herbivores or fungal decomposers. No generalization fits comfortably, however, because the fungi may be saprophytes, biotrophic or necrotrophic symbionts, or predators (Harley, 1971; Lewis, 1973). Although necrotrophic symbionts may be considered "living tissue saprophytes" for purposes of ecosystem modeling, the biotrophic symbionts such as mycorrhizal fungi pose a special problem: the structural and physiological interdependence of fungus and host preclude their conceptual separation either physically or functionally.

One solution to this problem is to consider mycorrhizal fungi as extensions of the host root system, since the fungi are essential to nutrient absorption by the host and the fungi obtain their energy from the host much as if they were indeed root tissue. Thus the mycorrhizal fungi can be regarded as organs of producer plants. Mycophagists that feed on mycorrhizal fungi are then grazers. Grazing of mycorrhizal fungi accounts for about 75 percent of the small mammal mycophagy in terms of the reports listed later in this paper, evaluated as known or presumed mycorrhizal taxa (Trappe, 1962, 1971). If only reports of stomach contents are considered, the percentage is considerably higher. Mycophagy of saprophytes, necrotrophic symbionts, or predators, on the other hand, can be regarded as part of the decomposer process and accounts for about 25 percent of the reports. Many, perhaps most mycophagists feed on both categories of fungi and thereby participate in both processes.

Small mammal mycophagy seems surely to be a significant energy transfer process in ecosystems, but it remains to be quantitatively assessed. We are aware of only one paper with enough information to present a small mammal-fungus carbon budget for a short period: C. Smith (1965) estimated the caloric consumption of different food sources by the red squirrel (*Tamiasciurus hudsonicus*) for the month of August in western Washington. During this time, mycophagy accounted for about 73 percent of the squirrels' caloric intake, even though coniferous seeds and blueberries were also available (Fig. 1). Data from several other papers were not usable since percent volume and frequency values reported are useless without estimates of food consumption, weight, caloric value, stomach size, food density, etc. Modeling can not proceed until integrated studies on mycophagist food habits and fungal production are completed.

Despite inadequate information, mycophagy is clearly important to mammals and fungi alike. More extensive and definitive information is badly needed, particularly for understanding the dynamics of small mammal populations and for ecosystem modeling; nutrient flow is not a simple matter of diffusion through arbitrarily generalized functional groups such as primary producers, consumers, or decomposers.

Figure 1. *Tamiasciurus hudsonicus* caloric (kcal. 1000 gm<sup>-1</sup>) consumption for August. (Smith, C., 1965) Percent of August consumption enclosed in parentheses.



### The Reciprocal Lists

The literature reports of small mammal mycophagy are compiled in the following lists by taxa of both fungi and mammals. The lists are intended to provide a broad overview indexed to the original sources. Because any such listing is subject to limitations, restraint should be used in drawing detailed inferences from the lists themselves. Use them only as portals to the original literature. Keep in mind that the lists exclude lichens; that while coverage is almost complete for the North American literature, several references from other parts of the world could not be located; and that some taxa undoubtedly have been misidentified by some original authors.

No single system of taxonomy enjoys complete acceptance for either the fungi or the mammals. We have accordingly followed the classification and nomenclature of the more comprehensive compendia where possible, recognizing that many readers may prefer a somewhat different system. The fungal classification has been drawn from various sources, including Singer (1962) for Agaricales, Overholts (1953) for Polyporaceae, and Smith (1951) for Gasteromycetes. Mammal classification follows Hall and Kelson (1959), Jackson (1928), Johnson (1968), and Johnson and Ostenson (1959) for North American species and Ognev (1940, 1950) for Eurasian species.

The lists are alphabetical except for classes and orders. All known synonyms of a taxon have been cross-indexed to a single name; we do not intend any novel taxonomic decisions in this process, since the conclusions were drawn entirely from the literature, not from personal examination of types or other specimens.

Fungi listed with asterisk have been reported as demonstrated or probable mycorrhizal species in Trappe (1962, 1971), Gerdemann and Trappe (1974), or elsewhere.

### Fungi and the Animals That Eat Them

#### CLASS PHYCOMYCETES

Unspecified taxa: *Potorous tridactylus apicalis* (Guiler, 1971).

#### ORDER OOMYCETALES

See Class Phycomycetes. Apparently this order has never been used.

#### ORDER MUCORALES

#### FAMILY ENDOGONACEAE

- Endogone* spp.: *Blarina brevicauda* (Hamilton, 1941b; Linzey and Linzey, 1973; Whitaker, 1962; Whitaker and Ferraro, 1963); *Clethrionomys gapperi gapperi* (Linzey and Linzey, 1973; Whitaker, 1962; Williams and Finney, 1964); Cricetidae (Dowding, 1959). *Eutamias* spp. Dowding, 1955, 1959; Whitaker, 1962); *Lagurus curtatus* (Whitaker, 1962); *Microtus chrotorrhinus* (Linzey and Linzey, 1973); *M. longicaudus* (Williams and Finney, 1964; *M. ochrogaster* (Zimmermann, 1965, 1966); *M. oeconomus macfarlandi* (Dowding, 1955; Whitaker, 1962); *M. pennsylvanicus* (Whitaker, 1962; Zimmermann, 1966); *M. pinetorum* (Linzey and Linzey, 1973; Whitaker, 1962); *Mus musculus* (Calhoun, 1941; Whitaker, 1962, 1966); *Napaeozapus insignis insignis* (Linzey and Linzey, 1973; Whitaker, 1962, 1963a); *Ochotona* spp. (Dowding, 1959); *O. princeps* (Whitaker, 1962; Zimmermann, 1965); *Onychomys leucogaster* (Whitaker, 1962); *Oryzomys palustris* (Negus, Gould, and Chipman, 1961); *Peromyscus gossypinus* (Calhoun, 1941; Whitaker, 1962); *P. leucopus* (Calhoun, 1941; Linzey and Linzey, 1973; Whitaker, 1962, 1963b, 1966); *P. leucopus noveboracensis* (Hamilton, 1941b); *P. maniculatus* (Dowding, 1955; Linzey and Linzey, 1973; Whitaker, 1962; Williams and Finney, 1964); *P. maniculatus bairdii* (Whitaker, 1966); *P. maniculatus gracilis* (Hamilton, 1941b); *Phenacomys intermedius* (Williams and Finney, 1964), Rodentia (Ingold, 1961, 1966; Kessler and Blank, 1972); *Sorex* spp. (Thaxter, 1922); *S. bendirii* (Whitaker and Maser, 1976); *S. cinereus* (Whitaker, 1962); *S. fumeus* (Linzey and Linzey, 1973; Whitaker, 1962); *S. trowbridgii* (Whitaker and Maser, 1976); *S. vagrans* (Whitaker and Maser, 1976); *S. vagrans pacificus* (Whitaker and Maser, 1976); *S. vagrans yaquinae* (Whitaker and Maser, 1976); *Soricidae* (Dowding, 1959); *Synaptomys cooperi cooperi* (Conner, 1960; Hamilton, 1941a, 1942; Hatt, 1930; Linzey and Linzey, 1973; Whitaker, 1962); *S. cooperi gossii* (Burt, 1928; Hamilton, 1941b); *Talpidae* (Dowding, 1959); *Zapus hudsonicus* (Whitaker, 1962); *Z. princeps* (Williams and Finney, 1964); *Z. trinotatus* (Whitaker, 1962).
- \**E. flammicorona* Trappe and Gerdemann: *Clethrionomys californicus californicus* (Gerdemann and Trappe, 1974); *Peromyscus maniculatus* (Gerdemann and Trappe, 1974).
- \**E. lactiflua* Berk. and Broome: *Clethrionomys californicus californicus* (Gerdemann and Trappe, 1974); *Peromyscus maniculatus* (Gerdemann and Trappe, 1974); *Zapus trinotatus* (Gerdemann and Trappe, 1974).
- E. pisiformis* Link ex Fries: *Clethrionomys californicus californicus* (Gerdemann and Trappe, 1974); *Peromyscus maniculatus* (Gerdemann and Trappe, 1974).
- E. macrospora*: See *G. macrocarpa*. (*E. macrospora* was an erroneous citation for *E. macrocarpa*).
- \**Glomus caledoniensis* (Nicol. and Gerd.) Trappe and Gerdemann: *Peromyscus maniculatus* (Gerdemann and Trappe, 1974).
- \**G. fasciculatus* (Thaxt. sensu Gerdemann) Gerdemann and Trappe: *Clethrionomys californicus californicus* (Gerdemann and Trappe, 1974); *Lagurus curtatus* (Dowding, 1955); *Microtus pennsylvanicus* (Bakerspigel, 1956, 1958; Whitaker, 1962); *Ochotona princeps* (Dowding, 1955); *Onychomys leucogaster* (Dowding, 1955); *Peromyscus maniculatus* (Bakerspigel, 1956, 1958; Dowding, 1955; Whitaker, 1962); *Sciurus aberti* (Stephenson, 1975); *Zapus princeps* (Bakerspigel, 1956; Whitaker, 1962); *Z. trinotatus* (Gerdemann and Trappe, 1974).
- \**G. macrocarpus* Tul. and Tul.: *Blarina* spp. (Hamilton, 1941b); *B. brevicauda* (Diehl, 1939); *Clethrionomys* spp. (Hamilton, 1941b); *C. gapperi gapperi* (Diehl, 1939); *Peromyscus* spp. (Hamilton, 1941b); *P. leucopus noveboracensis* (Diehl, 1939); *P. maniculatus* (Harling and McClaren, 1970); *Sorex cinereus* (Diehl, 1939; Hamilton, 1941b); *S. fumeus* (Diehl, 1939; Hamilton, 1941b); *Synaptomys cooperi cooperi* (Diehl, 1939); *S. cooperi gossii* (Hamilton, 1941b).
- \**G. macrocarpus* Tul. and Tul. var. \**macrocarpus*: *Clethrionomys californicus californicus* (Gerdemann and Trappe, 1974); *Peromyscus maniculatus* (Gerdemann and Trappe, 1974).
- \**G. microcarpus* Tul. and Tul.: *Clethrionomys californicus californicus* (Gerdemann and Trappe, 1974); *Peromyscus maniculatus* (Gerdemann and Trappe, 1974).
- \**G. monosporus* Gerdemann and Trappe: *Clethrionomys californicus californicus* (Gerdemann and Trappe, 1974); *Zapus trinotatus* (Gerdemann and Trappe, 1974).
- G. pulvinatus* (Henn.) Trappe and Gerdemann: *Zapus princeps* (Dowding, 1955; Whitaker, 1962).
- \**Sclerocystis rubiformis* Gerdemann and Trappe: *Clethrionomys californicus californicus* (Gerdemann and Trappe, 1974); *Peromyscus maniculatus* (Gerdemann and Trappe, 1974); *Zapus trinotatus* (Gerdemann and Trappe, 1974).

CLASS ASCOMYCETES

ORDER PLECTASCALES

FAMILY ELAPHOMYCETACEAE

*Elaphomyces* spp.: *Neotoma fuscipes annectens* (Parks, 1922); Rodentia (Ingold, 1961, 1966; Reess and Fisch, 1887); *Sciurus aberti* (Stephenson, 1975).

\**E. granulatus* Fr.: Leporidae (Hawker, 1954); Rodentia (Hastings and Mottram, 1916; Hawker, 1954; Ingold, 1953); Sciuridae (Bulle, 1922; Dowding, 1959); *Sciurus carolinensis carolinensis* (Ingold, 1973); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).

ORDER DOTHIIDIALES

FAMILY DOTHIIDACEAE

*Dibotryon* spp.: *Sciurus carolinensis carolinensis* (Dudderar, 1967).

ORDER HYPOCREALES

FAMILY HYPOCREACEAE

*Hypomyces* spp.: *Tamiasciurus hudsonicus hudsonicus* (Seton, 1928).

ORDER PEZIZALES

FAMILY HELVELLACEAE

*Helvella* spp.: *Oryctolagus cuniculus* (Kumerlove, 1956).

*H. lacunosa* Afz. ex Fr.: *Neotoma fuscipes* (Linsdale and Tevis, 1951).

*H. mitra*: See *H. lacunosa*.

*Paxina acetabulum* (L. ex St. Amans) Kuntze: *Neotoma fuscipes* (Linsdale and Tevis, 1951).

FAMILY MORCHELLACEAE

*Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).

*Morchella* spp.: *Lepus europaeus* (Kumerlove, 1956).

FAMILY PEZIZACEAE

\**Geopora cooperi* Harkness: *Sciurus aberti aberti* (Stephenson, 1975).

*Peziza* spp.: *Neotoma fuscipes* (Linsdale and Tevis, 1951); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).

*P. emileia* Cke: *Neotoma fuscipes* (Linsdale and Tevis, 1951).

ORDER TUBERALES

*Peromyscus boylei* (Jameson, 1952); *Spermophilus* spp. (Bessey, 1935); *Tamiasciurus douglasi* (Smith, C., 1965); *T. hudsonicus hudsonicus* (Smith, C., 1965).

FAMILY TUBERACEAE

Leporidae (Kavaler, 1965); *Neotoma* spp. (Smith, 1938); Rodentia (Christenson, 1951; Kavaler, 1965; Lilly and Barnett, 1951; Mehrotra, 1967; Smith, 1938); Sciuridae (Kavaler, 1965).

*Balsamia* spp.: *Clethrionomys glareolus* (Drozd, 1966).

\**B. vulgaris* Vitt.: *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916); *S. vulgaris* (DuReau, 1912; Hatt, 1929).

\**Fischerula subcaulis* Trappe: *Microtus oregoni* (Trappe, 1975).

*Genea* spp.: *Clethrionomys glareolus* (Drozd, 1966).

*Hydnotrya* spp.: *Clethrionomys glareolus* (Drozd, 1966).

*Tuber* spp.: *Clethrionomys glareolus* (Drozd, 1966); Geomyidae (Parks, 1922); *Neotoma fuscipes* (Linsdale and Tevis, 1951); Rodentia (Ingold, 1961, 1966); *Sciurus aberti aberti* (Keith, 1965; Stephenson, 1975).

\**T. aestivum* Vitt.: Rodentia (Hawker, 1954).

*T. candidum* Harkn: see *T. rufum* var. *nitidum*.

\**Tuber rufum* var. *nitidum* (Vitt.) Fischer: *Neotoma* spp. (Parks, 1921); Rodentia (Parks, 1921).

CLASS BASIDIOMYCETES

ORDER UREDINALES

*Peromyscus leucopus* (Calhoun, 1941).

FAMILY PUCCINIACEAE

*Peridermium harknessii* Moore: *Tamiasciurus hudsonicus hudsonicus* (Smith, C. 1965).

FAMILY MELAMPORACEAE

*Cronartium ribicola* A. Fisch.: Sciuridae (Spaulding, 1922); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).

ORDER TREMALES

FAMILY DACRYMYCETACEAE

*Calocera viscosa* (Fr.) Fr.: *Sciurus vulgaris* (Ognev, 1940; Stakhrovskii, 1932).

ORDER AGARICALES

*Sciurus vulgaris leucourus* (Barrett-Hamilton and Hinton, 1921).

FAMILY AGARICACEAE

*Neotoma cinerea orolestes* (Finley, 1958); *N. floridana magister* (Newcombe, 1930); *N. fuscipes annectens* (Parks, 1922); *Sciurus carolinensis carolinensis* (Styan, 1946); *S. niger niger* (Baumgartner, 1939a, 1939b).

*Agaricus* spp.: *Neotoma floridana magister* (Newcombe, 1930); *Sciurus aberti aberti* (Keith, 1965).

*A. campestris* L. ex Fr.: *Potorous tridactylus apicalis* (Guiler, 1971); *Sciurus vulgaris* (Ognev, 1940; Stakhrovskii, 1932); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).

*Coprinus* spp.: *Blarina brevicauda* (Hamilton, 1930); *Didelphis virginiana* (Hamilton, 1958).

*Lepiota* spp.: *Sciurus aberti aberti* (Keith, 1965).

*L. naucinoides*: See *Leucoagaricus naucinus*.

- Leucoagaricus naucinus* (Fr.) Sing.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929; Klugh, 1927).
- Mycena epipterygia* (Scott. ex Fr.) S. F. Gray: *Leporidae* (Hastings and Mottram, 1916).
- Psalliota campestris*: See *Agaricus campestris*.
- FAMILY AMANITACEAE
- Amanita* spp.: Mammalia (Miller and Halls, 1969); Sciuridae (Miller and Halls, 1969); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1920, 1922; Hamilton, 1939, Smith, 1968); *T. hudsonicus minnesota* (Hamilton, 1943).
- \**A. muscaria* (L. ex Fr.) Pers. ex Hooker: Rodentia (Hastings and Mottram, 1916); Sciuridae (Hastings and Mottram, 1916); *Sciurus aberti aberti* (Keith, 1965); *S. carolinensis carolinensis* (Hastings and Mottram, 1916); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929; Klugh, 1927; Krieger, 1967; Metcalf, 1925; Rue, 1967).
- \**A. phalloides* (Vaill. ex Fr.) Secr.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).
- \**A. rubescens* (Pers. ex Fr.) S. F. Gray: *Leporidae* (Hastings and Mottram, 1916).
- \**A. vaginata* (Bull. ex Fr.) Vitt.: *Sciurus aberti aberti* (Keith, 1965); *Tamias striatus lysteri* (Allen, 1938, Fraleigh, 1929); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).
- Amanitopsis vaginata variabilis*: See *Amanita vaginata*.
- Pluteus atricapillus* (Secr.) Sing.: *Neotoma fuscipes* (Linsdale and Tevis, 1951); *Sciurus carolinensis carolinensis* (Nixon, Worley, and McClain, 1968); *S. niger niger* (Nixon, Worley, and McClain, 1968).
- P. cervinus*: See *P. atricapillus*.
- P. nanus* (Pers. ex Fr.) Kumm.: *Sciurus vulgaris exalbidus* (Ognev, 1940).
- FAMILY BOLETACEAE
- Neotoma fuscipes annectens* (Parks, 1922); Rodentia (Miller and Halls, 1969); Sciuridae (Miller and Halls, 1969).
- Boletinus cavipes*: See *Suillus cavipes*.
- \**B. pictus* (Peck) Peck: *Tamiasciurus hudsonicus hudsonicus* (Hardy, 1949).
- Boletus* spp.: *Eutamias quadrimaculatus* (Tevis, 1952); *Lemmus lemmus* (Ognev, 1948); *Neotoma floridana magister* (Newcombe, 1930; Poole, 1940); Sciuridae (Odell, 1926); *Sciurus aberti aberti* (Keith, 1965); *S. carolinensis carolinensis* (Cross, 1942; Nixon, Worley, and McClain, 1968); *S. niger niger* (Nixon, Worley, and McClain, 1968); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922; Klugh, 1927; Krieger, 1967; Seton, 1929; Smith, 1968).
- B. badius*: See *Xerocomus badius*.
- B. bovinus*: See *Suillus bovinus*.
- \**B. calopus* Fr.: Sciuridae (Cooke, 1890); *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916).
- B. castaneus*: See *Gyroporus castaneus*.
- B. chrysenteron*: See *Xerocomus chrysenteron*.
- \**B. edulis* Bull. ex Fr.: Rodentia (Hastings and Mottram, 1916); *Sciurus vulgaris* (Ognev, 1940).
- B. flavidus*: See *Suillus flavidus*.
- \**B. impolitus* Fr.: Sciuridae (Cooke, 1890); *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916).
- B. luteus*: See *Suillus luteus*.
- \**B. mirabilis* Murr.: *Tamiasciurus douglasi douglasi* (Smith, C., 1965); *T. hudsonicus hudsonicus* (Smith, C., 1965).
- B. pachypus*: See *B. calopus*.
- B. pallidus* Frost: *Neotoma floridana magister* (Poole, 1940).
- B. piperatus*: See *Suillus piperatus*.
- B. rufus*: See *Leccinum aurantiacum*.
- B. scaber*: See *Leccinum scabrum*.
- B. subtomentosus*: See *Xerocomus subtomentosus*.
- B. variegatus*: See *Suillus variegatus*.
- B. versipellis* Fr.: See *Leccinum versipellis*.
- \**Gyroporus castaneus* (Bull. ex Fr.) Quel.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).
- \**Leccinum aurantiacum* (Bull.) S. F. Gray: *Sciurus vulgaris* (Ognev, 1940); *S. vulgaris exalbidus* (Ognev, 1940; Stakhrovskii, 1932).
- \**L. scabrum* (Fr.) S. F. Gray: Sciuridae (Masse, 1913); *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916); *S. vulgaris* (Ognev, 1940; Stakhrovskii, 1932); *S. vulgaris exalbidus* (Ognev, 1940); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1920, 1922; Hatt, 1929).
- \**L. versipellis* (Fr.) Snell: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).
- \**Suillus bovinus* (L. ex Fr.) Kuntze: Rodentia (Hastings and Mottram, 1916); *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916); *S. vulgaris exalbidus* (Ognev, 1940); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).
- \**S. cavipes* (Opat.) A. H. Sm. and Thiers: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).
- \**S. flavidus* (Fr.) Sing.: Rodentia (Hastings and Mottram, 1916).
- \**S. granulatus* (Fr.) Kuntze: *Tamiasciurus douglasi douglasi* (Smith, C., 1965); *T. hudsonicus hudsonicus* (Smith, C., 1965).
- \**S. lakei* (Murr.) A. H. Sm. and Thiers: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).
- \**S. luteus* (L. ex Fr.) S. F. Gray: *Sciurus vulgaris* (Ognev, 1940; Stakhrovskii, 1932); *S. vulgaris exalbidus* (Ognev, 1940); *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).

- \**S. piperatus* (Bull. ex Fr.) Kuntze: *Sciurus vulgaris exalbidus* (Ognev, 1940).
- \**S. subolivaceus* A. H. Sm. and Thiers: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).
- \**S. tomentosus* (Kauffm.) Sing., Snell, and Dick: *Tamiasciurus douglasi douglasi* (Smith, C., 1965); *T. hudsonicus hudsonicus* (Smith, C., 1965).
- \**S. variegatus* (Sw. ex Fr.) Kuntze: Rodentia (Hastings and Mottram, 1916); *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).
- \**Xerocomus badius* (Fr.) Kühner ex Gilbert: Leporidae (Buller, 1922); Rodentia (Hastings and Mottram, 1916).
- \**X. chryseron* (Bull. ex St. Amans) Quél.: Rodentia (Hastings and Mottram, 1916); *Sciurus vulgaris exalbidus* (Ognev, 1940).
- \**X. submentosus* (L. ex Fr.) Quél.: *Sciurus vulgaris* (Ognev, 1940; Stakhrovskii, 1932); *S. vulgaris exalbidus* (Ognev, 1940).

#### FAMILY COPRINACEAE

- Hypholoma* spp.: See *Psathyrella* spp.
- H. fasciculare*: See *Naematoloma fasciculare* in Strophariaceae.
- Psathyrella* spp.: *Sciurus aberti aberti* (Keith, 1965); *Tamiasciurus hudsonicus hudsonicus* (Seton, 1929).

#### FAMILY CORTINARIACEAE

- Cortinarius* spp.: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1920, 1922; Hardy, 1949; Hatt, 1929; Seton, 1929; Smith, C., 1965).
- \**C. collinitus* (Pers. ex Fr.) Fr.: Rodentia (Hastings and Mottram, 1916).
- \**C. distans* Peck: Rodentia (Miller and Halls, 1969).
- \**C. varius* (Schaeff. ex Fr.) Fr.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).
- Flammula sapinea*: See *Gymnopilus sapineus*.
- Gymnopilus sapineus* (Fr.) Maire: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).
- Inocybe* spp.: *Neotoma floridana magister* (Newcombe, 1930; Poole, 1940); *Tamiasciurus hudsonicus hudsonicus* (Hardy, 1949).
- \**Rozites caperata* (Pers. ex Fr.) Karst.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).

#### FAMILY CREPIDOTACEAE

- Crepidotus malachius*: See *C. nephrodes*.
- C. nephrodes* (Berk. and Curt.) Sacc.: *Sciurus carolinensis carolinensis* (Nixon, Worley, and McClain, 1968); *S. niger niger* (Nixon, Worley, and McClain, 1968).

#### FAMILY GOMPHIDACEAE

- Chroogomphus rutilus*: See *Gomphidius rutilus*.
- C. tomentosus*: See *Gomphidius tomentosus*.
- \**Gomphidius glutinosus* (Schaeff. ex Fr.) Fr.: *Sciurus vulgaris exalbidus* (Ognev, 1940).
- \**G. rutilus* (Schaeff. ex Fr.) Lund. and Nannf.: *Tamiasciurus douglasi douglasi* (Smith, C., 1965); *T. hudsonicus hudsonicus* (Smith, C., 1965).
- \**G. subroseus* Kauffm.: *Tamiasciurus douglasi douglasi* (Smith, C., 1965); *T. hudsonicus hudsonicus* (Smith, C., 1965).
- \**G. tomentosus* Murr.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).

#### FAMILY HYGROPHORACEAE

- Hygrocybe miniata* (Scop. ex Fr.) Kumm.: *Sciurus carolinensis carolinensis* (Nixon, Worley, and McClain, 1968); *S. niger niger* (Nixon, Worley, and McClain, 1968).
- \**Hygrophorus chrysodon* (Batsch ex Fr.) Fr.: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).
- \**H. hypothejus* (Fr.) Fr.: Rodentia (Hastings and Mottram, 1916); *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).
- H. miniatus*: See *Hygrocybe miniata*.
- H. pudorinus* (Fr.) Fr.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929; Odell, 1925).

#### FAMILY PAXILLACEAE

- Hygrophoropsis aurantiaca* (Wulf ex Fr.) Maire: Leporidae (Hastings and Mottram, 1916); Rodentia (Hastings and Mottram, 1916); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).
- Paxillus atrotomentosus* (Batsch ex Fr.) Fr.: Rodentia (Hastings and Mottram, 1916).
- \**P. involutus* (Batsch ex Fr.) Fr.: Leporidae (Hastings and Mottram, 1916); Rodentia (Hastings and Mottram, 1916); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).

#### FAMILY RUSSULACEAE

- Clethrionomys rutilus mikado* (Murata, 1976).
- Lactarius* spp.; Rodentia (Miller and Halls, 1969); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).
- \**L. deliciosus* (L. ex Fr.) S. F. Gray: *Sciurus carolinensis carolinensis* (Barkalow and Shorten, 1973); *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).
- \**L. lignyotus* (Fr. ex Fr.) Fr.: *Clethrionomys gapperi gapperi* (Hamilton, 1941b).
- \**L. necator* (Bull. em. Pers. ex Fr.) Karst.: Rodentia (Hastings and Mottram, 1916).
- \**L. piperatus* (L. ex Fr.) S. F. Gray: Rodentia (Miller and Halls, 1969); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1920, 1922; Hardy, 1949; Hatt, 1929).
- \**L. rufus* (Scop. ex Fr.) Fr.: Rodentia (Hastings and Mottram, 1916).
- \**L. torminosus* (Schaeff. ex Fr. S. F. Gray: *Sciurus vulgaris* (Ognev, 1940; Stakhrovskii, 1932).
- L. turpis*: See *Lactarius necator*.
- \**L. uvidus* (Fr. ex Fr.) Fr.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).
- Russula* spp.: *Clethrionomys glareolus* (Drożdż 1966); *Lemmus lemmus* (Ognev, 1948); *Napa-*

- eozapus insignis insignis* (Sheldon, 1934); *Neotoma floridana magister* (Newcombe, 1930; Poole, 1940); Sciuridae (Dowding, 1955); *Sciurus aberti aberti* (Keith, 1965); *S. carolinensis carolinensis* (Cross, 1942; Dudderar, 1967; Murrill, 1910); *Tamias striatus lysteri* (Allen, 1938; Fraleigh, 1929; Sheldon, 1936); *Tamiasciurus hudsonicus fremonti* (Hatt, 1943); *T. hudsonicus hudsonicus* (Buller, 1920, 1922; Klugh, 1927).
- \**R. adusta* (Pers. ex Fr.) Fr.: *Tamiasciurus hudsonicus hudsonicus* (Hardy, 1949).  
 \**R. alutacea* (Pers. ex Fr.) Fr.: *Tamiasciurus hudsonicus hudsonicus* (Hardy, 1949).  
 \**R. bicolor* Burl.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
*R. bicolor* var. *constans*: See *Russula bicolor*.  
*R. bicolor* var. *reticulispora*: See *Russula bicolor*.  
 \**R. blackfordae* Peck: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
 \**R. brevipes* Peck: *Tamiasciurus douglasi douglasi* (Smith, C., 1965).  
 \**R. consobrina* (Fr. ex Fr.) Fr.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
 \**R. decorans* Fr.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
*R. delicata*: see *R. brevipes*.  
*R. depallens*: See *Russula pulchella*.  
*R. disparilis* Burl.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
 \**R. emetica* (Schaeff. ex Fr.) Pers. ex Fr.: Rodentia (Hastings and Mottram, 1916); *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).  
 \**R. emetica* ssp. \**fragilis* (Pers. ex Fr.) Sing.: *Tamiasciurus hudsonicus* (Smith, C., 1965).  
*R. fragilis*: See *Russula emetica* ssp. *fragilis*.  
*R. mariae* Peck: *Tamiasciurus hudsonicus hudsonicus* (Hardy, 1949).  
 \**R. nigricans* (Bull. ex Fr.) Fr.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
 \**R. ochroleuca* (Pers. ex Fr.): *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916).  
*R. pallidostraminea* nom. prov.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
 \**R. pulchella* Borsc.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
 \**R. rosea* Quél.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
*R. rosipes* (Secr.) Bres.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).  
 \**R. rubra* (Fr.) Fr.: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).  
 \**R. xerampelina* (Schaeff. ex Secr.) Fr.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
*R. xerampelina* var. *isabelliniceps*: See *R. xerampelina*.
- FAMILY STROPHARIACEAE  
*Kuehneromyces mutabilis* (Schaeff. ex Fr.) Sing. and A. H. Sm.: *Sciurus vulgaris* (Ognev, 1940; Stakhrovskii, 1932).  
*Naematoloma fasciculare* (Huds ex Fr.) Karst.: *Sciurus vulgaris* (Moffat, 1923); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1920, 1922; Hatt, 1929).  
*Pboliota* spp.: *Sciurus vulgaris* (Ognev, 1940).  
*P. carbonaria* (Fr.) Sing.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
*P. discolor* (Pk.) Sacc.: *Sciurus carolinensis carolinensis* (Nixon, Worley, and McClain, 1968).  
*P. lenta* (Pers. ex Fr.) Sing.: *Tamiasciurus douglasi douglasi* (Smith, C., 1965).  
*P. mutabilis*: See *Kuehneromyces mutabilis*.
- FAMILY TRICHOLOMATACEAE  
*Armillaria* spp.: *Sciurus vulgaris* (Ognev, 1940).  
*A. mellea* (Vahl in Fl. Dan. ex Fr.) Karst.: Sciuridae (Miller and Halls, 1969); *Sciurus vulgaris* (Ognev, 1940; Stakhrovskii, 1932); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1920, 1922; Hatt, 1929).  
*Clitocybe* spp.: *Eutamias quadrimaculatus* (Tevis, 1952); *Sciurus vulgaris* (Ognev, 1940); *Tamiasciurus hudsonicus hudsonicus* (Hardy, 1949).  
*C. brumalis* (Fr.) Quél.: Leporidae (Hastings and Mottram, 1916); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).  
*C. maxima* (Fl. Wett. ex Fr.) Kumm.: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).  
*C. monodelpha* Morg.: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).  
 \**C. odora* (Bull. ex Fr.) Kumm.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).  
*C. virens*: See *C. odora*.  
*Collybia dryophila* (Bull. ex Fr.) Kumm.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).  
*C. maculata* (A. and S. ex Fr.) Quél.: Rodentia (Hastings and Mottram, 1916); *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).  
*C. radicata*: See *Oudemansiella radicata*.  
*C. velutipes*: See *Flammulina velutipes*.  
*Flammulina velutipes* (Curt. ex Fr.) Sing.: *Sciurus carolinensis carolinensis* (Nixon, Worley, and McClain 1968).  
*Lentinus lepideus* (Fr. ex Fr.) Fr.: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).  
 \**Lepista personata* (Fr. ex Fr.) Cke.: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).  
*Lyophyllum ulmarium* (Bull. ex Fr.) Kühner: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).  
*Marasmius* spp.: *Orcytolagus cuniculus* (Kumerloeve, 1956).  
*Oudemansiella radicata* (Relh. ex Fr.) Sing.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929; Klugh, 1927).  
*Tricholoma* spp.: *Tamiasciurus hudsonicus hudsonicus* (Hardy, 1949).  
*T. equestre*: See *T. flavovirens*.



\**T. flavovirens* (Pers. ex Fr.) Lund.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929; Odel, 1925).

*T. personatum*: See *Lepista personata*.

\**T. terreum* (Schaeff. ex Fr.) Kumm.: *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916).

ORDER APHYLLOPHORALES  
FAMILY CANTHARELLACEAE

*Cantharellus aurantiacus*: See *Hygrophoropsis aurantiaca* in Paxillaceae.

\**C. cibarius* Fr.: *Sciurus vulgaris* (Ognev, 1940; Stakhrovskii, 1932); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922; Hatt, 1929; Klugh, 1927).

*C. floccosus*: See *Gomphus floccosus*.

\**Gomphus floccosus* (Schw.) Sing.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).

FAMILY CLAVARIACEAE

*Clavaria* spp.: *Neotoma floridana magister* (Newcombe, 1930; Poole, 1930); *Tamiasciurus hudsonicus hudsonicus* (Smith, 1968).

*C. aurea*: See *Ramaria aurea*.

*C. flava*: See *Ramaria flava*.

*C. pyxidata*: See *Clavicornia pyxidata*.

*Clavicornia pyxidata* (Fr.) Poty: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).

\**Ramaria aurea* (Fr.) Quél.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929; Klugh, 1927).

\**R. flava* (Fr.) Quél.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).

*Sparassis crispa* Fr.: Leporidae (Hastings and Mottram, 1916).

FAMILY HYDNACEAE

*Hericium caput-ursi* (Fr.) Banker: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).

*Hydnum* spp.: *Tamiasciurus hudsonicus hudsonicus* (Smith, 1968).

*H. caput-ursi*: See *Hericium caput-ursi*.

*H. erinaceus* Bull.: *Sciurus carolinensis carolinensis* (Nixon, Worley, and McClain, 1968); *S. niger niger* (Nixon, Worley, and McClain, 1968).

*H. fulgineo-violaceum* Kalchb. in Fr.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).

\**H. imbricatum* L. ex Fr.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).

\**H. repandum* L. ex Fr.: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).

FAMILY POLYPORACEAE

*Sciurus griseus griseus* (Grinnel and Storer, 1924).

*Daedalea* spp.: *Sciurus carolinensis carolinensis* (Dudderar, 1967).

*Fomes fomentarius* (L. ex Fr.) Kickx: *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).

*F. pinicola* (Swartz ex Fr.) Cooke: *Tamiasciurus hudsonicus hudsonicus* (Hardy, 1949).

*Pleurotus* spp.: *Sciurus niger niger* (Packard, 1956).

*P. ostreatus* (Jacq. ex Fr.) Kumm.: *Sciurus carolinensis carolinensis* (Nixon, Worley, and McClain, 1968); *S. niger niger* (Nixon, Worley, and McClain, 1968); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929; Klugh, 1927).

*P. ulmarius*: See *Lyophyllum ulmarium*.

*Polyporus* spp.: *Neotoma cinerea orolestes* (Warren, 1920); *Sciurus carolinensis carolinensis* (Dudderar, 1967); *Spermophilus lateralis* (Gordon, 1943); *Sylvilagus palustris paludicola* (Blair, 1936); *Tamiasciurus hudsonicus fremonti* (Hatt, 1943); *T. hudsonicus hudsonicus* (Hardy, 1949; Smith, 1968); *T. hudsonicus loquax* (Layne, 1954).

*P. amarus* Hedgc.: Sciridae (Boyce, 1920); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).

*P. betulinus* Bull. ex Fr.: Leporidae (Hastings and Mottram, 1916); *Sciurus vulgaris* (Ognev 1940); *Tamiasciurus hudsonicus hudsonicus* (Buller, 1922).

*P. picipes* Fr.: *Sciurus carolinensis carolinensis* (Nixon, Worley, and McClain, 1968); *S. niger niger* (Nixon, Worley, and McClain, 1968).

*P. sulphureus* Bull. ex Fr.: *Sciurus carolinensis carolinensis* (Nixon, Worley, and McClain, 1968); *S. niger niger* (Nixon, Worley, and McClain, 1968).

*P. varius* Fr.: *Neotoma fuscipes* (Linsdale and Tevis, 1951).

*Polysticus* spp.: See *Polyporus* spp.

FAMILY STEREEACEAE

*Stereum hirsutum* Willdenow ex Fr.: *Sciurus carolinensis carolinensis* (Murrill, 1910).

ORDER HYMENOGASTRALES

FAMILY HYMENOGASTRACEAE

*Rodentia* (Gäumann and Dodge, 1928).

\**Alpova diplophloea* (Zeller and Dodge) Trappe: *Tamiasciurus douglasi douglasi* (Smith, C., 1965); *T. hudsonicus hudsonicus* (Smith, C., 1965).

*Gautieria* spp.: *Neotoma* spp. (Parks, 1919); *N. fuscipes annectens* (Parks, 1922); *Sciurus aberti aberti* (Stephenson, 1975); *S. griseus griseus* (Stienecker and Browning, 1970).

\**G. graveolens* Vitt.: *Tamiasciurus douglasi douglasi* (Smith, C., 1965); *T. hudsonicus hudsonicus* (Smith, 1965; Smith, C., 1965).

\**G. morchelliformis* Vitt.: *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).

*Hydnangium* spp.: *Neotoma fuscipes annectens* (Parks, 1922).

*Hymenogaster* spp.: *Blarina brevicauda* (Whitaker, 1962; Whitaker and Ferraro, 1963); *Clethrionomys gapperi gapperi* (Whitaker, 1962); *C. glareolus* (Drozd, 1966); *Microtus oeconomus macfarlandi* (Whitaker, 1962); *Napaeozapus insignis insignis* (Whitaker, 1962, 1963a);

- Neotoma fuscipes annectens* (Parks, 1922); Rodentia (Ingold, 1961, 1966); *Zapus hudsonius* (Whitaker, 1962).
- \**H. tener* Berk.: *Tamiasciurus hudsonicus hudsonicus* (Hardy, 1949).
- Martellia* spp.: *Sciurus aberti aberti* (Stephenson, 1975).
- Rhizopogon* spp.: *Clethrionomys glareolus* (Drozd, 1966); Rodentia (Ingold, 1966; Miller and Halls, 1969); Sciuridae (Miller and Halls, 1969); *Sciurus aberti aberti* (Stephenson, 1975); *S. griseus griseus* (Stienecker and Browning, 1970).
- R. diplophloeus*: See *Alpova diplophloeus*.
- \**R. luteolus* Fr. and Nordh.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).
- \**R. maculatus* Zeller and Dodge: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).
- \**R. occidentalis* Zeller and Dodge: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).
- R. pachyphloeus*: See *Alpova diplophloeus*.
- R. provincialis*: See *R. vulgaris*.
- \**R. roseolus* Corda sensu A. H. Sm.: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).
- \**R. rubescens* (Tul.) Tul.: *Tamiasciurus douglasi douglasi* (Smith, C., 1965); *T. hudsonicus hudsonicus* (Smith, C., 1965).
- \**R. vinicolor* Sm.: *Tamiasciurus douglasi douglasi* (Fogel, 1976).
- \**R. viridis* Zeller and Dodge: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).
- \**R. vulgaris* (Vitt.) M. Lange: *Tamiasciurus hudsonicus hudsonicus* (Smith, C., 1965).
- \**Truncocolumella citrina* Zeller: *Tamiasciurus douglasi douglasi* (Fogel, 1976); *T. hudsonicus hudsonicus* (Smith, C., 1965).
- ORDER HYSTERANGIALES  
FAMILY HYSTERANGIACEAE  
Rodentia (Gäumann and Dodge, 1928).
- Hysterangium* spp.: *Neotoma* spp.: (Parks, 1919); *N. fuscipes annectens* (Parks, 1922); *Sciurus aberti aberti*: (Stephenson, 1975); *S. griseus griseus* (Stienecker and Browning, 1970).
- FAMILY MELANOGASTRACEAE  
*Melanogaster* spp.: *Clethrionomys glareolus* (Drozd, 1966); *Neotoma* spp. (Parks, 1919); *N. fuscipes* (Linsdale and Tevis, 1951); *N. fuscipes annectens* (Parks, 1922); *Sciurus griseus griseus* (Stienecker and Browning, 1970).
- \**M. ambiguus* (Vitt.) Tul.: Sciuridae (Langham, 1916); *Sciurus carolinensis carolinensis* (Hastings and Mottram, 1916); *Tamiasciurus hudsonicus hudsonicus* (Hatt, 1929).
- M. broomeianus* Berk. in Tul.: Leporidae (Hawker, 1954).
- M. variegatus* (Vitt.) Tul. emend. Zeller and Dodge: *Microtus pennsylvanicus* (Bakerspigel, 1958); *Peromyscus maniculatus* (Bakerspigel, 1958).
- M. variegatus* var. *broomeianus*: See *M. broomeianus*.
- ORDER LYCOPERDALES  
*Dasytus novemcinctus mexicanus* (Kalmbach, 1944); Phascolomidae (Troughton, 1947).
- FAMILY GEASTERACEAE  
*Geastrum* spp.: *Tamias striatus griseus* (Schmidt, 1931).
- FAMILY LYCOPERDACEAE  
*Neotoma albigula warreni* (Finley, 1958); *Tamias striatus fisheri* (Tunis, 1971).
- Castoreum* spp.: Leporidae (Cleland, 1935); Marsupialia (Cleland, 1935).
- C. radicum* Cke. and Mass.: Marsupialia (Cunningham, 1942).
- Lycoperdon* spp.: *Apodemus sylvaticus* (Miller, 1954); *Neotoma cinerea* (Dalquest, 1948); *N. cinerea occidentalis* (Taylor, 1920); *N. floridana magister* (Poole, 1940); *Oryctolagus cuniculus* (Kumerloeve, 1956); *Sciurus aberti aberti* (Keith, 1965); *S. carolinensis carolinensis* (Hastings and Mottram, 1916); *Tamiasciurus hudsonicus loquax* (Connor, 1960).
- \**L. gemmatum* Batsch: *Sciurus carolinensis carolinensis* (Nixon, Worley, and McClain, 1968); *S. niger niger* (Nixon, Worley, and McClain, 1968).
- L. pyriforme* Pers.: *Tamiasciurus hudsonicus hudsonicus* (Hardy, 1949).
- Mesopbellia* spp.: Leporidae (Cleland, 1936; Cunningham, 1942); Marsupialia (Cleland, 1935; Cunningham, 1942).
- M. arenaria* Berk.: Leporidae (Cunningham, 1942); Macropodidae (Cunningham, 1942); Marsupialia (Cunningham, 1942); Permelidae (Cunningham, 1942).
- M. pachytrix* (Cke. and Mass.) Lloyd: Leporidae (Cleland, 1935).
- ORDER SCLERODERMATALES  
FAMILY SCLERODERMATACEAE  
*Scleroderma* spp.: *Oryctolagus cuniculus* (Kumerloeve, 1956).
- \**S. citrinum* Pers.: *Neotoma floridana magister* (Newcombe, 1930).
- S. hypogaeum* Zeller: *Neotoma fuscipes* (Linsdale and Tevis, 1951).
- S. vulgata*: See *S. citrinum*.
- MISCELLANEOUS REPORTS  
Unnamed fungi: *Apodemus flavicollis* (Drozd, 1966; Zemanek, 1972); *A. sylvaticus* (Hansson, 1970; 1971; Holíšová, 1960; Southern, 1964); *Clethrionomys* spp. (Palmer, 1954; Walker, 1964); *C. gapperi abbascae* (Harper, 1956); *C. gapperi gapperi* (Connor, 1960; Doutt, Heppenstall, and Guilday, 1967; Jackson, 1961; Larrison, 1970; Linzey and Linzey, 1973); *C. gapperi paludicola* (Doutt, Heppenstall, and Guilday, 1967); *C. gapperi rhoadsii* (Connor, 1953); *C. gapperi rupicola* (Doutt, Heppenstall, and Guilday, 1967); *C. glareolus* (Hansson, 1969, 1971; Holíšová, 1971, 1972; Miller, 1954; Ognev, 1950; Southern, 1964; Zem-

anek, 1972); *C. rufocanus* (Hansson, 1969); *Dasybus novemcinctus mexicanus* (Fitch, Goodrum, and Newman, 1952; Kalmbach, 1944; Miller and Halls, 1969); *Dicrostonyx groenlandicus* (Dunaeva and Kucheruk, 1941; Watson, 1956); *D. torquatus* (Ognev, 1948); *Didelphis virginiana* (Palmer, 1964; Sumner and Dixon, 1953); *Dipodomys* spp. (Cahalane, 1947; Palmer, 1954); *Eutamias* spp. (Cahalane, 1947; Gordon, 1943; Hamilton, 1939; Olin, 1961; Orr, 1949; Palmer, 1954; Walker, 1964); *E. cinereicollis cinereicollis* (Bailey, 1931); *E. minimus neglectus* (Hamilton, 1943); *E. minimus pictus* (Bailey, 1936); *E. quadrivittatus quadrivittatus* (Bailey, 1931; MacClintock, 1970); *E. speciosus* (Ingles, 1947; Tevis, 1953); *E. townsendii* (Larrison, 1970); *E. townsendii senex* (Bailey, 1936); *Glaucomys* spp. (Palmer, 1954); *G. sabrinus* (Cahalane, 1947; MacClintock, 1970; McKeever, 1960); *G. sabrinus fuscus* (Hamilton, 1943); *G. sabrinus macrotis* (Connor, 1960; Jackson, 1961); *G. volans volans* (Cahalane, 1947; Corbet, 1966; MacClintock, 1970; Ognev, 1940; Schwartz and Schwartz, 1959); *G. volans volans* (Connor, 1960); *Isoden obesulus* (Brazenor, 1950); *Lemmus lemmus* (Corbet, 1966; Marsden, 1964; Ognev, 1948); Leporidae (Rolfe and Rolfe, 1925); *Lepus europaeus* (Hansson, 1970); Macropodidae (Cleland, 1934); *Mastomys* spp. (Walker, 1964); *Microdipodops* spp. (Cahalane, 1947); *Microtus agrestis* (Hansson, 1970, 1971); *M. pennsylvanicus* (Golley, 1960); *Neotoma* spp. (Hamilton, 1939); *N. cinerea* (Dalquest, 1948); *N. cinerea occidentalis* (Taylor, 1920); *N. cinerea orolestes* (Warren, 1942); *N. floridana* (Hall, 1955; Schwartz and Schwartz, 1959); *N. floridana attwateri* (Bailey, 1903; Rainey, 1956); *N. floridana magister* (Hamilton, 1943; Newcombe, 1930; Paradiso, 1969; Roberts and Early, 1952); *N. fuscipes* (Linsdale and Tevis, 1951); *N. fuscipes macrotis* (Howell, 1926); *N. lepida lepida* (Barnes, 1922, 1927); *Papio cyncephalus* (Altman and Altman, 1970); *Perognathus parvus* (Johnson, 1961); *Peromyscus boyleyi* (Jameson, 1952); *P. leucopus* (Miller and Halls, 1969); *P. leucopus noveboracensis* (Hamilton, 1941b); *P. maniculatus* (Dalquest, 1948; Jameson, 1952; Miller and Halls, 1969; Schwartz and Schwartz, 1959; Walton, 1898); *P. maniculatus artemisiae* (Williams, 1959); *P. maniculatus bairdii* (Whitaker, 1966); *P. maniculatus gracilis* (Connor, 1960; Hamilton, 1941b); *P. maniculatus osgoodi* (Williams, 1959); *P. truei* (Bradford, 1974); Phascologomidae (Carter, Hill, and Tate, 1945); Peramelidae (Cleland, 1934); *Phenacomys intermedius mackenzii* (Harper, 1956); *Pitymys subterraneanus* (Holišová, 1965); *P. taticus* (Holišová, 1965); *Potorous tridactylus* (Brazenor, 1950); *P. tridactylus apicalis* (Guiler, 1971); Rodentia (Miller and Halls, 1969; Rolfe and Rolfe, 1925); Sciuridae (Brehm, 1895; Hamilton, 1939; Miller and Halls, 1969; Rolfe and Rolfe, 1925); *Sciurus* spp. (Walker, 1964); *S. aberti aberti* (Bailey, 1931; Cahalane, 1947; MacClintock, 1970; Palmer, 1954; Patton, 1975; Seton, 1929; Stephenson, 1974); *S. carolinensis carolinensis* (Bailey, 1946; Ballou, 1927; Barber, 1954; Bernard, 1937; Burt, 1957; Cahalane, 1947; Cranbrook and Payn, 1962; Cross, 1942, 1946; Goodrum, 1940; Halls and Stransky, 1971; Hoffmeister and Mohr, 1957; Lyon, 1936; MacClintock, 1970; Martin et al., 1951; Miller and Halls, 1969; Schwartz and Nelson, 1918; Palmer, 1954; Peterson, 1966; Robinson and Cowan, 1954; Schwartz and Schwartz, 1959; Shorten, 1962; Shorten and Courtier, 1955; Southern, 1964); *S. carolinensis hypophaeus* (Jackson, 1961; Larrison, 1970); *S. carolinensis pennsylvanicus* (Doutt, Heppenstall, and Guilday, 1967); *S. griseus griseus* (Cahalane, 1947; Miller and Halls, 1969; Sumner and Dixon, 1953); *S. kaibabensis* (Cahalane, 1947; Goldman, 1928; Hall, 1967; MacClintock, 1970; Nelson, 1918; Palmer, 1954); *S. niger cinereus* (Bailey, 1946); *S. niger niger* (Anthony and McSpadden, 1917; Allen, 1943; Barber, 1954; Baumgartner, 1939a, 1939b; Burt, 1957; Cahalane, 1947; Golley, 1962; Hall, 1955; Hoffmeister and Mohr, 1957; MacClintock, 1970; Martin et al., 1951; Moore, 1943; Palmer, 1954; Shafeldt, 1920); *S. niger rufiventris* (Larrison, 1970; Over and Churchill, 1941); *S. vulgaris* (Bourliere, 1956; Cranbrook and Payn, 1962; Lampio, 1967; Southern, 1964; Van Gelder, 1969); *S. vulgaris leucourus* (Shorten, 1959; 1962); *S. vulgaris orientis* (Abe, 1967); Soricidae (Martin et al., 1951); *Spermophilus* spp. (Bourliere, 1956; Walker, 1964); *S. beecheyi* (Fitch, 1948; Linsdale, 1946); *S. lateralis chrysodeirus* (Bailey, 1936; Howell, 1938; Sumner and Dixon, 1953); *S. lateralis lateralis* (Bailey, 1931; Cahalane, 1947; Dalquest, 1948; Gordon, 1943; MacClintock, 1970; Seton, 1929); *S. richardsonii* (Hansen and Ueckert, 1970); *S. undulatus kennicottii* (Bee and Hall, 1956); *S. undulatus parryii* (Howell, 1938; Seton, 1971); *Spilogale* spp. (Howell, 1906); *S. putorius* (Bailey, 1946; Martin et al., 1951); *Synaptomys cooperi cooperi* (Doutt, Heppenstall, and Guilday, 1967); *S. cooperi stonoi* (Doutt, Heppenstall, and Guilday, 1967); Talpidae (Martin et al., 1951); *Tamias* spp. Dalquest, 1948; Hamilton, 1939; Martin et al., 1951); *T. striatus fisheri* (Bailey, 1946; Tunis, 1971); *T. striatus griseus* (Jackson, 1961); *T. striatus lysteri* (Allen, 1938; Doutt, Heppenstall, and Guilday, 1967; Gifford and Whitebread, 1951; Mearns, 1899; Seton, 1929); *T. striatus striatus* (Bailey, 1946; Cahalane, 1947; Golley, 1962; MacClintock, 1970; Schwartz and Schwartz, 1959); *Tamiasciurus* spp. (Bourliere, 1956; Palmer, 1954); *T. douglasi albolimbatus* (Bailey, 1936; Stevens, 1906); *T. douglasi douglasi* (Bailey, 1936; Broadbrooks, 1958; Cahalane, 1947; Dalquest, 1948; MacClintock, 1970; Nelson, 1918; Orr, 1949; Seton, 1929); *T. douglasi mollipilosus* (Bailey, 1936); *T. hudsonicus abieticola* (Golley, 1926); *T. hudsonicus dakotensis* (Over and Churchill, 1941); *T. hudsonicus fremonti* (Cahalane, 1947; Hatt, 1943; Olin, 1961); *T. hudsonicus hudsonicus* (Anthony and McSpadden, 1917; Burt, 1957; Cahalane, 1947; Cameron, 1956; Cram, 1924; Dalquest, 1948; Dice, 1921);

Hamilton, 1939; Jackson, 1931-1932; MacClintock, 1970; Marie, 1927; Martin *et al.*, 1951; Merriam, 1884; Murrill, 1902; Nelson, 1918; Peterson, 1966; Rand, 1948; Seagars, 1949, 1950; Seton, 1929; Van Gelder, 1969; Walton, 1903; Zim and Hoffmeister, 1955); *T. hudsonicus loquax* (Bailey, 1946; Connor, 1960; Doutt, Heppenstall, and Guilday, 1967; Grim and Roberts, 1950; Layne, 1954; Lyon, 1936; Paradiso, 1969; Roberts and Early, 1952); *T. hudsonicus lychnuchus* (Bailey, 1931); *T. hudsonicus minnesota* (Hamilton, 1943; Jackson, 1961; Over and Churchill, 1941); *T. hudsonicus mogollonensis* (Bailey, 1931); *T. hudsonicus preblei* (Harper, 1956); *T. hudsonicus richardsoni* (Bailey, 1936); *T. hudsonicus streatori* (Rust, 1946); *T. hudsonicus ventorum* (Barnes, 1922, 1927); *Vombatus hirsutus* (Brazenor, 1950).

Hypogeous fungi, "truffles," hypogeous Gasteromycetes, basidiomycetous tubers, hypogeous Basidiomycetes, etc.

*Apodemus sylvaticus* (Watts, 1968); *Clethrionomys glareolus* (Watts, 1968); *Eutamias* spp. (Dowding, 1959); *E. amoenus* (Broadbrooks, 1958; Tevis, 1952); *E. quadrimaculatus* (Tevis, 1952, 1953); *E. townsendii* (Tevis, 1952, 1953); *Glaucomys sabrinus* (McKeever, 1960; Tevis, 1953); *Lepus europaeus* (Southern, 1964); Macropodidae (LeSouef and Burrell, 1926; Troughton, 1947); *Neotoma* spp. (Zeller, 1939); *N. cinerea* (Tevis, 1953); *Peromyscus boyleyi* (Tevis, 1953); *Phascolomys* spp. (LeSouef and Burrell, 1926); *Potorous* spp. (Troughton, 1943, 1944); Rodentia (Burnett, 1968; Gwynne-Vaughan, 1937); Sciuridae (Lohwag, 1932); *Sciurus griseus griseus* (Grinnel and Storer, 1924; Tevis, 1953); *Spermophilus* spp. (Dowding, 1959); *S. beecheyi* (Tevis, 1953); *Tamiasciurus douglasi douglasi* (Zeller, 1939).

### Animals and the Fungi They Eat

#### CLASS MAMMALIA

##### ORDER MARSUPIALIA

*Mesophellia* spp.; *M. arenaria*; *Castoreum* spp.; *C. radicum*.

##### FAMILY DIDELPHIDAE

*Didelphis virginiana* Kerr—opossum: *Coprinus* spp.; fungi.

##### FAMILY PERAMELIDAE

*Mesophellia arenaria*; fungi.

*Isoden obesulus* (Shaw)—short-nosed bandicoot: fungi.

##### FAMILY PHASCOLOMIDAE

Lycoperdales; fungi.

*Phascolomys* spp.: hypogeous fungi.

*Vombatus hirsutus* (Perry)—common wombat: fungi.

##### FAMILY MACROPODIDAE

*Mesophellia arenaria*; hypogeous fungi; fungi.

*Potorous* spp.: hypogeous fungi.

*P. tridactylus* Kerr—dark rat-kangaroo: fungi.

*P. tridactylus apicalis* (Gould): *Agaricus campestris*; *Oomycetales*; fungi.

##### ORDER INSECTIVORA

##### FAMILY SORICIDAE

*Endogone* spp.

*Blarina* spp.: *Glomus macrocarpus*.

*B. brevicauda* (Say)—short-tailed shrew: *Coprinus* spp.; *Endogone* spp.; *Glomus macrocarpus*;

*Hymenogaster* spp.

*Sorex* spp.: fungi.

*S. bendirii* (Merriam)—Pacific water shrew: fungi.

*S. cinereus* Kerr—masked shrew: *Endogone* spp.; *Glomus macrocarpus*.

*S. fumeus* Miller—smoky shrew: *Endogone* spp.; *Glomus macrocarpus*.

*S. pacificus*: See *Sorex vagrans pacificus*.

*S. trowbridgii* Baird—Trowbridge's shrew: *Endogone* spp.

*S. vagrans* Baird—vagrant shrew: *Endogone* spp.

*S. vagrans pacificus* Coues: fungi.

*S. vagrans yaquinae* Jackson: *Endogone* spp.

*S. yaquinae*: see *Sorex vagrans yaquinae*.

##### FAMILY TALPIDAE

*Endogone* spp.; fungi.

##### ORDER PRIMATES

##### FAMILY CYNOPITHECINAE

*Papio cynocephalus* L.—yellow baboon: fungi.

##### ORDER ENDENTATA

##### FAMILY DASYPODIDAE

*Dasybus novemcinctus mexicanus* Peters—nine-banded armadillo: Lycoperdales, fungi.

*D. novemcinctus texanus*: See *Dasybus novemcinctus mexicanus*.

##### ORDER LAGOMORPHA

##### FAMILY LEPORIDAE

*Amanita rubescens*; *Castoreum* spp.; *Clitocybe brumadis*; *Elaphomyces granulatus*; *Hygrophoropsis aurantiaca*; *Melanogaster ambiguus*; *Mesophellia* spp.; *M. arenaria*; *M. pachythrix*; *Mycena epipterygia*; *Paxillus involutus*; *Polyporus betulinus*; *Sparassis crispa*; *Tuberaceae*; *Xerocomus badius*.

*Lepus europaeus* Pallus—brown hare: *Morchella* spp.; fungi.

*Oryctolagus cuniculus* (L.)—European rabbit: *Helvella* spp., *Lycoperdon* spp., *Marasmius* spp., *Scleroderma* spp.

*Sylvilagus palustris paludicola* (Miller and Bangs)—marsh rabbit: *Polyporus* spp.

#### FAMILY OCHOTONIDAE

*Ochotona* spp.: *Endogone* spp.

*O. princeps* (Richardson)—pika: *Endogone* spp.; *Glomus fasciculatus*.

#### ORDER RODENTIA

*Amanita muscaria*; *Boletus edulis*; *Collybia maculata*; *Cortinarius collinitus*; *Elaphomyces granulatus*; *Hygrophoropsis aurantiaca*; *Hygrophorus hypobezus*; *Lactarius necator*; *L. rufus*; *Paxillus atrotomentosus*; *P. involutus*; *Russula emetica*; *Suillus bovinus*; *S. flavidus*; *S. variegatus*; *Tuber* spp.; *T. aestivum*; *Xerocomus badius*; *X. chrysenteron*; fungi.

#### FAMILY SCIURIDAE

*Amanita muscaria*; *Boletus* spp.; *Cronartium ribicola*; *Elaphomyces granulatus*; *Hygrophorus pudorinus*; *Leccinum scabrum*; *Polyporus* spp.; *Tricholoma flavovirens*; *Tuberaceae*.

*Callospermophilus* spp.: See *Spermophilus* spp.

*Citellus* spp.: See *Spermophilus* spp.

*C. parryi parryi*: See *Spermophilus undulatus parryi*.

*C. richardsonii*: See *Spermophilus richardsonii*.

*Eutamias* spp.; *Endogone* spp.; fungi.

*E. amoenus* (J. A. Allen)—yellow-pine chipmunk: *Tuberales*; hypogeous fungi.

*E. cinereicollis cinereicollis* (J. A. Allen)—gray-colored chipmunk: fungi.

*E. minimus jacksoni*: See *E. minimus neglectus*.

*E. minimus neglectus* (J. A. Allen)—least chipmunk: fungi.

*E. minimus pictus* (J. A. Allen): fungi.

*E. quadrimaculatus* (Gray)—long-eared chipmunk: *Boletus* spp.; *Clitocybe* spp.; *Tuberales*; hypogeous fungi; fungi.

*E. quadrivittatus quadrivittatus* (Say)—Colorado chipmunk: fungi.

*E. speciosus* (Merriam)—lodgepole chipmunk: fungi.

*E. townsendii* (Bachman)—Townsend's chipmunk: *Tuberales*, hypogeous fungi.

*E. townsendii senex* (J. A. Allen): fungi.

*Glaucomyces sabrinus* (Shaw)—northern flying squirrel: hypogeous fungi; fungi.

*G. sabrinus fuscus* Miller: fungi.

*G. sabrinus macrotis* (Mearns): fungi.

*G. volans volans* (L.)—southern flying squirrel: fungi.

*Pteromys volans*: See *Glaucomyces volans*.

*Sciurus* spp.: fungi.

*S. aberti aberti* Woodhouse—Abert's squirrel: *Agaricus* spp.; *Amanita muscaria*; *A. vaginata*; *Boletus* spp.; *Elaphomyces* spp.; *Gautieria* spp.; *Geopora cooperi*; *Glomus fasciculatus*; *Hysterangium* spp.; *Lepiota* spp.; *Lycoperdon* spp.; *Martellia* spp.; *Psathyrella* spp.; *Rhizopogon* spp.; *Russula* spp.; *Tuber* spp.; fungi.

*S. carolinensis carolinensis* Gmelin—eastern gray squirrel: *Agaricaceae*; *Amanita muscaria*; *Balsamia vulgaris*; *Boletus* spp.; *B. calopus*; *B. impolitus*; *Collybia maculata*; *Crepidotus malachias*; *Daedalea* spp.; *Dibotryon* spp.; *Elaphomyces granulatus*; *Flammulina velutipes*; *Hydnum erinaceus*; *Hygrocybe miniata*; *Hygrophorus hypobezus*; *Lactarius deliciosus*; *Leccinum scabrum*; *Lycoperdon* spp.; *Melanogaster ambiguus*; *Paxillus involutus*; *Pholiota discolor*; *Pleurotus ostreatus*; *Pluteus atricapillus*; *Polyporus* spp.; *P. picipes*; *P. sulphureus*; *Russula* spp.; *R. emetica*; *R. ochroleuca*; *Stereum hirsutum*; *Suillus bovinus*; *Tricholoma terreum*; fungi.

*S. carolinensis hypophaeus* Merriam: fungi.

*S. carolinensis pennsylvanicus* Ord.: fungi.

*S. cinereus*: See *S. carolinensis carolinensis*.

*S. douglasii*: See *Tamiasciurus douglasii*.

*S. fremonti lychnuchus*: See *Tamiasciurus hudsonicus lychnuchus*.

*S. fremonti mogollonensis*: See *Tamiasciurus hudsonicus mogollonensis*.

*S. griseus griseus* Ord—western gray squirrel: *Gautieria* spp.; *Hysterangium* spp.; *Melanogaster* spp.; *Polyporaceae*; *Rhizopogon* spp.; hypogeous fungi; fungi.

*S. hudsonicus*: See *Tamiasciurus hudsonicus hudsonicus*.

*S. hudsonicus ventorum*: See *Tamiasciurus hudsonicus ventorum*.

*S. leucourus*: See *Sciurus vulgaris leucourus*.

*S. kaibabensis* Merriam—Kaibab squirrel: fungi.

*S. niger cinereus* L.—eastern fox squirrel: fungi.

*S. niger neglectus*: See *S. niger cinereus*.

*S. niger niger* L.: *Agaricaceae*; *Boletus* spp.; *Crepidotus nephrodes*; *Flammulina velutipes*; *Hydnum erinaceus*; *Hygrocybe miniata*; *Lycoperdon gemmatum*; *Pholiota discolor*; *Pluteus atricapillus*; *Pleurotus* spp.; *P. ostreatus*; *Polyporus picipes*; *P. sulphureus*; fungi.

- S. niger rufiventer* E. Geoffroy St.-Hilaire: fungi.  
*S. vulgaris* (L.)—European red or brown squirrel: *Agaricus campestris*; *Amanita* spp.; *Armillaria* spp.; *A. mella*; *Balsamia vulgaris*; *Boletus edulis*; *Cantharellus cibarius*; *Calocera viscosa*; *Kuehneromyces mutabilis*; *Lactarius torminosus*; *Leccinum aurantiacum*; *L. scabrum*; *Naematoloma fasciculare*; *Polyporus betulinus*; *Suillus luteus*; *Xerocomus*; fungi.  
*S. vulgaris exalbidus* Pallas: *Clitocybe* spp.; *Ganoderma* spp.; *Gomphidius glutinosus*; *Leccinum aurantiacum*; *L. scabrum*; *Pholiota* spp.; *Pluteus nanus*; *Suillus bovinus*; *S. luteus*; *S. piperatus*; *Xerocomus chrysenteron*; *X. subimentosus*.  
*S. vulgaris leucourus* Kerr.: Agaricales; fungi.  
*S. vulgaris orientis* Thomas: fungi.  
*S. vulgaris varius*: See *Sciurus vulgaris*.  
*Spermophilus* spp.: Tuberales; hypogeous fungi; fungi.  
*S. beecheyi* (Richardson)—California ground squirrel: hypogeous fungi; fungi.  
*S. lateralis* (Say)—golden-mantled ground squirrel: *Polyporus* spp.; Tuberales; fungi.  
*S. lateralis chrysodermus* (Merriam): fungi.  
*S. richardsonii* (Sabine)—Richardson's ground squirrel: fungi.  
*S. undulatus kennicottii* (Ross)—arctic ground squirrel: fungi.  
*S. undulatus parryi* (Richardson): fungi.  
*Tamias* spp.; fungi.  
*T. striatus fisheri* A. H. Howell—eastern chipmunk: Lycoperdaceae; fungi.  
*T. striatus griseus* Mearns: *Geastrum* spp.; fungi.  
*T. striatus lysteri* (Richardson): *Amanita vaginata*; *Russula* spp.; fungi.  
*T. striatus striatus* (L.): fungi.  
*Tamiasciurus douglasi albolimbatus* (J. A. Allen)—Douglas' squirrel: fungi.  
*T. douglasi cascadenis*: See *T. douglasi mollipilosus*.  
*T. douglasi douglasi* (Bachman): *Alpova diplobloesus*; *Boletus mirabilis*; *Gautieria graveolens*; *Gomphidius rutilus*; *G. subroseus*; *Pholiota lenta*; *Rhizopogon rubescens*; *R. vinicolor*; *Russula delicata*; *Suillus granulatus*; *S. tomentosus*; *Truncocolumella citrina* var. *citrina*; Tuberales; fungi; hypogeous fungi.  
*T. douglasi mollipilosus* (Aud. and Bachman): fungi.  
*T. fremonti*: See *T. hudsonicus fremonti*.  
*T. hudsonicus abieticola* (A. H. Howell)—red squirrel: fungi.  
*T. hudsonicus dakotensis* (J. A. Allen): fungi.  
*T. hudsonicus fremonti* (Aud. and Bachman): *Polyporus* spp.; *Russula* spp.; fungi.  
*T. hudsonicus hudsonicus* (Erxleben): *Alpova diplobloesus*; *Amanita* spp.; *A. muscaria*; *A. phalloides*; *Amillaria mellea*; *Boletinus pictus*; *Boletus* spp.; *B. mirabilis*; *Cantharellus cibarius*; *Clitocybe* spp.; *C. brumalis*; *C. maxima*; *C. monodelpha*; *C. odora*; *Clavicornia pyxidata*; *Collybia dryophila*; *C. maculata*; *Corinariis* spp.; *C. varius*; *Elaphomyces granulatus*; *Fomes fomentarius*; *F. pinicola*; *Gautieria graveolens*; *G. morchelliformis*; *Gomphidius rutilus*; *G. subroseus*; *G. tomentosus*; *Gomphus floccosus*; *Gymnopilus sapineus*; *Gyroporus castaneus*; *Hericiium caput-ursi*; *Hydnum* spp.; *H. fuligineo-violaceum*; *H. imbricatum*; *H. repandum*; *Hygrophoropsis aurantiaca*; *Hygrophorus hypobryus*; *H. pudorinus*; *Hymenogaster tener*; *Inocybe* spp.; *Lactarius deliciosus*; *L. piperatus*; *L. uvidus*; *Leccinum scabrum*; *L. versipellis*; *Lentinus lepideus*; *Leucoagaricus naucinus*; *Lycoperdon pyriforme*; *Lyophyllum ulmarium*; *Melanogaster ambigua*; *Morchella conica*; *M. esulenta*; *Naematoloma fasciculare*; *Oudemansiella radiata*; *Paxillus involutus*; *Peridermium barknessii*; *Pholiota carbonaria*; *Pleurotus ostreatus*; *Polyporus* spp.; *P. amarus*; *P. betulinus*; *Psathyrella* spp.; *Ramaria aurea*; *R. flava*; *Rhizopogon luteolus*; *R. maculatus*; *R. occidentalis*; *R. roseolus*; *R. rubescens*; *R. viridis*; *R. vulgaris*; *Rozites caperata*; *Russula* spp.; *R. adusta*; *R. alutacea*; *R. bicolor*; *R. blackfordae*; *R. consobrina*; *R. decolorans*; *R. disparilis*; *R. emetica*; *R. emetica* ssp. *fragilis*; *R. mariae*; *R. nigricans*; *R. pulchella*; *R. rosea*; *R. rosipes*; *R. xerampelina*; *Suillus bovinus*; *S. cavipes*; *S. granulatus*; *S. lakei*; *S. luteus*; *S. subolivaceus*; *S. tomentosus*; *S. variegatus*; *Tricholoma* spp.; *T. flavovirens*; *Truncocolumella citrina*; Tuberales; fungi.  
*T. hudsonicus loquax* (Bangs): *Lycoperdon* spp.; *Polyporus* spp.; fungi.  
*T. hudsonicus lychnuchus* (Stone and Rehn): fungi.  
*T. hudsonicus minnesota* (J. A. Allen): *Amanita* spp.; fungi.  
*T. hudsonicus mogollonensis* (Mearns): fungi.  
*T. hudsonicus preblei* A. H. Howell: fungi.  
*T. hudsonicus richardsoni* (Bachman): fungi.  
*T. hudsonicus streatorum* (J. A. Allen): fungi.  
*T. hudsonicus ventorum* (J. A. Allen): fungi.

FAMILY GEOMYIDAE

*Tuber* spp.

FAMILY HETEROMYIDAE

*Dipodomys* spp.—kangaroo rats: fungi.

*Microdipodops* spp.—kangaroo mice: fungi.

*Perognathus parvus* (Peale)—Great Basin pocket mouse: fungi.

FAMILY CASTORIDAE

*Castor canadensis* Kuhl—beaver: fungi.

FAMILY CRICETIDAE

- Endogone* spp.  
*Apodemus sylvaticus* (L.)—wood-mouse: *Lycoperdon* spp.; *Paxillus involutus*; *Xerochomus badius*; fungi.  
*A. flavicollis* Milch.—yellow-necked field mouse: fungi.  
*Clethrionomys* spp.; *Glomus macrocarpus*; fungi.  
*C. californicus californicus* (Merriam)—California red-backed vole: *Endogone flammicorona*; *E. lactiflua*; *E. pisiformis*; *Glomus fasciculatus*; *G. macrocarpus* var. *macrocarpus*; *G. microcarpus*; *G. monosporus*; *Sclerocystis rubiformis*.  
*C. gapperi atabascae* (Preble)—Gapper's red-backed vole: fungi.  
*C. gapperi gapperi* (Vigors): *Endogone* spp.; *Glomus macrocarpus*; *Hymenogaster* spp.; *Lactarius lignyotus*; fungi.  
*C. gapperi paludicola* Doult: fungi.  
*C. gapperi rhoadsii* (Stone): fungi.  
*C. gapperi rupicola* E. L. Poole: fungi.  
*C. glareolus* Schreb.—European bank vole: *Balsamia* spp.; *Genea* spp.; *Hydnotrya* spp.; *Hymenogaster* spp.; *Melanogaster* spp.; *Rhizopogon* spp.; *Russula* spp.; *Tuber* spp.; fungi.  
*C. rufocanus* Sund.—red-gray vole: fungi.  
*C. rutilus mikado* Thomas—Emperor vole: Boletaceae Russulaceae.  
*Dicrostonyx groenlandicus* (Traill)—Greenland varying lemming: fungi.  
*D. torquatus* Pallas—collared lemming: fungi.  
*Lagurus curtatus* (Cope)—sagebrush vole: *Endogone* spp.; *Glomus fasciculatus*.  
*Lemmiscus curtatus*: See *Lagurus curtatus*.  
*Lemmus lemmus* L.—Norwegian lemming: *Boletus* spp.; *Russula* spp.; fungi.  
*Microtus agrestis* (L.)—North Europe dark vole: fungi.  
*M. chrotorrhinus* (Miller)—rock vole: *Endogone* spp.  
*M. longicaudus* (Merriam)—long-tailed vole: *Endogone* spp.  
*M. ochrogaster* (Wagner)—prairie vole: *Endogone* spp.  
*M. oeconomus macfarlandi* Merriam—tundra vole: *Endogone* spp.; *Hymenogaster* spp.  
*M. oregoni* (Bachman)—Oregon meadow mouse: *Fischerula subcaulis*.  
*M. pennsylvanicus* (Ord)—meadow vole: *Endogone* spp.; *Glomus fasciculatus*; *Melanogaster variegatus*; fungi.  
*M. pinetorum* (Le Conte)—pine vole: *Endogone* spp.  
*Neotoma* spp.: *Gautieria* spp.; *Hymenogastrales*; *Hysterangium* spp.; *Melanogaster* spp.; *Tuber rufum* var. *nitidum*; hypogeous fungi.  
*N. albigula warreni* Merriam—white-throated wood rat: Lycoperdaceae.  
*N. cinerea* (Ord)—bushy-tailed wood rat: *Lycoperdon* spp.; fungi; hypogeous fungi.  
*N. cinerea occidentalis* Baird: *Lycoperdon* spp.; fungi.  
*N. cinerea orolestes* Merriam: Agaricaceae; *Polyporus* spp.; fungi.  
*N. desertorum*: See *N. lepida lepida*.  
*N. floridana atwateri* Mearns—eastern wood rat: fungi.  
*N. floridana floridana* (Ord): fungi.  
*N. floridana magister* Baird: Agaricaceae; *Agaricus* spp.; *Boletus* spp.; *B. pallidus*; *Clavaria* spp.; *Inocybe* spp.; *Lycoperdon* spp.; *Russula* spp.; *Scleroderma citrinum*; fungi.  
*N. fuscipes annectens* Elliot—dusky-footed wood rat: Agaricaceae; Boletaceae; *Elaphomyces* spp.; *Gautieria* spp.; *Hydnangium* spp.; *Hymenogastrales*; *Hymenogaster* spp.; *Hysterangium* spp.; *Melanogaster* spp.  
*N. fuscipes fuscipes* Baird: *Helvella lacunosa*; *Melanogaster* spp.; *Morchella conica*; *Paxina acetabulum*; *Peziza* spp.; *P. emileia*; *Pluteus cervinus*; *Polyporus varius*; *Scleroderma hypogaeum*; *Tuber* spp.  
*N. fuscipes macrotis* Thomas: fungi.  
*N. lepida lepida* Thomas—desert wood rat: fungi.  
*N. magister*: See *N. floridana magister*.  
*N. mexicana fallax* Merriam—Mexican wood rat: fungi.  
*N. pennsylvanica*: See *N. floridana magister*.  
*Onychomys leucogaster* (Wied-Neuwied)—northern grasshopper mouse: *Endogone* spp.; *Glomus fasciculatus*.  
*Oryzomys palustris* (Harlan)—marsh rice rat: *Endogone* spp.  
*Peromyscus* spp.: *Glomus macrocarpus*.  
*P. boylii* (Baird)—brush mouse: Tuberales; hypogeous fungi.  
*P. gossypinus* (Le Conte)—cotton mouse: *Endogone* spp.  
*P. leucopus* (Rafinesque)—white-footed mouse: *Endogone* spp.; Uredinales; fungi.  
*P. leucopus noveboracensis* (Fisher): *Endogone* spp.; *Glomus macrocarpus*; fungi.  
*P. maniculatus* (Wagner)—deer mouse: *Endogone* spp.; *E. flammicorona*; *E. lactiflua*; *E. pisiformis*; *Glomus caledonius*; *G. fasciculatus*; *G. macrocarpus*; *G. macrocarpus* var. *macrocarpus*; *G. microcarpus*; *Melanogaster variegatus*; *Sclerocystis rubiformis*; fungi.  
*P. maniculatus artemisiae* (Rhoads): fungi.  
*P. maniculatus bairdii* (Hoy and Kennicott); *Endogone* spp.; fungi.  
*P. maniculatus gracilis* (Le Conte): *Endogone* spp.; fungi.  
*P. maniculatus osgoodi* Mearns: fungi.  
*P. truei* (Shufeldt)—piñon mouse: fungi.

*Phenacomys intermedius* Merriam—heather vole: *Endogone* spp.  
*P. intermedius mackenzii* Preble: fungi.  
*P. ungava mackenzii*: See *P. intermedius mackenzii*.  
*Pitymys pinetorum*: See *Microtus pinetorum*.  
*P. subterraneus* de Sél.-Long.—European ground vole: fungi.  
*P. tataricus* Kratochvil—Tatra pine vole: fungi.  
*Synaptomys cooperi cooperi* Baird—southern bog lemmings: *Endogone* spp.; *Glomus macrocarpus*; fungi.  
*S. cooperi gossii* (Coues): *Endogone* spp.; *Glomus macrocarpus*.  
*S. cooperi stonei* Rhoads: fungi.  
**FAMILY MURIDAE**  
*Mastacomys* spp.—broad-toothed rats: fungi.  
*Mus musculus* L.—house mouse: *Endogone* spp.  
**FAMILY ZAPODIDAE**  
*Napaeozapus insignis insignis* (Miller)—woodland jumping mouse: *Endogone* spp.; *Hymenogaster* spp.; *Russula* spp.  
*Zapus hudsonius* (Zimmermann)—meadow jumping mouse: *Endogone* spp.; *Hymenogaster* spp.  
*Z. princeps* J. A. Allen—western jumping mouse: *Endogone* spp.; *Glomus fasciculatus*; *G. pulvinatus*.  
*Z. trinotatus* Rhoads—pacific jumping mouse: *Endogone* spp.; *E. lactiflua*; *Glomus fasciculatus*; *G. monosporus*; *Sclerocystis rubiformis*.  
**ORDER CARNIVORA**  
**FAMILY MUSTELIDAE**  
*Spilogale* spp.: fungi.  
*S. putorius* (L.)—eastern spotted skunk: fungi.

### Acknowledgments

We gratefully acknowledge the advice and suggested improvements provided by Chris Maser, University of Puget Sound, Tacoma, Washington. Preparation of this paper was supported by National Science Foundation Grant BMS 76-02656 to the Coniferous Forest Biome (contribution no. 271), DEB 76-10188, and done in cooperation with the Pacific Northwest Forest and Range Experiment Station, Forestry Sciences Laboratory, Corvallis, Oregon. This is contribution no. 1,126 from the Forest Research Laboratory, Oregon State University.

### Literature Cited

- Abe, H. 1967. Notes on the ecology of *Sciurus vulgaris orientis* Thomas. Mammal. Soc. Jap., J. 3: 118-124.  
Allen, D. L. 1943. Michigan Fox Squirrel Management. Game Div. Dept. Conserv., Lansing, Mich. Game Div. Publ. 100. 404 pp.  
Allen, E. G. 1938. The Habits and Life History of the Eastern Chipmunk, *Tamias striatus lysteri*. N.Y. St. Mus. Bull. 314. 122 pp.  
Allen, J. M. 1952. Gray and fox squirrel management in Indiana. Ind. Dep. Conserv. Bull. 1. 112 pp.  
Altman, S. A., and J. Altman. 1970. Baboon Ecology. Univ. Chicago Press, Chicago. 200 pp.  
Andreotti, R. and U. Casoli. 1968. Composizione chimica e tecnologia di conservazione del tartufo. Industr. Conserve 43: 215-219.  
Anthony, H. E., and J. W. Spadler (eds). 1917. Mammals of America. University Soc., N.Y. 335 pp.  
Bailey, J. W. 1946. The Mammals of Virginia. Williams Printing Co., Richmond, Va. 416 pp.  
Bailey, V. 1905. Biological Survey of Texas. N. Amer. Fauna 25. 222 pp.  
———. 1931. Mammals of New Mexico. N. Amer. Fauna 53. 412 pp.  
———. 1936. The Mammals and Life Zones of Oregon. N. Amer. Fauna 55. 416 pp.  
Bakerspigel, A. 1956. *Endogone* in Saskatchewan rodents. Amer. J. Bot. 43: 471-475.  
———. 1958. The spores of *Endogone* and *Melanogaster* in the digestive tracts of rodents. Mycologia 50: 440-442.  
Ballou, W. H. 1927. Squirrels as mushroom eaters. J. Mammal. 8: 57-58.  
Barber, H. L. 1954. Gray and fox squirrel food habit investigations. SE Assoc. Game Fish Comm. Proc. 8: 92-94.  
Barger, N. R. 1947. The chipmunk. Wisc. Conserv. Bull. 12: 29-30.  
Barkalow, F. S., Jr., and M. Shorten 1973. The World of the Grey Squirrel. J. B. Lippincott Co., N.Y. 160 pp.  
Barnes, C. T. 1922. Mammals of Utah. Univ. Utah, Bull. 12: 1-166.



- . 1927. Utah Mammals. Univ. Utah, Bull. 17: 1-183.
- Barrett-Hamilton, G. E. H., and M. C. A. Hinton. 1921. A history of British mammals. Gurney & Jackson, London, 21: 696-720.
- Baumgartner, L. L. 1939. Food of the fox squirrel in Ohio. Ohio Wildl. Res. Sta. Release 108. 7 pp. mimeo.
- . 1939 a. Foods of the fox squirrel in Ohio. N. Amer. Wildl. Conf., Trans. 4: 579-584.
- Bee, J. W., and E. R. Hall. 1956. Mammals of Northern Alaska. Univ. of Kans. Mus. Natur. Hist. Misc. Publ. 8. 309 pp.
- Benedict, R. G., V. E. Tyler, Jr., and L. R. Brady. 1967. Studies on spore germination and growth of some mycorrhizal-associated Basidiomycetes. Mycopath. Mycol. Appl. 31: 319-326.
- Bergmann, W. 1953. The plant sterols. Ann. Rev. Plant Physiol. 4: 383-426.
- Bernard, H. 1937. L'écureuil gris. Soc. Can. d'Hist. Nat., Biblio Juenes Nat. tract 33: 1-4.
- Bessey, E. A. 1935. Text-book of Mycology. P. Blakistan's Son & Co., Philadelphia. 495 pp.
- Blair, W. F. 1936. The Florida marsh rabbit. J. Mammal. 17: 197-207.
- Bourliere, F. 1956. The Natural History of Mammals. Second edition. (transl. from French by H. M. Parschley). Alfred A. Knopf, N.Y. 364 pp., xi.
- Boyce, J. S. 1920. The dry-rot in incense cedar. U.S. Dep. Agr. Bur. Plant Ind. Bull. 871: 10.
- Bradford, D. F. 1974. Water stress of free-living *Peromyscus truei*. Ecology 55: 1407-1414.
- Brazenor, C. W. 1950. The Mammals of Victoria. Brown, Prior, Anderson Pty., Melbourne. 125 pp.
- Brehm, A. E. 1895. Brehm's Life of Animals. Vol. I. Mammalia. A. N. Marquis & Co., Chicago. 616 pp.
- Broadbrooks, H. E. 1958. Life history and ecology of the chipmunk, *Eutamias amoenus*, in eastern Washington. Univ. Mich. Mus. Zool. Misc. Publ. 103: 1-42.
- Brown, L. G., and L. E. Yeager. 1945. Fox squirrels and grey squirrels in Illinois. Ill. Nat. Hist. Surv. Bull. 23: 449-536.
- Buller, A. H. R. 1920. The red squirrel of North America as a mycophagist. Brit. Mycol. Soc. Trans. 6: 355-362.
- . 1922. The red squirrel of North America as a mycophagist. In: Researches on Fungi. Longmans, Green and Co., London. Vol. 2: 195-211.
- Burdsall, H. H., Jr. 1968. A revision of the genus *Hydnocystis* (Tuberales) and of the hypogeous species of *Geopora* (Pezizales). Mycologia 60: 496-525.
- Burnett, J. H. 1968. Fundamentals of Mycology. St. Martin's Press, N.Y. 546 pp.
- Burt, W. H. 1928. Additional notes on the life history of the Goss lemming mouse. J. Mammal. 9: 212-216.
- . 1957. Mammals of the Great Lakes Region. Univ. Mich. Press, Ann Arbor. 246 pp.
- Cahalane, V. H. 1942. Caching and recovery of food by the western fox squirrel. J. Wildl. Manage. 6: 338-352.
- . 1947. Mammals of North America. MacMillan Co., N.Y. 682 pp.
- Calhoun, J. B. 1941. Distribution and food habits of mammals in the vicinity of the Reelfoot Lake Biological Station. Tenn. Acad. Sci. J. 16: 177-185, 207-225.
- Cameron, A. W. 1956. A Guide to Eastern Canadian Mammals. Nat. Mus. Canada, Ottawa. 72 pp.
- . 1958. Canadian Mammals. Nat. Mus. Canada, Ottawa. 81 pp.
- Carter, T. D., J. E. Hill, and G. H. Tate. 1945. Mammals of the Pacific World. MacMillan Co., N.Y. 227 pp.
- Catalfomo, J., and J. M. Trappe. 1970. Ectomycorrhizal fungi: A phytochemical survey. Northw. Sci. 44: 19-24.
- Ceruti, A., F. Montacchini, and C. Duployez. 1967. Ecologia dei funghi ipogei dell' "Arboretum Taurinense". Allionia 13: 89-105.
- Chapman, F. B., and L. L. Baumgartner. 1939. Winter feeding of squirrels. Ohio Cons. Bull. 3: 6-7.
- Christensen, C. M. 1951. The Molds and Man. Univ. Minnesota Press, Minneapolis. 284 pp.
- Cleland, J. B. 1934, 1935. Toadstools and mushrooms and other larger fungi of South Australia. Adelaide; Government paper; Part I p. 1-178; Part II (1935) p. 179-362.
- Cochrane, V. W. 1963. Physiology of fungi. John Wiley & Sons, N.Y. 524 pp.
- Connor, P. F. 1953. Notes on the mammals of a New Jersey pine barrens area. J. Mammal. 34: 227-234.
- . 1960. The small mammals of Otsogo and Schoharie Counties. New York. N.Y. St. Mus. Bull. 382: 1-84.
- Cooke, M. C. 1890. Animal mycophagists. Grevillea 19: 54.
- Cooke, W. B. 1955. Fungi, lichens and mosses in relation to vascular plant communities in Eastern Washington and adjacent Idaho. Ecol. Monogr. 25: 119-180.
- Corbet, G. B. 1966. The Terrestrial Mammals of Western Europe. G. T. Foulis & Co., London. 264 pp.
- Cram, W. E. 1924. The red squirrel. J. Mammal. 5: 37-41.
- Cranbrook, Earl of, and W. H. Payn. 1962. Squirrel survey. Suffolk Nat. Soc. Trans. 12: 184-185.
- Cross, R. H., Jr. 1942. A Study of the Habits and Management of the Gray Squirrel in Virginia. Va. Polytech. Inst., Blacksburg, M. S. thesis. 40 pp.
- . 1946. The grey squirrel. Va. Wildl. 7: 14, 20.

- . 1941. On the occurrence of *Synaptomys cooperi* in forested regions. J. Mammal. 21: 195.
- . 1941 a. Food of small forest mammals in eastern United States. J. Mammal. 21: 250-263.
- . 1943. The Mammals of Eastern United States. Comstock Publ. Co., N.Y. 432 pp.
- . 1958. Life history and economic relations of the opossum (*Didelphis marsupialis virginiana*) in New York State. N.Y. St. Coll. Agr. Mem. 354: 1-48.
- Hansen, R. M., and P. N. Ueckert. 1970. Dietary similarity of some primary consumers. Ecology 51: 640-648.
- Hansson, L. 1969. Spring populations of small mammals in central Swedish Lapland in 1964-68. Oikos 20: 431-450.
- . 1970. Methods of morphological diet micro-analysis in rodents. Oikos 21: 255-266.
- . 1971. Small rodent food, feeding and population dynamics. Oikos 22: 183-198.
- Hardy, G. A. 1949. Squirrel cache of fungi. Can. Field Nat. 63: 86-87.
- Harley, J. L. 1969. The Biology of Mycorrhizae. Leonard Hill, London. 334 pp.
- . 1971. Fungi in ecosystems. J. Ecol. 59: 653-668.
- Harling, J., and M. McClaren. 1970. The occurrence of *Endogone macrocarpa* in stomachs of *Peromyscus maniculatus*. Sysis 3: 155-159.
- Harper, F. 1956. The Mammals of Keewatin. Univ. Kans. Mus. Nat. Hist. Misc. Publ. 12. 94 pp.
- Hastings, S., and J. C. Mottram. 1961. Observations upon the edibility of fungi for rodents. Brit. Mycol. Soc. Trans. 5: 364-378.
- Hatt, R. T. 1929. The red squirrel: its life history and habits, with special reference to the Adirondacks of New York and the Harvard Forest. Roosevelt Wild Life Annal 2: 7-146.
- . 1930. The biology of the voles of New York. Roosevelt Wild Life Bull. 5: 513-623.
- . 1943. The pine squirrel in Colorado. J. Mammal. 24: 311-345.
- Hawker, L. E. 1954. British hypogeous fungi. Roy. Soc. (London) Phil. Trans. B. 237: 430-546.
- Hering, T. F. 1966. The terricolous higher fungi of four Lake District woodlands. Brit. Mycol. Soc. Trans. 49: 369-383.
- Hodgman, C. D., R. C. Weast, R. S. Shankland, and S. M. Selby. 1963. Handbook of Chemistry and Physics. Chemical Rubber Publ. Co., Cleveland. 3604 pp.
- Hoffmeister, D. F., and C. O. Mohr. 1957. Fieldbook of Illinois Mammals. Ill. Nat. Hist. Survey Manual 4. 233 pp.
- Holišová, V. 1960. Potrava myšice křovinné *Apodemus sylvaticus* L. no Českomoravské vrchovině. Zool. Listy 9: 135-158.
- . 1965. The food of *Pitymys subterraneus* and *P. taticus* (Rodentia, Microtidae) in the mountain zone of the Sorbeto-Piceetum. Zool. Listy 14: 15-28.
- . 1971. The food of *Clethrionomys glareolus* at different population densities. Acta Sci. Nat. Brno 5: 1-34.
- . 1972. The food of *Clethrionomys glareolus* in a reed swamp. Zool. Listy 21: 293-307.
- Howell, A. B. 1926. Anatomy of the Wood Rat. Baltimore, Williams and Wilkins Co. Mongr. Amer. Soc. Mammal. 1. 225 pp.
- Howell, A. H. 1906. Revision of the skunks of the genus *Spilogale*. N. Amer. Fauna. 26: 1-55.
- . 1938. Revision of the North American ground squirrels with a classification of the North American Sciuridae. N. Amer. Fauna 56: 1-256.
- Ingles, L. G. 1947. Mammals of California. Stanford Univ. Press, Stanford. 258 pp.
- Ingold, C. T. 1953. Dispersal in Fungi. Clarendon Press, Oxford. 197 pp.
- . 1961. The Biology of Fungi. Hutchinson Educational, London. 144 pp.
- . 1966. Spore Release. In: Ainsworth, G. C., and A. S. Sussman, (eds.). The Fungi. Vol. 2: 679-707. John Wiley & Sons, N.Y. 805 pp.
- . 1973. The gift of a truffle. Brit. Mycol. Soc. Bull. 7: 32-33.
- Jackson, H. H. T. 1928. A taxonomic review of the American long-tailed shrews (genera *Sorex* and *Microsorex*). N. Amer. Fauna 51: 1-238.
- . 1931-1932. The red squirrel, chatterer of the spruces. Home Geogr. Monthly 1-2: 19-24.
- . 1961. Mammals of Wisconsin. Univ. Wisconsin Press, Madison. 504 pp.
- Jameson, E. W., Jr. 1952. Food of deer mice, *Peromyscus maniculatus* and *P. boylei*, in the northern Sierra Nevada, California. J. Mammal. 33: 50-60.
- Johnson, D. R. 1961. The food habits of rodents on rangelands of southern Idaho. Ecology 42: 407-410.
- Johnson, M. L., and B. T. Ostenson. 1959. Comments on the nomenclature of some mammals of the Pacific Northwest. J. Mammal. 40: 571-577.
- . 1968. Application of blood protein electrophoretic studies to problems in mammalian taxonomy. Syst. Zool. 17: 23-30.
- Kalmbach, E. R. 1944. The Armadillo in Its Relation to Agriculture and Game. Texas Game Fish and Oyster Comm. Bull. 60 pp.
- Kavaler, L. 1965. Mushrooms, Molds, and Miracles. New American Library, N.Y. 256 pp.
- Keith, J. O. 1965. The Abert squirrel and its dependence on ponderosa pine. Ecology 46: 150-163.
- Kessler, K. J., Jr., and R. W. Blank. 1972. *Endogone* sporocorps associated with sugar maple. Mycologia 64: 634-638.
- Klugh, A. B. 1927. Ecology of the red squirrel. J. Mammal. 8: 1-32.

- Korf, R. P. 1973. Sparassoid ascocarps in Pezizales and Tuberales. Rep. Tottori Mycol. Inst., Japan 10: 389-403.
- Krieger, L. C. C. 1967. The Mushroom Handbook. Dover Publ., N.Y. 560 pp.
- Kumerloeve, H. 1956. Kännchen, *Oryctolagus cuniculus* (Linne, 1758) und Hasen, *Lepus europaeus* Pallas, 1778, als Pilzfresser. Säug. Mitt. 4: 125-126.
- Lamb, R. J., and B. N. Richards. 1974. Survival potential of sexual and asexual spores of ectomycorrhizal fungi. Brit. Mycol. Soc. Trans. 62: 181-191.
- Lampio, T. 1967. Sex ratios and the factors contributing to them in the squirrel, *Sciurus vulgaris*, in Finland. Riistat. Julk. 29: 1-69.
- Langham, C. L. 1916. Squirrel eating *Melanogaster ambiguus*. Irish Nat. 28: 136.
- Larrison, E. J. 1970. Washington Mammals. Seattle Audubon Soc., Seattle. 243 pp.
- Layne, J. N. 1954. The biology of the red squirrel *Tamiasciurus hudsonicus loquax* (Bangs), in central New York. Ecol. Monogr. 24: 227-267.
- Le Souef, A. S., and H. Barrell. 1926. The Wild Animals of Australasia. George C. Harrap and Co., London, 288 pp.
- Lewis, D. H. 1973. Concepts in fungal nutrition and the origin of biotrophy. Biol. Rev. 48: 261-278.
- Lilly, V. G., and H. L. Burnett. 1951. Physiology of the Fungi. McGraw-Hill Book Co., N.Y. 464 pp.
- Linsdale, J. M. 1946. The California Ground Squirrel. Univ. California Press, Berkeley. 475 pp.
- , and L. P. Tevis, 1951. The Dusky-Footed Wood Rat. Univ. California Press, Berkeley. 664 pp.
- Linzey, D. W., and A. V. Linzey. 1973. Notes on food of small mammals from Great Smoky Mountains National Park, Tennessee-North Carolina. J. Elisha Mitchell Sci. Soc. 89: 6-14.
- Litten, W. 1975. The most poisonous mushrooms. Sci. Amer. 232: 91-101.
- Lohweg, H. 1932. Über Truffelverkommen. Verhandl. Zool. Bot. Ges. Wien. 82: 117-123.
- Lyon, M. W., Jr. 1936. Mammals of Indiana. Amer. Midl. Nat. 17: 1-384.
- MacClintock, D. 1970. Squirrels of North America. Van Nostrand Reinhold Co., N.Y. 184 pp.
- Marie, O. J. 1927. The Alaska red squirrel providing for winter. J. Mammal. 8: 37-40.
- Marks, G. C., N. Ditchburne, and R. C. Foster. 1968. Quantitative estimates of mycorrhiza populations in radiata pine forests. Austral. For. 32: 26-38.
- Marsden, W. 1964. The Lemming Year. Chatto and Windus, London. 252 pp.
- Martin, A. C., H. S. Zim, and A. L. Nelson. 1951. American Wildlife and Plants: a Guide to Wildlife Food Habits. McGraw-Hill Book Co., N.Y. 500 pp.
- Massee, G. 1911. British Fungi, with a Chapter on Lichens. George Routledge Sons, London. 551 pp.
- McCorkindale, N. J., S. A. Hutchinson, B. A. Pursey, W. T. Scott, and R. Wheeler. 1969. A comparison of the types of sterol found in spores of the Saproleginales and Leptomitales with those found in some other Phycomycetes. Phytochemistry 8: 861-867.
- McKeever, S. 1960. Food of the northern flying squirrel in northeastern California. J. Mammal. 41: 270-271.
- . 1964. Food habits of the pine squirrel in northeastern California. J. Wildl. Manage. 28: 402-403.
- Mearns, E. A. 1899. Mammals of the Catskill Mountains. U.S. Nat. Mus. Proc. 21: 352.
- Mehrotra, B. S. 1967. The Fungi. Intern. Publ. House, Allahabad. 331 pp.
- Melin, E., and H. Nilsson. 1957. Transport of C<sup>14</sup> labelled photosynthate to the fungal associate of pine mycorrhiza. Svensk Bot. Tidskr. 51: 166-186.
- Mendel, L. B. 1898. Chemical composition and nutritive value of some edible American fungi. Amer. J. Physiol. 1: 225-238.
- Merriam, C. H. 1884. The vertebrates of the Adirondack region of northeastern New York. Linn. Soc. N.Y. Trans. 2: 1-214.
- Metcalf, M. M. 1925. *Amanita muscaria* in Maine. Science, N. S. 61: 567.
- Milazzo, F. H. 1965. Sterol production by some wood rotting Basidiomycetes. Can. J. Bot. 43: 1347-1353.
- Miller, H. A., and L. K. Halls. 1969. Fleshy Fungi Commonly Eaten by Southern Wildlife. South. Forest Exp. Sta., New Orleans, La. 28 pp. (USDA Forest Serv. Res. Pap. SO-49.)
- Miller, R. S. 1954. Food habits of the wood-mouse, *Apodemus sylvaticus* (Linne, 1758), and the bank vole, *Clethrionomys glareolus* (Schreber, 1780), in Wytham Woods, Berkshire. Säug. Mitt. 2: 109-113.
- Moffat, C. B. 1923. Food of the Irish squirrel. Irish Nat. 32: 77-82.
- Moore, G. E. 1943. Food habits of squirrels. Mo. Conserv. 4: 14 (Dec.).
- Murata, Y. 1976. Spores of higher fungi found in the stomach of *Clethrionomys rutilus mikado* Thomas, a kind of vole. [In Japanese.] Mycol. Soc. Jap., Trans. 17: 85-87.
- Murrill, W. A. 1902. Animal mycophagists. Torreya 2: 25-26.
- . 1910. On fungi eaten by gray squirrels. Mycologia 2: 96.
- Negus, N. C., E. Gould, and R. K. Chipman. 1961. Ecology of the rice rat, *Oryzomys palustris* (Harlow), on Breton Island, Gulf of Mexico, with a critique of the social stress theory. Tulane Stud. Zool. 8: 93-123.
- Nelson, E. W. 1918. Wild animals of North America. National Geographic Soc., Washington, D.C. Pp. 385-612.

- Newcombe, C. L. 1930. An ecological study of the Allegheny cliff rat (*Neotoma pennsylvanica* Stone). J. Mammal. 11: 204-211.
- Nixon, C. M., D. M. Worley, and M. W. McClain. 1968. Food habits of squirrels in southeast Ohio. J. Wildl. Manage. 32: 294-305.
- Odell, W. S. 1925. Squirrels eating *Amanita muscaria*. Can. Field Nat. 39: 180-181.
- . 1926. Further observations on squirrels eating *Amanita*. Can. Field Nat. 40: 184.
- Ognev, S. I. 1940-1950. Mammals of the USSR and adjacent countries. Vols. IV, V, VI, and VII. Transl. by Jean Salkind. Smithsonian Inst. and NSF, Washington, D.C.
- Olin, G. 1961. Mammals of the Southwest Mountains and Mesas. Southw. Parks Monuments Assoc., Globe, Ariz. 126 pp.
- Orr, R. T. 1949. Mammals of Lake Tahoe. Calif. Acad. Sci., San Francisco. 127 pp.
- Over, W. H., and E. P. Churchill. 1941. Mammals of South Dakota. Mus. Dept. Zool., Univ. S. Dak., Vermillion. 56 pp.
- Overholts, L. E. 1953. The Polyporaceae of the United States, Alaska, and Canada. Univ. Michigan Press, Ann Arbor. 466 pp.
- Ovington, J. D. 1961. Some aspects of energy flow in plantations of *Pinus sylvestris* L. Ann. Bot., London N. S. 25: 12-20.
- Packard, R. L. 1956. The Tree Squirrels of Kansas: Ecology and Economic Importance. Mus. Nat. Hist. St. Biol. Surv., Univ. Kans. Misc. Publ. 11: 67 pp.
- Palenzona, M. 1969. Sintesi micorrizica tra *Tuber aestivum* Vitt., *Tuber brumale* Vitt., *Tuber melanosporum* Vitt. e semenzali di *Corylus avellana* L. Allionia 15: 121-132.
- Palmer, R. S. 1954. The Mammal Guide. Doubleday and Co., Garden City, N.Y. 384 pp.
- Paradiso, J. L. 1969. Mammals of Maryland. N. Amer. Fauna 66. 193 pp.
- Parks, H. E. 1919. Notes on California fungi. Mycologia 11: 15-20.
- . 1921. Californian hypogeous fungi. Tuberaceae. Mycologia 13: 301-314.
- . 1922. The genus *Neotoma* in the Santa Cruz Mountains. J. Mammal. 3: 241-253.
- Patton, D. R. 1975. Abert Squirrel Cover Requirements in Southwestern Ponderosa Pine. USDA, Forest Serv., Rocky Mt. Forest Range Exp. Sta. Res. Pap. RM-145. 12 pp.
- Peterson, R. L. 1966. The Mammals of Eastern Canada. Oxford Univ. Press, Toronto. 465 pp.
- Pilát, A., and O. Usak. 1958. Mushrooms. Spring Books, London. 340 pp.
- Poole, E. L. 1940. A life history sketch of the Allegheny wood rat. J. Mammal. 21: 249-270.
- Proudfit, F. T., and C. H. Robinson. 1961. Normal and Therapeutic Nutrition. MacMillan, N.Y. 858 pp.
- Rainey, D. G. 1956. Eastern woodrat, *Neotoma floridana*: life history and ecology. Univ. Kans. Mus. Nat. Hist. Publ. 8: 535-646.
- Rajala, P., and T. Lampio. 1963. The food of the squirrel (*Sciurus vulgaris*) in Finland in 1945-1961. Suomen Riista, Helsinki 16: 155-185.
- Rand, A. L. 1948. Mammals of the Eastern Rockies and Western Plains of Canada. Nat. Mus. Can. Bull. 108. 237 pp.
- Reess, M., and C. Fisch. 1887. Untersuchungen unter Bau und Lebengeschichte der Hirschrüffel, *Elaphomyces*. Bibl. Bot. 7: 1-24.
- Richardson, M. J. 1970. Studies on *Russula emetica* and other agarics in a Scots pine plantation. Brit. Mycol. Soc. Trans. 55: 217-229.
- Roberts, H. A., and R. C. Early. 1952. Mammal Survey of Southeastern Pennsylvania. Penn. Game Comm., Harrisburg. 70 pp.
- Robinson, D. J., and I. McT. Cowan. 1954. An introduced population of the gray squirrel (*Sciurus carolinensis gmelin*) in British Columbia. Can. J. Zool. 32: 261-282.
- Rolfe, R. T., and F. W. Rolfe. 1925. The Romance of the Fungus World. Chapman and Hall Ltd., London. 309 pp.
- Rue, L. L. 1967. Pictorial Guide to the Mammals of North America. Thos. Y. Crowell Co., N.Y. 299 pp.
- Rust, H. J. 1946. Mammals of northern Idaho. J. Mammal. 27: 308-327.
- Schmidt, F. J. W. 1931. Mammals of western Clark County, Wisconsin. J. Mammal. 12: 99-117.
- Schwartz, C. W., and E. R. Schwartz. 1959. The Wild Mammals of Missouri. Univ. Missouri Press, Kansas City. 341 pp.
- Seagers, C. 1949-1950. The red squirrel. N.Y. St. Conserv. 4: 40-44 (Dec.).
- Seton, E. T. 1909. Life-Histories of Northern Animals. Charles Scribner's Sons, N.Y. Vol. 1.
- . 1911. The Arctic Prairies. Charles Scribner's Sons, N.Y. 415 pp.
- . 1929. Lives of Game Animals. Doubleday, Doran and Co., N.Y. 4: 1-440.
- Sheldon, C. 1934. Studies on the life histories of *Zapus* and *Napaeozapus* in Nova Scotia. J. Mammal. 15: 290-300.
- . 1936. The mammals of Lake Kedgemakooge and vicinity, Nova Scotia. J. Mammal. 17: 207-215.
- Shemakhanova, N. M. 1967. Mycotrophy of Woody Plants. Israel Prog. Sci. Transl., Jerusalem, U.S. Dep. Commerce TT 66-51073. 329 pp.
- Shivrina, A. N., E. G. Platonova, and Y. P. Cherotchenko. 1968. Higher Basidiomycetes as sources of ergosterol. Mikol. Fitopatol. 2: 65-69.
- Shorten, M. 1959. Squirrels in Britain. In: Flyger, V. (ed.) Symposium on the Gray Squirrel. Chesapeake Biol. Lab., Contrib., Salomono, Md. Pp. 375-378.
- . 1962. Squirrels; Their Biology and Control. H. M. Stationary Off., London. 44 pp.

- \_\_\_\_\_, and F. A. Courtier. 1955. A population study of the grey squirrel (*Sciurus carolinensis*) in May 1954. *Ann. Appl. Biol.* 43: 494-510.
- Shufeldt, R. W. 1920. Four-footed foresters—the squirrels. *Amer. For.* 26: 37-44.
- Singer, R. 1961. *Mushrooms and Truffles*. Interscience Publ., N.Y. 269 pp.
- \_\_\_\_\_. 1962. *The Agaricales in Modern Taxonomy*. J. Cramer, Weinheim. 915 pp.
- Smith, A. H. 1949. *Mushrooms in Their Natural Habitats*. Sawyer's Inc., Portland, Or. 626 pp.
- \_\_\_\_\_. 1951. *Puffballs and Their Allies in Michigan*. Univ. Michigan Press, Ann Arbor. 131 pp.
- \_\_\_\_\_. 1965. New and unusual Basidiomycetes with comments on hyphal and spore wall reactions with Melzer's solution. *Mycopathol. Mycol. Appl.* 26: 385-402.
- Smith, C. C. 1965. Interspecific competition in the genus of tree squirrels *Tamiasciurus*. Univ. Wash., Seattle. Ph.D. thesis. 269 pp.
- \_\_\_\_\_. 1970. The coevolution of pine squirrels (*Tamiasciurus*) and conifers. *Ecol. Monogr.* 40: 349-371.
- Smith, G. M. 1938. *Cryptogamic Botany*. McGraw-Hill Book Co., N.Y. Vol. 1. 545 pp.
- Smith, M. C. 1968. Red squirrel responses to spruce cone failure in interior Alaska. *J. Wildl. Manage.* 32: 305-317.
- Southern, H. N. (ed.). 1964. *The Handbook of British Mammals*. Blackwell Sci. Publ., Oxford. 465 pp.
- Spaulding, P. 1922. Investigations of the white-pine blister rust. U.S. Dep. Agr. Bur. Plant Ind. Bull. 957: 35-36.
- Stakhrovskii, V. C. 1932. The biology of squirrels kept in open air cages. (In Russian.) *Zool. Zh.* 11: 82-104.
- Stark, N. 1972. Nutrient cycling pathways and litter fungi. *Bioscience* 22: 355-360.
- Stephens, F. 1906. *California Mammals*. West Coast Publ. Co., San Diego, Calif. 351 pp.
- Stephenson, R. L. 1974. Seasonal food habits of Abert's squirrels *Sciurus aberti*. *J. Ariz. Acad. Sci.* 9(suppl.): 8.
- \_\_\_\_\_. 1975. Reproductive Biology and Food Habits of Abert's Squirrels in Central Arizona. Arigona State Univ., M. S. thesis. 66 pp.
- Stienecker, W., and B. M. Browning. 1970. Food habits of the western gray squirrel. *Calif. Fish Game.* 56: 36-48.
- Stoner, D. 1918. The Rodents of Iowa. *Iowa Geol. Surv. Bull.* 5. 172 pp.
- Styan, K. E. 1946. Squirrel with a toadstool. *Field*, London 187: 49 (letter).
- Sumner, L., and J. S. Dixon. 1953. *Birds and Mammals of the Sierra Nevada*. Univ. California Press, Los Angeles. 484 pp.
- Sussman, A. S., and H. O. Halvorson. 1966. *Spores: Their Dormancy and Germination*. Harper and Row, N.Y. 354 pp.
- Taylor, W. P. 1920. The wood rat as collector. *J. mammal.* 1: 91-92.
- Tevis, L. 1952. Autumn foods of chipmunks and golden-mantled ground squirrels in the northern Sierra Nevada. *J. Mammal.* 33: 198-205.
- \_\_\_\_\_. 1953. Stomach contents of chipmunks and mantled squirrels in northeastern California. *J. Mammal.* 34: 316-324.
- Thaxter, R. 1922. A revision of the Endogonaceae. *Amer. Acad. Arts Sci., Proc.* 57: 291-341.
- Theodorou, C., and G. D. Brown. 1973. Inoculation of seeds and soil with basidiospores of mycorrhizal fungi. *Soil Biol. Biochem.* 5: 765-771.
- Trappe, J. M. 1962. Fungus associates of ectotrophic mycorrhizae. *Bot. Rev.* 28: 538-606.
- \_\_\_\_\_. 1971. Mycorrhiza-forming Ascomycetes. In: Hacskeylo, E. (ed.), *Mycorrhizae*. U.S. Dep. Agric., Misc. Publ. 1189: 19-37.
- \_\_\_\_\_. 1971a. A synopsis of the Carbobymycetaceae and Terfeziaceae (Tuberales). *Brit. Mycol. Soc. Trans.* 57: 85-92.
- \_\_\_\_\_. 1975. The genus *Fischerula* (Tuberales). *Mycologia* 67: 934-941.
- \_\_\_\_\_, and C. Maser. 1976. Germination of spores of *Glomus macrocarpus* (Endogonaceae) after passage through a rodent digestive tract. *Mycologia* 68: 433-436.
- Troughton, E. 1943. The kangaroo family. Rat kangaroos. *I. Austl. Mus. Mag.* 8: 171-175.
- \_\_\_\_\_. 1944. The kangaroo family. Hare wallabies. *Austl. Mus. Mag.* 8: 229-233.
- \_\_\_\_\_. 1947. *Furred Animals of Australia*. Charles Scribner's Sons, N. Y. 374 pp.
- Tunis, E. 1971. *Chipmunks on the Doorstep*. Thos. Y. Crowell Co., N.Y. 70 pp.
- Uhlig, H. C. 1955. *The Gray Squirrel: Its Life History, Ecology, and Population Characteristics in West Virginia*. W. Va. Conserv. Comm. 175 pp.
- VanGelder, R. G. 1969. *Biology of Mammals*. Charles Scribner's Sons, N.Y. 197 pp.
- Walker, E. P. 1964. *Mammals of the World*. Johns Hopkins Press, Baltimore, Maryland. Vol. 2. 1500 pp.
- Walton, M. A. 1898. The red squirrel. *Forest Stream*. P. 43.
- \_\_\_\_\_. 1903. *A hermit's wild friends*. D. Estes and Co., Boston. 304 pp.
- Warren, E. R. 1920. Notes on wood rat work. *J. Mammal.* 1: 233-235.
- \_\_\_\_\_. 1942. *The Mammals of Colorado*, Second Edition. Univ. Oklahoma Press, Norman. 330 pp.
- Watson, A. 1956. Ecological notes on the lemmings *Lemmus trimucronatus* and *Dicrostonyx groenlandicus* in Baffin Island. *J. Animal Ecol.* 25: 289-302.

- Watts, C. H. S. 1968. The foods eaten by woodmice (*Apodemus sylvaticus* and bank voles (*Clethrionomys glareolus*) in Wytham Woods, Berkshire. *J. Anim. Ecol.* 37: 25-41.
- Whitaker, J. O., Jr. 1962. *Endogone*, *Hymenogaster*, and *Melanogaster* as small mammal foods. *Amer. Midl. Nat.* 67: 152-156.
- . 1963. Food, habitat and parasites of the woodland jumping mouse in central New York. *J. Mammal.* 44: 316-321.
- . 1963 a. Food of 120 *Peromyscus leucopus* from Ithaca, New York. *J. Mammal.* 44: 418-419.
- . 1966. Food of *Mus musculus*, *Peromyscus maniculatus bairdi* and *Peromyscus leucopus* in Vigo County, Indiana. *J. Mammal.* 47: 473-486.
- , and M. G. Ferraro. 1963. Summer food of 220 short-tailed shrews from Ithaca, New York. *J. Mammal.* 44: 419.
- , and C. Maser. 1976. Food habits of five Western Orgeon shrews. *Northw. Sci.* 50: 102-107.
- Wilkins, W. H., and G. C. Harris. 1946. The ecology of the larger fungi. V. An investigation into the influence of rainfall and temperature on the seasonal production of fungi in a beechwood and a pinewood. *Ann. Appl. Biol.* 33: 179-188.
- Williams, O. 1959. Food habits of the deer mouse. *J. Mammal.* 40: 415-419.
- , and B. A. Finney. 1964. *Endogone*—food for mice. *J. Mammal.* 45: 265-271.
- Winton, A. L., and K. B. Winton. 1932. *The Structure and Composition of Foods*. John Wiley and Sons, N.Y. Vol. I. 710 pp.
- , and ———. 1935. *The Structure and Composition of Foods*. John Wiley and Sons, N.Y. Vol. II. 904 pp.
- Zeller, S. M. 1939. Developmental Morphology of *Alpova*. *Oreg. St. Monog., Stud. Bot.* 2. 19 pp.
- Zemanek, M. 1972. Food and feeding habits of rodents in a deciduous forest. *Acta Theriol.* 17: 315-325.
- Zim, H. S., and D. F. Hoffmeister. 1955. *Mammals*. Simon and Schuster, N.Y. 160 pp.
- Zimmerman, E. G. 1965. A comparison of habitat and food of two species of *Microtus*. *J. Mammal.* 46: 605-612.
- . 1965a. A comparison of food habits of two species of *Microtus*. *Ind. Acad. Sci. Proc.* 75: 281.

Received June 16, 1976

Accepted for publication February 7, 1977