
Measurement of Forest Disturbance and Regrowth With Landsat and Forest Inventory and Analysis Data: Anticipated Benefits From Forest and Inventory Analysis' Collaboration With the National Aeronautics and Space Administration and University Partners

Sean Healey¹, Gretchen Moisen², Jeff Masek³, Warren Cohen⁴, Sam Goward⁵, Scott Powell⁶, Mark Nelson⁷, Dennis Jacobs⁸, Andrew Lister⁹, Robert Kennedy¹⁰, and John Shaw¹¹

Abstract.—The Forest Inventory and Analysis (FIA) program has partnered with researchers from the National Aeronautics and Space Administration, the University of Maryland, and other U.S. Department of Agriculture Forest Service units to identify disturbance patterns across the United States using FIA plot data and time series of Landsat satellite images. Spatially explicit predictions of biomass loss and gain from 1972 to 2002 will be produced in 2-year intervals using 25 Landsat scenes distributed throughout the country. The map-based analyses that will be made possible through this collaboration will complement FIA's current ability to track disturbances at the county and State level.

Overview of the Collaboration

The Forest Inventory and Analysis (FIA) program has entered into a collaborative agreement with a diverse team of scientists for the purpose of using historical Landsat data to measure forest disturbance and regrowth since 1972. FIA analysts across the country are working with other U.S. Department of Agriculture (USDA) Forest Service scientists and collaborators from the National Aeronautics and Space Administration (NASA), the University of Maryland, and Oregon State University to create biennial maps of forest biomass change. The project has the following three stated goals:

1. To characterize disturbance regimes for forests across the United States and portions of Canada.
2. To evaluate the variability of post-disturbance forest regrowth.
3. To develop techniques that enable FIA analysts to study the disturbance history of any forested area in the country.

This collaboration has the potential to significantly improve FIA's capacity to monitor the forest changes resulting from disturbance. Historical Landsat imagery has been used to map the occurrence of several types of forest disturbances,

¹ Research Ecologist, U.S. Department of Agriculture (USDA), Forest Service, Rocky Mountain Research Station, 507 25th Street, Ogden, UT 84401. E-mail: seanhealey@fs.fed.us.

² Research Ecologist, USDA Forest Service, Rocky Mountain Research Station, 507 25th Street, Ogden, UT 84401. E-mail: gmoisen@fs.fed.us.

³ Research Scientist, National Aeronautics and Space Administration, Goddard Space Flight Center, Greenbelt, MD 20771. E-mail: Jeffrey.G.Masek@nasa.gov.

⁴ Research Forester, USDA Forest Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis, OR 97331. E-mail: wcohen@fs.fed.us.

⁵ Professor of Geography, University of Maryland, 2181 LeFrak Hall, College Park, MD 20782. E-mail: sgoward@geog.umd.edu.

⁶ Research Forester, USDA Forest Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis, OR 97331. E-mail: scottpowell@fs.fed.us.

⁷ Research Forester, USDA Forest Service, North Central Research Station, 1992 Folwell Avenue, St. Paul, MN 55108. E-mail: mdnelson@fs.fed.us.

⁸ Research Forester, USDA Forest Service, Southern Research Station, 4700 Old Kingston Pike, Knoxville, TN 37919. E-mail: djacobs@fs.fed.us.

⁹ Research Forester, USDA Forest Service, Northeastern Research Station, 11 Campus Boulevard, Suite 200, Newtown Square, PA 19073. E-mail: alister@fs.fed.us.

¹⁰ Research Forester, USDA Forest Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis, OR 97331. E-mail: robertkennedy@fs.fed.us.

¹¹ Forester, USDA Forest Service, Rocky Mountain Research Station, 507 25th Street, Ogden, UT 84401. E-mail: jdshaw@fs.fed.us.

including harvest (e.g., Cohen *et al.* 2002, Healey *et al.* 2005, Sader 1995), fire (e.g., Cocke *et al.* 2005), insect activity (e.g., Skakun *et al.* 2003), and storm events (e.g., McMaster 2005). In addition to mapping the occurrence and extent of disturbances, a few studies have attempted to measure their effect, either in general classes of tree mortality (Franklin *et al.* 2000, Jin and Sader 2005, Skakun *et al.* 2003) or as continuous variables representing change in an element of forest structure (Collins and Woodcock 1996, Healey *et al.* 2006b, Olsson 1994). The scope and precision of the maps to be produced through the current collaboration are unique. Biennial estimates of biomass loss and gain will be produced for areas across the continental United States and portions of Canada at the resolution of the Landsat pixel (~ 30 m). This production of highly specific (in time, space, and degree) estimates of change over an area of almost 1 million square km is only possible through the combined expertise of the assembled partners. The following section describes the relevant capacities of each of the partners and the contributions they are expected to make. The final section of this paper contains a discussion of the possible benefits to FIA of the products and techniques resulting from this effort.

Collaborators

NASA

The Landsat Ecosystem Disturbance Adaptive Processing System (LEDAPS) is a NASA-funded program based at the Goddard Space Flight Center. The goal of this program (http://ledaps.nascom.nasa.gov/ledaps/ledaps_NorthAmerica.html) is to map forest disturbance and regrowth across the

North American continent using three dates of Landsat imagery (1975, 1990, 2000). In meeting the significant logistical challenges of processing such a large number of images, LEDAPS has developed several automated algorithms for critical tasks such as removal of atmospheric effects, radiometric normalization, orthorectification, and disturbance detection (table 1). These algorithms will support not only the processing of the imagery needed in this project, but will also, once validated, be available to FIA for use in other projects.

The LEDAPS continentwide disturbance maps will be available by mid 2006. Because of the decadal sampling interval for these maps, it is likely that a portion of disturbances will not be detected. Vegetation regrowth following a disturbance can mask the disturbance's spectral signal if the sampling frequency is low (Healey *et al.* 2005, Jin and Sader 2005). Through the current collaboration, disturbance rates will be identified in more than 25 Landsat scenes across the country with imagery acquired at 2-year intervals from 1972 to the present using methods discussed below. These scenes will be chosen in a national-scale sampling framework so that the resultant disturbance maps may be used in concert with the LEDAPS product to improve national-level estimates of forest disturbance rates.

FIA

The national network of inventory plots maintained by FIA has a sampling intensity of at least one plot per 6,000 acres (approximately 1,400 plots per Landsat scene). In addition to measuring biometric characteristics such as biomass and basal

Table 1.—*Relevant processing algorithms under development by Landset Ecosystem Disturbance Adaptive Processing System.*

Algorithm name	Description	Current status
Indcal	Landsat-5 and Landsat-7 calibration and conversion to top-of-atmosphere reflectance	Operational
Indsr	Aerosol retrieval, atmospheric correction, conversion to surface reflectance	Operational/still in testing
Indcsm	Create cloud/shadow/snow mask	Prototype exists
Indreg	Precision image-to-image matching via Ground Control Points and orthorectification	Operational/still in testing
Inddm	Disturbance mapping using Disturbance Index (Healey <i>et al.</i> 2005)	Operational/still in testing
Indcom	Direct surface reflectance compositing across multiple acquisitions without Bidirectional Reflectance Distribution Function adjustment	Prototype exists

area at each plot, FIA records the likely cause and estimated year of forest disturbances occurring at each plot between measurements. FIA's plot data may be used in several ways to train and validate satellite-based forest change detection algorithms. A plot may be viewed categorically according to its binary FIA plot-level disturbance attribute, in which case it could be used to support the mapping of the location, but not the intensity, of disturbances. If a plot has been measured both before and after the identified disturbance, then the degree of damage may be assessed in terms of change in a stand attribute such as live volume or biomass. In this case, predictive models of disturbance intensity may be built using the relationship between the degree of measured physical change and the spectral differences seen in pre- and post-disturbance imagery. Plots that have not been revisited may still support efforts to map disturbance intensity. Measurements of attributes such as biomass from any date may be associated with contemporaneous imagery, and, if there is adequate radiometric normalization among images across time, a date-invariant predictive spectral model for that attribute may be produced. If that model is uniformly applied to normalized imagery from different dates, differences in predicted conditions may contain significant information about the intensity of local disturbances (Healey *et al.* 2006b).

FIA scientists, having long had access to the spatial coordinates of the Nation's largest forest inventory, have made important strides both in the modeling of biophysical variables using remotely sensed data (e.g., Blackard *et al.* 2006, Frescino *et al.* 2001, Lister *et al.* 2004, McRoberts *et al.* 2002, Moisen and Frescino 2002) and in the assessment of those models (Czaplewski and Patterson 2001, Edwards *et al.* 1998, Patterson and Williams 2003). In this respect, it is likely that FIA personnel will be instrumental both in interpreting information from FIA plot records and in modeling that information. FIA analysts will also be instrumental at the local level in helping identify the causes of mapped disturbances. Finally, FIA will have a role in communicating the results of this project as disturbance trends are included in regional and national reports.

Other USDA Forest Service and University Partners

While FIA's scientists have used satellite imagery and plot data to map forest conditions across large areas, the program has little experience in mapping forest changes. In contrast, other USDA Forest Service collaborators and those from the University of Maryland and Oregon State University have a good deal of experience in developing (Cohen *et al.* 1998, Huang *et al.* 2000, Powell 2004), testing (Cohen and Fiorella 1998, Healey *et al.* 2005), and applying (Cohen *et al.* 2002) Landsat-based change detection algorithms. Because of this experience, USDA Forest Service and university partners will have a leading role in developing methods for mapping change. As stated earlier, these methods will use historical Landsat imagery to both measure the intensity of forest disturbances and plot the regrowth of disturbed stands. Illustrating the degree to which this project will draw on the strengths of all collaborators, the change detection algorithms developed by the USDA Forest Service and university researchers will rely on both the mass preprocessing techniques designed by NASA personnel and the plot data and modeling techniques provided by FIA.

Benefits to FIA

This collaboration will greatly increase the spatial precision with which FIA can characterize disturbance across the United States. FIA's sample-based estimates of forest conditions typically are made at the county or State level to assure the consideration of a statistically adequate number of plots. Although the estimation of forest attributes at the county level using FIA plot data is statistically straightforward, it precludes more spatially explicit analyses. This project will produce estimates of forest change at the scale of the Landsat pixel (~ 30 m), permitting fine-scale analyses of disturbances such as harvests, fires, and wind events that are not possible using the sample-based paradigm. Study of the spatial patterns of forest recovery will likewise be possible. This section will describe several potential applications of this project's change products that may be of use to FIA.

Harvest Detection


The basic products resulting from this project will be spatially explicit biennial estimates of biomass loss or gain within approximately 25 Landsat scenes. To translate these pixel-scale predictions into maps of disturbance, likely causes of each predicted disturbance will have to be identified. This process will likely focus on spatially contiguous patches of pixels displaying abrupt drops in estimated biomass. Assignment of disturbance type may be automated using rules regarding the size, shape, spatial complexity, or texture of each patch (e.g., Cohen *et al.* 2002). Formulation of these rules will have to be made in consultation with FIA analysts and other local experts. Once the sources of individual disturbances are identified, spatial and temporal trends in harvests and other types of disturbance may be conducted.

Harvest is a significant cause of forest disturbance in many managed landscapes. Harvests that remove most or all of the trees in a stand have been mapped using Landsat imagery with relatively high accuracy (e.g., Cohen *et al.* 2002, Hall *et al.* 1989, Sader and Winne 1992). Several projects have also suggested the potential for the use of historical Landsat imagery to map partial harvests (Collins and Woodcock 1996, Olsson 1994, Sader *et al.* 2003). Landsat's short-wave infrared bands may be particularly useful in modeling the degree of canopy removal involved with a harvest (Healey *et al.* 2006b; Olsson 1994).

The production of spatially and temporally explicit maps of disturbance will enable FIA to augment its current timber output records. It will be possible to summarize harvest trends since 1972 by any combination of geographic variables, including landowner, forest type, topography, or climate. For example, in a study supporting the monitoring component of the Northwest Forest Plan, Healey *et al.* (2006a) reported trends in clearcut harvesting for both Federal and non-Federal landowners in Oregon and Washington from 1972 to 2002 (fig. 1). The study showed that while non-Federal forest owners continued to harvest at relatively high levels during the 1990s (the period coinciding with the Forest Plan), clearcutting of Federal forests virtually stopped.


It is technically possible to create similar estimates of harvest by geographic variables using only FIA data because many of these variables are stored as plot characteristics. Satellite-based estimates, however, have at least three advantages. First, while FIA survey protocols and designs may have changed over the past 30 years, the continuity of the Landsat series since 1972 will allow relatively uniform measurement of disturbance in all time periods. Second, the specificity of plot-based estimates of harvest levels is limited by the conditions represented in the sample; harvest levels by a particular type of owner on particular slopes may only be estimated if a sufficient number of plots share those conditions. Lastly, disturbance maps resulting from this project may also be used to support purely spatial analyses for which sample-based methods are poorly suited. Healey *et al.* [2006a] looked at the size of clearcuts across time and owners (fig. 2), showing that Federal forest administrators have consistently used clearcuts that are approximately half the size of non-Federal owners. FIA plot data alone could not support this type of study. Other spatial attributes of harvests that may be of interest to FIA are proximity to streams or population centers, spatial aggregation, and edge ratio. Thus, while FIA currently has the capacity to study harvest levels at the county or State level, the use of plot and Landsat data to create harvest maps will provide significant insight into how harvests are distributed across the landscape.

Figure 1.—Harvest rates in western Oregon and Washington on USDA Forest Service, BLM, and non-Federal lands, 1972–2002. Shown is the annualized percentage of all forest land harvested using clearcut methods.



BLM = Bureau of Land Management; FS = Forest Service.
Source: Data from Healey *et al.* (2006a).

Figure 2.—Mean patch size of clearcut harvest units on USDA Forest Service, BLM, and non-Federal lands in western Oregon and Washington, 1972–2002.



BLM = Bureau of Land Management; FS = Forest Service.

Source: Data from Healey et al. 2006a.

Fire Mapping

The USDA Forest Service, through its Remote Sensing Applications Center (RSAC), currently supports two large-scale fire monitoring programs. Burned Area Reflectance Classification maps are produced at RSAC for many forest fires using pre- and post-fire Landsat images. These maps categorize reflectance differences associated with fires, and these differences are then considered along with ancillary data to produce categorical maps of fire severity. These maps are not created for all fires, however, and do not produce an explicit estimate of forest lost. RSAC's other fire-monitoring program is the MODIS Active Fire Mapping Program, a collaboration with NASA Goddard, the University of Maryland, the National Interagency Fire Center, and the Missoula Fire Sciences Lab. This project identifies likely areas of fire activity using the thermal band from the MODIS instruments on the Terra and Aqua satellites. An effort is under way to further classify these active fire maps into fire severity classes. This classification may then be used in conjunction with data from FIA plots within each severity class to create rapid characterizations of the forest types affected by each level of fire severity.

The current Landsat-based project will complement the MODIS-based efforts in that, although fire loss estimates will not be as immediate, they will have greater spatial resolution and they will be in the form of discrete predictions of biomass reduction at the pixel level. In addition, maps will be available

for fires occurring in the pre-MODIS era. The information implicit in these maps regarding the spatial distribution of fire effects may have several applications. LANDFIRE, a collaboration between the USDA Forest Service and several other Federal and private partners (www.landfire.gov), is creating maps of fuel conditions in the West using FIA data to train Landsat imagery. Fire intensity maps from the current project may be used to update LANDFIRE fuel layers. Maps of fires and other types of disturbance may likewise be used to update habitat maps (e.g., Lint 2005). Because fires can create conditions favorable for some forest pathogens (Gara 1988), maps of fire damage may also be of use in guiding forest health monitoring activities.

Storm Damage Assessment

Hurricanes and other storms can cause widespread forest damage. Storm damage, however, may be localized; differential mortality rates may result from topography, stand structure, or other factors (Millward and Kraft 2004). Although FIA currently has the capacity to estimate volume loss at the county or multicounty level, it has no way to monitor local storm effects. Spatially explicit estimates of storm damage may provide insight into storm risk at the stand level, particularly in relation to local topography. Going forward, it may also be possible to use post-storm imagery in a “rapid response” mode. In areas where change detection algorithms have already been trained with plot data and historical Landsat imagery, obtaining forest change estimates would require only the normalization of a post-storm image and the application of the existing algorithm. Although ephemeral storm effects such as standing water may somewhat reduce the accuracy of damage estimates obtained immediately after a storm, such estimates may nevertheless have value in pinpointing areas of highest damage. This information may be used to direct salvage crews or damage assessment surveys. At least 2 of the 25 Landsat scenes to be processed through this project depict major storm-affected areas (1989's Hurricane Hugo in South Carolina and the 1999 Boundary Waters wind event in Minnesota). There is substantial FIA plot data from both before and after each of these storms, allowing assessment of this project's estimates of storm damage.

Post-Disturbance Recovery

Just as this project's estimates of biomass over time may be of use in identifying disturbances, the same estimates may allow measurement of the rate of subsequent biomass accumulation. Successional recovery following disturbance can be a highly variable process, both within and across ecosystems (Yang *et al.* 2005). The consequences of slow recovery after a disturbance may include erosion (Agee 1993) and the delay of timber production. Spatially explicit recovery information, like spatial disturbance information, may be used to complement FIA plot-based estimates. Historical recovery maps may be used to update fuel and habitat maps, and they may be considered with other geographic variables to create context-dependent models of recovery. Such models may be useful to managers considering the need for or likely success of active recovery efforts following large-scale disturbances such as fires or storms. The temporal resolution of the biennial Landsat imagery used in this project may be a particular benefit in the monitoring of post-disturbance stand dynamics. The 5- or 10-year remeasurement intervals used by FIA may be less suited than biennial Landsat imagery to characterizing the potentially rapid changes (Oliver and Larson 1996) occurring after a disturbance. Thus, though FIA plot-based estimates may be used to estimate recovery rates at the county or State level, Landsat-based maps of forest recovery may add detail to our understanding of how recovery is spatially and temporally distributed.

Summary

This collaboration represents an opportunity for FIA to greatly expand the spatial information that it can provide to stakeholders and clients. An approach to forest change detection is being developed specifically to take advantage of the existing FIA database. The intrinsically spatial information resulting from this approach will complement the program's current ability to make area-based estimates of disturbance. Several applications of this spatial information in the monitoring of harvests, fires, storms, and regrowth have been suggested in this paper.

The disturbance histories of 25 sample areas across the country are now being processed using Landsat imagery and FIA data. When these initial analyses have been completed, the change detection algorithm, the critical image processing tools developed by the LEDAPS program, and the maps of disturbance and regrowth will be available to FIA for future studies.

Literature Cited

- Agee, J.K. 1993. Fire ecology of Pacific Northwest forests. Washington: Island Press. 493 p.
- Blackard, J.A.; Moisen, G.G.; Tymcio, R.P. 2006. Modeling investigations into mapping of forest attributes with FIA data for the State of Wyoming. Proceedings, sixth annual forest inventory and analysis symposium. Gen. Tech. Rep. WO-70. Washington, DC: U.S. Department of Agriculture, Forest Service. 11 p.
- Cocke, A.E.; Fule, P.Z.; Crouse, J.E. 2005. Comparison of burn severity assessments using differenced normalized burn ratio and ground data. *International Journal of Wildland Fire*. 14(2): 189-198.
- Cohen, W.B.; Fiorella, M. 1998. Comparison of methods for detecting conifer forest change with thematic mapper imagery. In: Lunetta, R.S.; Elvidge, C.D., eds. Remote sensing change detection, environmental monitoring methods and applications. Chelsea, MI: Ann Arbor Press: 89-102.
- Cohen, W.B.; Fiorella, M.; Gray, J.; Helmer, E.; Anderson, K. 1998. An efficient and accurate method for mapping forest clearcuts in the Pacific Northwest using Landsat imagery. *Photogrammetric Engineering and Remote Sensing*. 64: 293-300.
- Cohen, W.B.; Spies, T.A.; Alig, R.J.; Oetter, D.R.; Maiersperger, T.K.; Fiorella, M. 2002. Characterizing 23 years (1972-1995) of stand-replacing disturbance in western Oregon forest with Landsat imagery. *Ecosystems*. 5: 122-137.

-
- Collins, J.B.; Woodcock, C.E. 1996. An assessment of several linear change detection techniques for mapping forest mortality using multitemporal Landsat TM data. *Remote Sensing of Environment*. 56: 66-77.
- Czaplewski, R.L.; Patterson, P.L. 2001. Accuracy of remotely sensed classifications for stratification of forest and nonforest lands. In: Reams, G.A.; McRoberts, R.E.; Van Deusen, P.C., eds. *Proceedings, second annual Forest Inventory and Analysis symposium*. Gen. Tech. Rep. SRS-47. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station: 32-42.
- Edwards, T.C.; Moisen, G.G.; Cutler, R.D. 1998. Assessing map accuracy in a remotely sensed, ecoregion-scale cover map. *Remote Sensing of Environment*. 63: 73-83.
- Franklin, S.E.; Moskal, L.M.; Lavigne, M.B.; Pugh, K. 2000. Interpretation and classification of partially harvested forest stands in the Fundy model forest using multitemporal Landsat TM. *Canadian Journal of Remote Sensing*. 26: 318-333.
- Frescino, T.S.; Edwards, T.C.; Moisen, G.G. 2001. Modeling spatially explicit forest structural attributes using generalized additive models. *Journal of Vegetation Science*. 12: 15-26.
- Gara, R.I. 1988. Interactions between fires, fungi, mountain pine beetles, and lodgepole pine in south-central Oregon. *The Northwest Environmental Journal*. 4: 355-358.
- Hall, R.J.; Kruger, A.R.; Scheffer, J.; Titus, S.J.; Moore, W.C. 1989. A statistical evaluation of Landsat TM and MSS for mapping forest cutovers. *Forest Chronicle*. 65: 441-449.
- Healey, S.P.; Cohen, W.B.; Yang, Z.; Krankina, O.N. 2005. Comparison of tasseled cap-based Landsat data structures for use in forest disturbance detection. *Remote Sensing of Environment*. 97(3): 301-310.
- Healey, S.P.; Cohen, W.B.; Yang, Z.; Kennedy, R.E. 2006a. Remotely sensed data in the mapping of forest harvest patterns. In: Wulder, M.; Franklin, S., eds. *Forest disturbance and spatial patterns: remote sensing and GIS approaches*. Boca Raton, FL: CRC Press: 63-84.
- Healey, S.P.; Yang, Z.; Cohen, W.B.; Pierce, J.D. 2006b. Application of two regression-based methods to estimate the effects of partial harvest on forest structure using Landsat data. *Remote Sensing of Environment*. 101: 115-126.
- Huang, C.; Townshend, J.R.G.; Zhan, X.; Hansen, M.; DeFries, R.; Sohlberg, R. 2000. Detecting land cover changes based on their trajectories in the spectral space. In: *Proceedings, IGARSS: IEEE International Geoscience and Remote Sensing symposium*. Honolulu, HI: 61-67.
- Jin, S.; Sader, S.A. 2005. Comparison of time series tasseled cap wetness and the normalized difference moisture index in detecting forest disturbances. *Remote Sensing of Environment*. 94: 364-372.
- Lint, J.B., tech. coord. 2005. Northwest forest plan—the first ten years (1994-2003): status and trend of northern spotted owl populations and habitat. Gen. Tech. Rep. PNW-GTR-648. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 176 p.
- Lister, A.; Hoppus, M.; Riemann, R. 2004. Technique for ranking potential predictor layers for use in remote sensing analysis. In: *Proceedings, Society of American Foresters 2003 national convention*. Bethesda, MD: Society of American Foresters: 402-409.
- McMaster, K.J. 2005. Forest blowdown prediction: a correlation of remotely sensed contributing factors. *Northern Journal of Applied Forestry*. 22(1): 48-53.
- McRoberts, R.E.; Nelson, M.D.; Wendt, D.G. 2002. Stratified estimation of forest area using satellite imagery, inventory data, and the k-nearest neighbors technique. *Remote Sensing of Environment*. 82(2-3): 457-468.

-
- Millward, A.A.; Kraft, C.E. 2004. Physical influence of landscape on a large-extent ecological disturbance: the northeastern North American ice storm of 1998. *Landscape Ecology*. 19(1): 99-111.
- Moisen, G.G.; Frescino, T.S. 2002. Comparing five modelling techniques for predicting forest characteristics. *Ecological Modelling*. 157(2-3): 209-226.
- Oliver, C.D.; Larson, B.C. 1996. *Stand dynamics*, update edition. New York: John Wiley. 520 p.
- Olsson, H. 1994. Changes in satellite-measured reflectances caused by thinning cuttings in boreal forest. *Remote Sensing of Environment*. 50: 221-230.
- Patterson, P.L.; Williams, M.S. 2003. Effects of registration errors between remotely sensed and ground data on estimators of forest area. *Forest Science*. 49(1): 110-118.
- Powell, S.L. 2004. *Conifer cover increase in the Greater Yellowstone ecosystem: rates, extent, and consequences for carbon*. Bozeman, MT: Montana State University. Ph.D. dissertation. 137 p.
- Sader, S.A. 1995. Spatial characteristics of forest clearing and vegetation regrowth as detected by Landsat thematic mapper imagery. *Photogrammetric Engineering and Remote Sensing*. 61: 1145-1151.
- Sader, S.A.; Bertrand, M.; Wilson, E.H. 2003. Satellite change detection of forest harvest patterns on an industrial forest landscape. *Forest Science*. 49(3): 341-353.
- Sader, S.A.; Winne, J.C. 1992. RGB-NDVI colour composites for visualizing forest change dynamics. *International Journal of Remote Sensing*. 13(16): 3055-3067.
- Skakun, R.S.; Wulder, M.A.; Franklin, S.E. 2003. Sensitivity of the thematic mapper enhanced wetness difference index to detect mountain pine beetle red-attack damage. *Remote Sensing of Environment*. 86: 433-443.
- Yang, Z.; Cohen, W.B.; Harmon, M.E. 2005. Patterns of early forest succession following clearcuts in western Oregon. *Canadian Journal of Forest Research*. 35: 1889-1900.

United States
Department of
Agriculture

Forest Service

Gen. Tech. Report
WO-77

August 2007



Proceedings of the Seventh Annual Forest Inventory and Analysis Symposium

Edited by

Ronald E. McRoberts

Gregory A. Reams

Paul C. Van Deusen

William H. McWilliams

Portland, ME

October 3–6, 2005

Sponsored by

USDA Forest Service, National Council for Air and Stream Improvement,
Society of American Foresters, International Union of Forestry Research
Organizations, and Section on Statistics and the Environment of the American
Statistical Association

McRoberts, Ronald E.; Reams, Gregory A.; Van Deusen, Paul C.; McWilliams, William H., eds. 2007. Proceedings of the seventh annual forest inventory and analysis symposium; 2005 October 3–6; Portland, ME. Gen. Tech. Report WO-77. Washington, DC: U.S. Department of Agriculture Forest Service. 319 p.

Documents contributions to forest inventory in the areas of sampling, remote sensing, modeling, information management and analysis for the Forest Inventory and Analysis program of the USDA Forest Service.

KEY WORDS: Sampling, estimation, remote sensing, modeling, information science, policy, analysis

Disclaimer

Papers published in these proceedings were submitted by authors in electronic media. Editing was done to ensure a consistent format. Authors are responsible for content and accuracy of their individual papers. The views and opinions expressed in this report are those of the presenters and authors and do not necessarily reflect the policies and opinions of the U.S. Department of Agriculture. The use of trade or firm names in this publication is for reader information and does not imply endorsement by the U.S. Department of Agriculture of any product or service.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.