Liane R. Davis^{1,2} and Klaus J. Puettmann, Department of Forest Science, Oregon State University, Corvallis, Oregon 97331

and

Gabriel F. Tucker, Castor International, 1604 8th Avenue S.W., Olympia, Washington 98502

Overstory Response to Alternative Thinning Treatments in Young Douglas-fir Forests of Western Oregon

Abstract

An increase in land dominated by young second-growth Douglas-fir forests in the Pacific Northwest has coincided with heightened concerns over loss of old-growth habitat. In search of options for managing young forests to provide late-successional forest structures, the Young Stand Thinning and Diversity Study was designed to test the effectiveness of modified thinning in acceleration of late-successional structural characteristics. Thinning treatments included: a control, a light thin (typical of standard commercial thins), a heavy thin (densities lower than typically prescribed), and a light thin with gaps (stands thinned lightly with the addition of 0.2 hectare patch cuts evenly spaced throughout the stand). Early response (maximum of 5-7 years post-treatment) of overstory vegetation was examined. Average growth of Douglas-fir increased in all thinned stands, but growth of the largest Douglas-fir trees was only accelerated in the heavy thin. After thinning, the canopy of all thinned treatments was initially more open than the control, but after 5-7 years the light thin was no longer significantly different from the control. The light with gaps thin had the highest variation in overstory canopy cover. Differentiation of vertical canopy structure among treatments was not evident. There was no difference in mortality among any of the treatments for most species tested; those that did had highest mortality in the control. Our results indicate that thinning can be effective in hastening development of some, but not all late-successional attributes, but such acceleration is not equivalent among the different thinning treatments.

Introduction

During recent decades, young managed forests have become a dominant feature in the Pacific Northwest landscape, especially in western Oregon and Washington. Mainly composed of planted Douglas-fir (Pseudotsuga menziesii) trees between the ages of 30-50 years, these stands have often replaced what was once dominated by late-successional or "old-growth" habitat (Bolsinger and Waddell 1993). These young stands often lack structural characteristics of old-growth forests, such as large living trees, snags, and a multilayered canopy that includes an abundant and heterogeneous understory (Franklin and Spies 1991a, 1991b; Spies 1991; Spies and Franklin 1991; Halpern and Spies 1995; Franklin et al. 2002). Without this suite of structural attributes, young stands may not provide the variety of habitats necessary to support a high diversity and abundance of native species (Spies 1991, Halpern and Spies 1995).

²Current Address: The Nature Conservancy,

The management of young stands in order to promote late-successional habitat has remained a topic of considerable debate. In response to this debate, the Young Stand Thinning and Diversity Study (YSTDS) was initiated in 1994. As a comprehensive and integrated long-term ecological study, the YSTDS was designed to test the efficacy of thinning young stands to accelerate development of late-successional habitat. Though retrospective studies have investigated whether thinning promotes late-successional habitat (Bailey et al. 1998, Thomas et al. 1999, Thysell and Carey 2000), few studies have implemented thinning with the intent of ecological enhancement (Thysell and Carey 2001). This paper examines short-term responses to thinning treatments designed to accelerate latesuccessional habitat characteristics.

By opening the canopy and releasing resources, thinning promotes growth of remaining overstory trees and establishment of a prominent understory layer, thereby adding complexity to these young stands and perhaps accelerating development of latesuccessional habitat (Muir et al. 2002). Enhancing stand complexity results in increased microhabitat heterogeneity and improvement of habitat suitability for many organisms (Carey and Johnson 1995,

Northwest Science, Vol. 81, No. 1, 2007 © 2007 by the Northwest Scientific Association. All rights reserved.

¹Author to whom correspondence should be addressed. Email: ldavis@tnc.org

⁷⁵⁰ Commercial St. #212, Astoria, OR 97130

Hagar et al. 1996, Bailey and Tappeiner 1998, Rambo and Muir 1998, Carey and Harrington 2001, Hagar et al. 2004). Over time, development of large trees and snags combined with a layered canopy and understory may result in thinned stands becoming more similar to old-growth faster than unthinned stands (Muir et al. 2002).

Thinning has not traditionally been used to achieve ecological objectives. It has mainly been a tool for enhancing timber production (for examples see the Level-of-Growing-Stock studies; Marshall and Curtis 2002). Thinning treatments implemented in this study were similar to traditional low thinnings used for timber production, but were modified to address the goal of accelerating several late-successional stand attributes. These attributes include tree species diversity and initial low density conditions of old-growth stand development (Tappeiner et al. 1997, Poage and Tappeiner 2002), and spatial diversity due to small-scale mortality patterns (Franklin and van Pelt 2004).

It is not known if these thinning prescriptions will be successful in acceleration of all late-successional features. For example, accelerating development of the dominant overstory component (Franklin and Spies 1991a) requires dominant trees to increase diameter and height growth following thinning (Staebler 1956, Miller and Williamson 1974, Oliver and Murray 1983). However, the intensity at which thinning reduces competition among the largest trees to accelerate their growth is uncertain. It is also not known whether thinning to very low densities results in high mortality from windthrow or other agents. While thinning may eventually promote establishment of a multilayered canopy by encouraging crown extension and understory release and regeneration (Bailey et al. 1998), in the short-term low thinning may simplify crown structure by removing many of the suppressed and intermediate trees (Smith et al. 1997). In addition, some conifer and hardwood species may be adapted to a shaded understory and experience high mortality upon canopy removal or as a result of harvest damage (Tucker and Emmingham 1977, Tucker et al. 1987).

To address these questions, the overall objective of this study was to characterize early overstory response following alternative thinning treatments in young Douglas-fir stands. Specifically, this includes a 5-7 year post-thinning comparison among four treatments of: (1) overstory cover (2) vertical crown structure (3) growth of all Douglas-fir trees

2 Davis, Puettmann, and Tucker

(4) growth of the largest Douglas-fir trees that may eventually provide the dominant component of late-successional habitat in the stand matrix and near gaps, and (5) differences in mortality of individual tree species due to stand instability, harvesting damage, and competition.

Methods

Study Design and Description

The study is a randomized block design comprised of four blocks with each block containing one replication of four treatments. Study blocks are designated as: Cougar Reservoir (CR), Christy Flats (CF), Sidewalk Creek (SC), and Mill Creek (MC). Blocks were selected for homogeneity in overstory composition, stand age, similar management history, and size (> 56 ha). All blocks are located in the Willamette National Forest on the western slope of the Cascade Range of Oregon (400 to 900 m elevation) and are composed of 40-50 year old (at the time of study initiation) planted Douglas-fir stands occurring within the Western Hemlock (Tsuga heterophylla) zone (Franklin and Dyrness 1973). Mean annual precipitation is 230 cm, with only 5% falling between July and October. The average yearly temperature is 10.1°C. Soils are generally well developed, ranging from thin silty loams/clay loams to thin gravelly loams.

Within each block, each of the four treatments were assigned randomly to a treatment unit, providing a total of 16 treatment units. Treatment units range from 15 to 53 ha in size, and have varied slope and aspects (Table 1). Within a block, treatment units were selected for homogeneity in size, elevation, slope, aspect, site index, soil type, and dominant plant association (Table 1).

In addition to Douglas-fir, stands contained a small component of other conifer (e.g., western hemlock) and hardwood (e.g., bigleaf maple (*Acer macrophyllum*)) species. Baseline pre-treatment stand exam data were collected in 1993 and showed pre-treatment basal areas (BA) and densities (trees per hectare (tph)) were comparable among treatment units within blocks (Table 2).

Treatment Description

The four treatments in each block are: Control, Light thin, Heavy thin, and Light with Gaps thin (hereafter abbreviated as C, LT, HT, and LG, re-

Block	Treatment	Area (Ha)	Age	Elev. (m)	Slope (%)	Aspect	Site Index ¹	Dominant Plant Association ²	Total# of Plots
CR	С	30.0	40	805	18.8	Е	107	Tshe/Gash	23
CR	HT	19.4	40	792	24	Е	105	Tshe/Bene	13
CR	LT	37.2	38	610	17.1	E	107	Tshe/Bene	26
CR	LG	14.6	40	792	16	ENE	105	Tshe/Bene	29 ³
MC	С	52.6	42	902	21.1	SSEE	105	Tshe/Bene	25
MC	HT	34.8	42	658	22.9	SE	105	Tshe/Bene	23
MC	LT	37.2	43	524	20	S	105	Tshe/Bene	30
MC	LG	19.8	42	439	8.9	SSW	106	Tshe/Bene	29 ³
CF	С	30.8	39	878	6.2	SE	117	Tshe/Bene	23
CF	HT	20.2	36	905	0	SE	120	Tshe/Bene	15
CF	LT	32.0	39	902	5.3	SE	117	Tshe/Bene	24
CF	LG	38.9	40	905	5.3	SSEE	118	Tshe/Bene	30
SC	С	51.0	37	634	11.4	N	114	Tshe/Rhma-Gash	17
SC	HT	19.0	35	652	16	NW	115	Tshe/Rhma-Gash	13
SC	LT	22.3	33	646	21.8	NNE	122	Tshe/Rhma-Gash	15
SC	LG	30.4	39	671	14.5	Ν	111	Tshe/Rhma-Gash	30

TABLE 1. Site characteristics for each treatment unit. (CR = Cougar Reservoir Block; MC = Mill Creek Block; CF = Christy Flats Block; SC = Sidewalk Creek Block).

Dominant tree height at 50 years, King's Site Index Tables

²Tshe = western hemlock; Bene = Oregon grape (*Berberis nervosa*); Rhma = western rhododendron (*Rhododendron macrophyllum*); Gash = salal (*Gaultheria shallon*)

³One stand matrix plot was removed from study because it was located out of study area.

TABLE 2. Description of thinning treatments.

х.,		Pre-treatment Density ¹	Post-treatment Density ¹	Pre-treatment BA1	Post- treatment BA1
Block	Treatment	(tph)	(tph)	(m² / ha)	(m^2 / ha)
CR	С	929	753	30	53
CR	HT	800	151	27	14
CR	LT	865	312	39	21
CR	LG	891	221	36	19
MC	С	402	655	35	40
MC	HT	466	283	36	13
MC	LT	339	415	40	25
MC	LG	335	346	36	18
CF	С	737	869	42	47
CF	HT	880	133	48	21
CF	LT	871	207	39	32
CF	LG	855	198	40	27
SC	С	756	792	28	39
SC	HT	820	165	25	12
SC	LT	800	277	26	20
SC	LG	743	225	30	15

¹Pre-treatment measures include all trees \geq 5 cm dbh. Post-treatment measures include all trees \geq 8 cm dbh. Pre-treatment data was also sampled differently from post-treatment data. Therefore, these numbers are not presented in order to make direct pre-/post-treatment comparisons, but to illustrate similarities of pre-treatment conditions within each block.

spectively). The LT reflected current conventional thinning for timber production, while the HT and

LG aimed at increasing understory vegetation by further reducing overstory densities using

Alternative Thinning in Douglas-fir 3

different spatial arrangements. Residual target densities (tph) for the thinning prescriptions were: C = unthinned (approximately 650 tph); LT =250-300 tph; HT = 125 tph; and LG = 250-300tph with additional cutting of 0.2 hectare circular gaps evenly dispersed every 2 ha (Table 2). Final post-treatment basal areas were close to those of the prescriptions (Table 2). Areas within the LG are stratified into 3 sub-treatments: (1) Gap: 0.2 ha gap; (2) Edge: a doughnut-shaped area surrounding the gap; (3) and Stand Matrix: the remainder of the treatment unit (Figure 1; hereafter abbreviated as G, E, and SM, respectively). More information on theses sub-treatments is provided under Sampling Methods.

Treatments were applied between 1995-1997. Due to the large treatment size, buffering between treatments was not always possible, but treatments were occasionally separated by roads or other terrain barriers. At the CR and MC blocks, thinning was done with a combination of tractor and skyline systems. A ground-based harvester and forwarder system was used at the CF block and a skyline system was used at the SC block.

All thinning treatments used a low thinning prescription with the added objective to leave species other than Douglas-fir. The C provided a reference for stand development without management intervention. The LT was similar to a typical commercial thin commonly used throughout western Oregon except for the retention of species other than Douglas-fir to encourage species mix. The HT opened the canopy substantially more than common commercial thins and reflected recent findings that some old-growth stands initiated at very low densities (Tappeiner et al. 1997). The LG was intended to provide spatial diversity by simulating gap-phase mortality and created open patches within a stand matrix thinned to the same density as the LT.

Sampling Methods

"First-year" post-thinning vegetation sampling occurred in the summer of 1995-1997, depending on the time of harvest completion. In most cases,



Figure 1. Schematic of sampling design in LG treatment.

4 Davis, Puettmann, and Tucker

this was the first growing season following harvest (for more information on harvest and sampling schedule refer to Beggs 2004). Resampling was completed during the summers of 1999 and again in 2001, depicting vegetation response 3-5 growing seasons and 5-7 growing seasons post-harvest. For ease of communication, these data will be used to depict "first-year", "third-year", and "fifth-year" post-treatment vegetation response, respectively.

Sampling was done using 0.1 ha (17.84 m radius) circular permanent plots randomly located along transects that were systematically placed through treatment units. In C, LT, and HT treatment units, approximately 7.5 % of the area was sampled (see Table1 for total plot numbers).

Sampling for each LG treatment unit used 30 plots in order to capture variation among the three sub-treatments. For each treatment unit, 10 gaps and 10 edges were randomly selected and one plot was placed in each. Gap plots were centered in the gap, allowing only the gap interior to be sampled by these plots. Edge plots were centered in a random direction at 35.7 m from gap center so that each edge plot extended 7.5 m into the gap and 28.2 m into the remainder of the stand. For each gap, only one plot was permited to be placed in the surrounding edge. Ten plots were also located in the stand matrix. These plots were randomly placed throughout the remainder of the treatment unit with the criteria that plot center was located at least 71.4 m from the center of any gap, permitting sampling of all area at least 53.5 m from gap center (Figure 1).

Within each plot, overstory cover was measured at plot center and at four locations 10.25 m from plot center in each cardinal direction using a "moosehorn" densiometer (Cook et al. 1995). Overstory cover included live foliage and tree bole, limbs, and snags. Diameter at breast height was also measured for of all trees ≥ 5 cm dbh in each plot. Overstory cover and dbh were measured during each year of sampling. In addition, a random subsample of trees in each treatment unit that were measured for dbh was also measured for height and crown length in 1999. Another random subsample of trees in each treatment unit that were selected for height and crown measurements was cored at breast height in 1999 to determine stand age.

Overstory Cover

With the exception of the LG treatment units, average overstory cover of each treatment unit was calculated by averaging the plot means. In the LG, the number of plots in each sub-treatment (G, E, and SM) was the same; however, each subtreatment did not occupy an equal proportion of the total treatment unit area. To adjust for this, a weighted average of sub-treatment means was used to calculate the treatment unit means. Weights for each sub-treatment were based on the proportion of areas in each sub-treatment to total treatment unit areas. The coefficient of variation (CV), an indicator of variation in overstory cover, was used to compare variation in overstory cover among treatments. The variation of cover was illustrated by comparing frequency diagrams of overstory cover in each treatment.

Vertical Canopy Structure

Assessment of vertical canopy structure required heights and crown lengths for all trees within the sample of the study. Because these measurements were only available for a subsample of the data, height and crown length were predicted for the remaining trees using species-specific non-linear regression. We used equation forms from Hanus et al. (1999) and Ritchie and Hann (1987), but estimated parameters from trees sampled in this study. Because of small sample size concerns, a species was combined with the species most closely resembling its growth pattern when fewer than 10 trees of the species were available. Therefore, grand fir (Abies grandis) was combined with Douglasfir; mountain hemlock (Tsuga mertensiana) was combined with western hemlock; Oregon ash (Fraxinus latifolia), cascara buckthorn (Rhamnus purshiana), and willow spp. (Salix spp.) were combined with bigleaf maple. The addition of these minor species did not affect parameter estimates for the dominant species.

These estimates were then used to calculate live crown ratios (LCR) and the foliage height diversity (FHD) index (MacArthur and MacArthur 1961) for each treatment unit. LCR gauges vertical length of the crown relative to tree height and assumes continuous vegetation throughout the entirety of the crown. The FHD index assesses diversity of vertical distribution of foliage using two components: richness and evenness, similar to the Shannon-Weiner diversity index. Richness, in

Alternative Thinning in Douglas-fir

this case, is the number of 5-meter layers occupied by tree crowns in the stand. A 5-meter interval was selected because smaller intervals would not have compensated for the error incurred with estimations of height and crown length measures. Evenness is the relative abundance of tree crowns within these intervals. Like the Shannon-Weiner index, FHD can be strongly influenced by unbalanced dominance of richness or evenness (Hill 1973). To examine if richness or evenness was controlling FHD, both were tested separately.

Diameter Growth

Annual adjusted diameter growth between firstyear and fifth-year post-thinning of all Douglas-fir trees \geq 5 cm dbh was compared among treatments. To calculate annual adjusted diameter growth (expressed as % increase from initial dbh), absolute diameter growth (cm/yr) was first calculated by subtracting first-year dbh from fifth-year dbh and dividing by the number of years between measurements, accounting for offsets in timing of first-year measurements. Annual adjusted diameter growth was then calculated by dividing absolute diameter growth by the first-year dbh and multiplying by 100. Adjusted diameter growth was used because absolute diameter growth did not account for initial differences in dbh among treatments that were an artifact of thinning (thinning removes smaller trees, inherently increasing average dbh in thinned treatments relative to the unthinned treatment regardless of differences in growth).

To specifically assess response of the largest Douglas-fir trees, i.e., trees that likely will make up the dominant stand structure, absolute diameter growth between first-year and fifth-year post-thinning of trees with the largest dbh was compared among treatments. This was done for the largest 10, 15, 20, 25, and 30 tph in order to simulate a range of large-tree densities typical of old-growth stands (Franklin and Spies 1991a). To examine the largest 10 tph, the largest tree from each 0.1 ha plot was selected. Likewise, the 2 largest trees were selected from each 0.1 ha plot to examine 20 tph, and the largest 3 for 30 tph. For the intermediate 15 and 25 tph, half of the plots were randomly selected and the largest 2 or 3 trees, respectively, from these plots were combined with the largest 1 or 2 trees, respectively, from all plots. Absolute diameter growth was calculated as previously described and averaged for each treatment unit. Absolute diameter growth was used instead of

6 Davis, Puettmann, and Tucker

adjusted diameter growth because average initial dbh of the largest trees did not differ among treatments.

In the LG, no Douglas-fir were present in the gaps. Thus, growth is representative only of trees in the SM and E. To evaluate potential effects of the gaps on tree development, adjusted diameter growth of all Douglas-fir trees ≥ 5 cm dbh and absolute diameter growth of the largest 10, 20, and 30 tph of Douglas-fir trees were compared between the E and SM sub-treatments. Calculations were done as previously described.

Mortality

Mortality of individual tree species as well as all hardwood species combined was compared among treatments. In each treatment unit, percent mortality of each species was computed between first-year and fifth-year sampling. Percent mortality was calculated for trees ≥ 5 cm dbh and for small trees (trees between 5 –10 cm dbh) in order to determine (a) if species experienced differences in mortality among treatments and (b) if a difference in mortality was limited to small trees, i.e., competition related. Due to concerns about small sample sizes, mortality for a species was only assessed if more than 10 trees of that species were present in each treatment during first and fifth-year sampling.

For all trees \geq 5 cm dbh, the following species were tested for differences in mortality among all treatments: bigleaf maple, golden chinquapin (Chrysolepis chrysophylla), Pacific dogwood (Cornus nuttallii), bitter cherry (Prunus emarginata), Douglas-fir, Pacific yew (Taxus brevifolia), western red cedar (Thuja plicata), and western hemlock. Fewer than 10 red alder (Alnus rubra) and incense cedar (Calocedrus decurrens) were observed in the HT, so comparisons of mortality were only made among the LT, LG, and C. The following species did not have enough trees in any treatment to permit separate comparisons and were only included in the combined hardwood analysis: Pacific madrone (Arbutus menziesii), Oregon ash, black cottonwood (Populus trichocarpa), cascara buckthorn, and willow species.

For small trees, bitter cherry had too few trees to permit separate treatment comparisons (in addition to the species listed above) and golden chinquapin did not have enough trees in the LT, so comparisons were only made among the C, HT, and LG. These species were still included in the combined hardwood analysis.

Data Analysis

All analyses were done using SAS v. 8.2 statistical software (SAS Institute 2001). Comparisons among treatments were performed with ANOVA using a randomized complete block model. The Tukey-Kramer adjustment was used for all multiple comparisons. Prior to ANOVA, all data were checked for normality. Small sample sizes (n =16 for comparisons among treatments; n = 12 for LG sub-treatment comparisons) made assessment of normal distributions difficult; however, under the central limit theorem, such distributions approach normality (Thysell and Carey 2000). Therefore, no transformations were performed. The significance level for all analyses was set at $P \le 0.05$, and $P \le 0.10$ was considered to be marginally significant.

For mortality comparisons, several species were more frequently observed in the C than thinned treatments, resulting in more precise mortality estimates of these species in the C. To account for decreased precision in thinned treatments, a weighted ANOVA was used to weigh average mortality by total trees (for each species) in each treatment.

Time trends in overstory cover were investigated using a repeated measures analysis. A time * treatment interaction was used to test whether changes in overstory cover were equal among treatments over time.

Results

Overstory Cover

As expected, thinning opened up the overstory canopy. During the first and third years post-thinning, all thinned treatments had less overstory cover than C (Figure 2; Year 1: P < 0.001 for HT,



Figure 2. Overstory cover over time (includes 90% confidence intervals). Letters indicate significant differences among treatments (treatments with same letter do not differ at $P \le 0.05$ level).

Alternative Thinning in Douglas-fir

LT, and LG; Year 3: P < 0.001 for HT and LG, P = 0.010 for LT). The thinning treatments differed, as HT had less overstory cover than the LT (Year 1: P = 0.004; Year 3: P = 0.005). Cover in the LG did not significantly differ from the LT (P > 0.100for both years) but was marginally higher than the HT (Year 1: P = 0.076; Year 3: P = 0.084). The trends in overstory canopy development differed among treatments (Time*Treatment interaction: P = 0.021; Figure 2) and five years after thinning, the HT and LG still had less average cover than the C (HT: P < 0.001; LG: P = 0.003), but no longer differed from each other (P = 0.121). Also, the LT was no longer significantly different from the C (P = 0.103; Figure 2). Within the LG, overstory cover was significantly less in the G than the E and SM during all years of measurement (P < 0.001for all years) while the E and SM did not differ from each other (P > 0.100 for all years).

Gap creation increased variation in overstory cover distribution. During the first year post-thinning, variation of overstory cover throughout the stand differed among all four treatments (P < 0.001). The C had the least amount of variation while the LG had the most. By third-year post-thinning, however, variation in the LT no longer differed from the C (P = 0.214). The HT and LG remained more variable than the C (P < 0.001 for HT and LG), and the LT (HT: P = 0.014; LG: P < 0.001) with variation in the LG being greater than variation in the HT (P = 0.006). These results did not change by the fifth-year post-thinning. While other treatments displayed normal distributions (for illustrations, see Beggs 2004), the greater variation in the LG is primarily a result of low overstory cover in the gaps and intermediate cover through the remainder of the stand (Figure 3).

Vertical Structure

Differentiation of crown layers did not seem to be significantly impacted by thinning. By third-year post thinning, FHD and LCR did not differ among treatments (P = 0.85 and P = 0.26, respectively; Table 3). No difference among treatments was found in richness or evenness (P = 0.783 and P = 0.473, respectively) of canopy layers.



Figure 3. Frequency distribution of overstory cover for LG treatment.

8 Davis, Puettmann, and Tucker

Treatment	FHD		LCR	ADG (% increase from initial dbh)	AG of Largest 15 tph (cm/yr)
С	1.57 ª (1.51 - 1.63)	-	0.53 ª (0.50 - 0.55)	1.3 ª (1.0 - 1.6)	0.70 ª (0.61 - 0.78)
HT	1.58 ª (1.52 - 1.64)		0.55 ° (0.52 - 0.58)	2.4 ^b (2.1 - 2.7)	0.94 ^b (0.84 - 1.0)
LT	1.57 ª (1.51 - 1.63)		0.51 ª (0.49 - 0.54)	1.8 ª (1.5 - 2.1)	0.78 ª (0.66 - 0.84)
LG	1.60 ª (1.54 - 1.65)		0.52 ª (0.50 - 0.55)	1.9 ª (1.6 - 2.3)	0.85 ª (0.72 - 0.89)
NOVA P-value ¹	0.854	٣	0.200	0.003	0.033

TABLE 3. Overstory Results: (1) FHD = Foliage Height Diversity Index; (2) LCR = Compacted Live Crown Ratio; (3) ADG = adjusted diameter growth of all Douglas-fir ≥ 5 cm dbh; (4) AG = absolute diameter growth of largest 15 tph Douglas-fir trees (results did not differ for ≥ 15 tph). Numbers in parentheses provide 90% confidence intervals of estimates.

¹P-value is for overall test of difference among treatment.

Diameter Growth

Heavy thinning was effective in increasing adjusted diameter growth of Douglas-fir. Adjusted diameter growth of Douglas-fir trees ≥ 5 cm dbh differed among treatments by the time of fifth-year measurement (P = 0.003; Table 3), but only the HT had significantly higher adjusted diameter growth than the C (P = 0.002). The increase in adjusted diameter growth in the LG compared to the C (P = 0.060) and in the HT compared to the LT (P = 0.057) was marginally significant. Within the LG, adjusted diameter growth in the E was slightly, but marginally significantly higher than in the SM (2.0 vs. 1.8, respectively; P = 0.064).

Large Douglas-firs also responded to improved growing conditions provided by heavy thinning. For the largest 15 tph, absolute growth was higher in the HT than in the C (P = 0.027) but did not differ between the LT and C (P = 0.854) or the LG and C (P = 0.405). Results were the same for the largest 20, 25, and 30 tph. For the largest 10 tph, absolute growth of the largest Douglas-fir trees did not differ significantly among any treatments (P = 0.228). Within the LG, absolute growth of the largest Douglas-firs did not differ between the E and the SM (10 tph: P = 0.648; 20 tph: P =0.460; 30 tph: P = 0.305).

Mortality

Tree mortality was reduced following thinning. For trees ≥ 5 cm dbh and small trees, mortality of Douglas-fir and combined hardwoods was higher in the C than all other treatments (Table 4). Within the LG, neither Douglas-fir nor combined hardwood species differed in mortality among sub-treatments (P > 0.150; comparisons only between E and SM sub-treamtens for Douglas-fir; for all combined hardwoods, comparisons made with and without G sub-treatment). The same pattern held for golden chinquapin ≥ 5 cm dbh except the difference between the C and LT was not significant (Table 4). Within the LG, mortality of golden chinquapin did not differ among subtreatments (P > 0.200; comparisons made with and without G sub-treatment). None of the other species tested showed significant differences in mortality among treatments.

Discussion

Our results indicated that thinning can place young managed stands on a trajectory to develop several late-successional stand attributes, such as large diameter trees. Other attributes, such as a diversified crown structure, were not rapidly accelerated by thinning; but favorable conditions for eventual development of such structure were created. In addition, leaving tree species other than Douglas-fir was effective in maintaining these species in the overstory and prevented initial simplification of canopy structure.

The thinning treatments differed in terms of their impact on overstory cover, and thus associated characteristics. Light thinning, similar to current timber management practices, did not maintain canopy opening beyond three years. Substantial reduction in stand density, like that of the HT treatment, is necessary to ensure canopy opening is maintained for several years. Open

Alternative Thinning in Douglas-fir

	Treatment	Douglas-fir	Golden chinquapin	Combined hardwoods
All Trees (≥ 5 cm d				
	С	14.0 ^a	27.1 ª	36.1 ª
		(12.3 - 15.8)	(19.4 - 34.9)	(28.4 - 43.8)
	HT	4.7 b	7.8 ^b	18.0 ^b
		(0.1 - 9.3)	(-1.1 - 16.7)	(8.9 - 27.0)
	LT	5.9 b	15.1 ab	15.3 b
		(3.1 - 8.8)	(1.5 - 28.8)	(9.0 - 21.6)
	LG	4.0 ^b	4.4 ^b	13.4 ^b
		(0.8 - 7.2)	(-3.6 - 12.5)	(7.0 - 19.7)
	ANOVA P-value ¹	< 0.001	0.010	0.001
Small Trees (dbh = $5 - 10$ cm)	С	43.9 a	27.1 a	43.3 ª
		(39.7 - 48.2)	(-25.8 - 80.0)	(33.7 - 52.8)
	HT	18.3 b	14.9 ª	22.5 ª
		(5.4 - 31.2)	(-49.8 - 79.5)	(11.2 - 33.8)
		(13.9 - 33.1)		(16.4 - 36.9)
	LG	14.7 ^b	14.6 ª	16.8 ^b
	ANOVA P-value ¹	< 0.001	0.781	0.004

TABLE 4. Comparisons of mortality (%) among treatments. Results only reported for species with significant differences among treatments. Numbers in parentheses provide 90% confidence intervals of estimates.

¹*P*-value is for overall test of difference among treatments

canopy conditions permit more light to reach the forest floor (Parker et al. 2001) and, with fewer trees transpiring, generally increase soil moisture (Everett and Sharrow 1985). This can result in enhanced development of understory shrubs and herbs and associated wildlife habitat (e.g., Alaback and Herman 1988, Thysell and Carey 2001, Hagar et al. 2004).

A concern with evenly spaced thinnings is that the uniformly open canopy will encourage a homogenous understory dominated by a few species instead of a patchy and heterogeneous understory (Tappeiner and Zasada 1993, Huffman and Tappeiner 1997, Thysell and Carey 2000, but see Thomas et al. 1999). Thinnings in this study increased variation in overstory cover relative to unthinned stands, especially when gaps were added. Similar to canopy openness, this effect disappeared within three years for the LT treatment, suggesting that traditional thinning may have limited or possibly even undesirable impacts on understory development. Variation in canopy cover following heavier thinnings or thinnings with gaps may prevent homogeneous dominance of a few understory species by ensuring uneven distribution of light (Franklin and Van Pelt 2004). Work currently underway by the authors on the understory vegetation response to the thinning treatments will examine these patterns in more detail.

Thinning treatments differed in their ability to accelerate diameter growth, important to development of the large-tree component of late-successional structure in west-side Douglas-fir forests of the PNW. The influence of residual density on overall tree growth was obvious, as the HT, which resulted in residual densities lower than most conventional thinnings, consistently reduced competition enough to permit residual trees to capitalize on elevated resource availability and increase their diameter growth (Oliver and Murray 1983, Marshall and Curtis 2002). If the difference in growth rates continues, development of large diameter trees will occur faster in heavily thinned stands than unthinned stands. Because these large trees are valuable nest sites for northern spotted owls (Forsman et al. 1984), provide substrate for several epiphytic species (Clement and Shaw 1999), and may eventually become the large snags and downed logs essential to several wildlife species (Hayes et al. 1997), acceleration of their development is key in acceleration of late-successional habitat and structure.

On the other hand, our study indicates that, despite reduction in stand density, competition among the largest trees in the LT (i.e., commonly used thinning regimes), remains too high for these trees to substantially increase their growth (Staebler 1956). However, when this thinning prescription is combined with gap creation, trees bordering gaps

10 Davis, Puettmann, and Tucker

seem to benefit from released resources (Gray et al. 2002) and experience slightly elevated growth. Five years may not be sufficient for trees to build up a crown and the increased resource availability may not be expressed in terms of diameter growth. Thus, the trend detectable for the edge trees may strengthen over time (Staebler 1956, Oliver and Murray 1983).

The very largest trees (10 tph) did not experience a release, even in the HT, indicating that they may already have been in a dominant position with minor competition from neighbors (Staebler 1956, Oliver and Murray 1983, D'Amato and Puettmann 2004). While more intense than traditional thinning, the residual density of the HT was still higher than densities at which some old growth stands may have initiated (Tappeiner et al. 1997, Poage and Tappeiner 2002, but see Winter et al. 2002). This seems to suggest that in some instances even more intensive thinnings may be necessary or desirable in order to accelerate growth of the largest trees. Other concerns, such as wind stability, may prohibit very open conditions or heavy thinning operations. In these cases variable density thinnings, which focus on reducing competition to the largest trees, may be more appropriate.

Amplified growth of non-dominant trees and understory vegetation may eventually strain resource availability, especially on drier sites (Messier and Mitchell 1994, Bennett et al. 2003). Other work, however, has shown that dominant trees in young stands tend to contribute the largest proportion of stand production up to the point of stand closure at which time dominance begins to diminish and smaller trees contribute proportionately more to stand growth (Binkley et al. 2003, Binkley 2004). It is hypothesized that when this dominance begins to relax, the trees are so large that their growth no longer balances with their greater control of resources (Binkley 2004). At this point, assuming the largest trees we investigated are dominant in our stands, the trees will be sufficiently large to support a variety of wildlife and other species.

Despite differences in growth and overstory cover, vertical structure was not affected by thinning in the short term. A multi-tiered canopy capable of providing a diversified microhabitat, like that common in old-growth stands (Franklin and Spies 1991a), has not yet begun to develop in these stands. Given the early post-treatment

response observed (2 to 4 years after thinning), however, significant changes in crown extension and epicormic branching were unlikely. Maybe more importantly, our results indicate that overstory crown structure in the thinned stands was not simplified, as commonly predicted in low thinnings (Smith et al. 1997). Leaving tree species other than Douglas-fir ensured that the lower layers, which are removed during a conventional low thin, were maintained. Retention of these other species also maintained valuable ecological components, such as hardwoods (Hagar et al. 1996, Rambo and Muir 1998, Rosso 2000). A mixture of overstory species that includes hardwoods can often support several species assemblages better than a forest lacking diversity in overstory composition (Hagar et al. 1996, Hayes et al. 1997, Rosso 2000). Bird species, such as the warbling vireo (Vireo gilvus), are related to hardwood presence (Hagar et al. 1996). Hardwood's also generally provide more nutrient rich leachate than conifers, making them important to forest floor bryophyte species that obtain the majority of their nutrients from leachate (Rambo and Muir 1998).

Concerns about loss of tree species other than Douglas-fir due to increased mortality in thinned stands were not warranted. Most mortality was related to competition, as mortality of Douglas-fir, golden chinquapin, and all combined hardwood species was higher in unthinned stands relative to thinned stands. The dense conditions of unthinned stands probably inflicted extreme competition for resources upon suppressed and intermediate trees, resulting in high mortality (Oliver and Larson 1996, Franklin et al. 2002). By removing several intermediate and suppressed trees, thinning likely relaxed resource competition among remaining trees, thereby decreasing mortality (Oliver and Larson 1996, Marshall and Curtis 2002). On the other hand, heavier thinnings did not result in unstable stand conditions over the measurement period.

In summary, thinning of the type implemented in this study can be an effective way of increasing complexity in young, managed stands. However, applying traditional thinning practices are likely not sufficient to accelerate the development of late successional structures. Instead, thinning practices need to be "customized" in terms of the specific structural components (e.g., large trees, diverse understory light conditions) that are desired.

Alternative Thinning in Douglas-fir 1

Acknowledgements

This study was established by John Tappeiner, Loren Kellogg, Brenda McComb, John Cissel, and James Mayo. Funding and cooperation comes from the USDA Forest Service Pacific Northwest Research Station, Willamette National Forest, Bureau of Land Management, Oregon State University, University of Oregon, and Cascade Center for Ecosystem Management. We would like to thank

Literature Cited

- Alaback, P. B., and F. R. Herman. 1988. Long-term response of understory vegetation to stand density in *Picea-Tsuga* forests. Canadian Journal of Forest Research 18:1522-1530.
- Bailey, J. D., C. Mayrsohn, P. S. Doescher, E. St Pierre, and J. C. Tappeiner. 1998. Understory vegetation in old and young Douglas-fir forests of western Oregon. Forest Ecology and Management 112:289-302.
- Bailey, J. D., and J. C. Tappeiner. 1998. Effects of thinning on structural development in 40- to 100-year-old Douglas-fir stands in western Oregon. Forest Ecology and Management 108:99-113.
- Beggs, L. R. 2004. Vegetation response following thinning in young Douglas-fir forests of western Oregon: can thinning accelerate development of late-successional structure and composition? M.S. Thesis. Oregon State University, Corvallis, Oregon.
- Bennett, J. N., L. L. Blevins, J. E. Barker, D. P. Blevins, and C. E. Prescott. 2003. Increases in tree growth and nutrient supply still apparent 10 to 13 years following fertilization and vegetation control of salal-dominated cedar-hemlock stands on Vancouver Island. Canadian Journal Forest Research 33:1516-1524.
- Binkley, D. 2004. A hypothesis about the interaction of tree dominance and stand production through stand development. Forest Ecology and Management 190:265-271.
- Binkley, D., U. Olsson, R. Rochelle, T. Stohlgren, and N. Nikolov. 2003. Structure, production and resource use in old-growth spruce/fir forests in the front range of the Rocky Mountains, USA. Forest Ecology and Management 172:271-279.
- Bolsinger, C. L., and K. L. Waddell. 1993. Area of old-growth forest in California, Oregon, and Washington. USDA Forest Service Resource Bulletin PNW-RB-197. Pacific Northwest Research Station, Portland, Oregon.
- Carey, A. B., and C. A. Harrington. 2001. Small mammals in young forests: implications for management for sustainability. Forest Ecology and Management 154:289-309.
- Carey, A. B., and M. L. Johnson. 1995. Small mammals in managed, naturally young, and old-growth forests. Ecological Applications 5:336-352.
- Clement, J. P., and D. C. Shaw. 1999. Crown structure and the distribution of epiphyte functional group biomass in old-growth *Pseudotsuga menziesii* trees. Ecoscience 6:243-254.

12 Davis, Puettmann, and Tucker

all field crews that participated in data collection that made this work possible. Helpful reviews on earlier versions of the manuscript were provided by Patricia Muir, James Agee, and Suzanne Schwab. Lisa Ganio also contributed to reviews in addition to providing assistance with data analysis and interpretation. We would also like to thank Gody Spycher, Maureen Duane, and Steven Garman for assistance with YSTDS background information and data management.

- Cook, J. G., T. W. Stutzman, C. W. Bowers, K. A. Brenner, and L. L. Irwin. 1995. Spherical densiometers produce biased estimates of forest canopy cover. Wildlife Society Bulletin 23:711-717.
- D'Amato, A. W. and K. J. Puettmann. 2004. The relative dominance hypothesis explains interaction dynamics in mixed species *Alnus rubra/Pseudotsuga menziesii* stands. Journal of Ecology. 92(3):450-463.
- Everett, R. L., and S. H. Sharrow. 1985. Soil water and temperature in harvested and nonharvested pinyon-juniper stands. USDA Forest Service Research Paper PNW-RP-342. Intermountain Forest and Range Experiment Station, Fort Collins, Colorado.
- Forsman, E. D., E. C. Meslow, and H. M. Wight. 1984. Distribution and biology of the spotted owl. Wildlife Monographs 87:1-64.
- Franklin, J. F., and C. T. Dyrness. 1973. The natural vegetation of Washington and Oregon. USDA Forest Service General Technical Report PNW-GTR-8. Pacific Northwest Research Station, Portland, Oregon.
- Franklin, J. F., and T. A. Spies. 1991a. Composition, function, and structure of old-growth Douglas-fir forests. USDA Forest Service General Technical Report PNW-GTR-285. Pacific Northwest Research Station, Portland, Oregon.
- Franklin, J. F., and T. A. Spies. 1991b. Ecological definitions of old-growth Douglas-fir forests. USDA Forest Service General Technical Report PNW-GTR-285. Pacific Northwest Research Station, Portland, Oregon.
- Franklin, J. F., T. A. Spies, R. Van Pelt, A. B. Carey, D. A. Thornburgh, D. R. Berg, D. B. Lindenmayer, M. E. Harmon, W. S. Keeton, D. C. Shaw, K. Bible, and J. Chen. 2002. Disturbances and structural development of natural forest ecosystems with silvicultural implications, using Douglas-fir forests as an example. Forest Ecology and Management 155:399-423.
- Franklin, J. F., and R. Van Pelt. 2004. Spatial aspects of structural complexity in old-growth forests. Journal of Forestry 102:22-28.
- Gray, A. N., T. A. Spies, and M. J. Easter. 2002. Microclimatic and soil moisture responses to gap formation in coastal Douglas-fir forests. Canadian Journal Forest Research 32:332-343.
- Hagar, J. C., W. C. McComb, and W. H. Emmingham. 1996. Bird communities in commercially thinned and unthinned Douglas-fir stands of western Oregon. Wildlife Society Bulletin 24:353-366.

- Hagar, J.C., S. Howlin, and L. Ganio. 2004. Short-term response of songbirds to experimental thinning of young Douglas-fir forests in the Oregon Cascades. Forest Ecology and Management 199:333-347.
- Halpern, C. B., and T. A. Spies. 1995. Plant species diversity in natural and managed forests of the Pacific Northwest. Ecological Applications 5:913-934.
- Hanus, M. L., D. D. Marshall, and D. W. Hann. 1999. Heightdiameter equations for six species in the coastal regions of the Pacific Northwest. Research Contribution 25, Forest Research Laboratory, Oregon State University, Corvallis, Oregon.
- Hayes, J. P., S. S. Chan, W. H. Emmingham, J. C. Tappeiner, L. D. Kellogg, and J. D. Bailey. 1997. Wildlife response to thinning young forests in the Pacific Northwest. Journal of Forestry 95:28-33.
- Hill, M. O. 1973. Diversity and evenness: a unifying notation and its consequences. Ecology 54:427-432.
- Huffman, D. W., and J. C. Tappeiner. 1997. Clonal expansion and seedling recruitment of Oregon grape (*Berberis* nervosa) in Douglas-fir (*Pseudotsuga menziesii*) forests: comparisons with salal (*Gaultheria shallon*). Canadian Journal of Forest Research 27: 1788-1793.
- MacArthur, R. H., and J. W. MacArthur. 1961. On bird species diversity. Ecology 42:594-598.
- Marshall, D. D., and R. O. Curtis. 2002. Levels-of-growing-stock cooperative study in Douglas-fir: Report No. 15-Hoskins: 1963-1998. USDA Forest Service Research Paper PNW-RP-537. Pacific Northwest Research Station, Portland, Oregon.
- Messier, C., and A. K. Mitchell. 1994. Effects of thinning in a 43-year-old Douglas-fir stand on above-and below-ground biomass allocation and leaf structure of understory *Gaultheria shallon*. Forest Ecology and Management 68:263-271.
- Miller, R. E., and R. L. Williamson. 1974. Dominant Douglas-fir respond to fertilizing and thinning in southwest Oregon. USDA Forest Service Research Note RN-PNW-216. Pacific Northwest Research Station, Portland, Oregon.
- Muir, P. S., R. L. Mattingly, J. C. Tappeiner, J. D. Bailey, W. E. Elliott, J. C. Hagar, J. C. Miller, E. B. Peterson, and E. E. Starkey. 2002. Managing for biodiversity in young Douglas-fir forests of western Oregon. USDI U.S. Geological Survey Biological Science Report. USGS/BRD/BSR-2002-0006.:76. Forest and Rangeland Ecosystem Science Center, Corvallis, Oregon.
- Oliver, C. D., and B. C. Larson. 1996. Forest Stand Dynamics. John Wiley & Sons, New York.
- Oliver, C. D., and M. D. Murray. 1983. Stand structure, thinning prescriptions, and density indexes in a Douglas-fir thinning study, Western Washington, U.S.A. Canadian Journal of Forest Research 13:126-136.
- Parker, W. C., K. A. Elliott, D. C. Dey, E. Boysen, and S. G. Newmaster. 2001. Managing succession in conifer plantations: converting young red pine (*Pinus resinosa* Ait.) plantations to native forest types by thinning and underplanting. The Forestry Chronicle 77:721-734.
- Poage, N. J., and J. C. Tappeiner. 2002. Long-term patterns of diameter and basal area growth of old-growth Doug-

las-fir trees in western Oregon. Canadian Journal of Forest Research 32:1232-1243.

- Rambo, T. R., and P. S. Muir. 1998. Forest floor bryophytes of *Pseudotsuga menziesiiI-Tsuga heterophylla* stands in Oregon: influence of substrate and overstory. Bryologist 101:116-130.
- Ritchie, M. W., and D. W. Hann. 1987. Equations for predicting height to crown base for fourteen tree species in southwest Oregon. Research Paper 50, Forest Research Lab, Oregon State University, Corvallis, Oregon.
- Rosso, A. L. 2000. Shrub epiphyte communities in relation to stand management in forests of western Oregon. Ph.D. Dissertation. Oregon State University, Corvallis, Oregon.
- SAS Institute Inc. 2001. SAS OnlineDoc Version 8.2 [computer program]. SAS Institute Inc., Carey, North Carolina.
- Smith, D. M., B. C. Larson, M. J. Kelty, and P. M. S. Ashton. 1997. The practice of silviculture: applied forest ecology, 9th edition. John Wiley & Sons, New York.
- Spies, T. A. 1991. Plant species diversity and occurrence in young, mature, and old-growth Douglas-fir stands in western Oregon and Washington. USDA Forest Service General Technical Report PNW-GTR-285. Pacific Northwest Research Station, Portland, Oregon.
- Spies, T. A., and J. F. Franklin. 1991. The structure of natural young, mature, and old-growth Douglas-fir forests in Oregon and Washington. *In*: L.F. Ruggiero, K.B. Aubry, A.B. Carey, and M.H. Huff (Editors). Wildlife and vegetation of unmanaged Douglas-fir forests. USDA Forest Service General Technical Report PNW-GTR-285. Pacific Northwest Research Station, Portland, Oregon. Pp. 91-110.
- Staebler, G. R. 1956. Effect of controlled release on growth of individual Douglas-fir trees. Journal of Forestry 54:567-568.
- Tappeiner, J. C., D. Huffman, D. Marshall, T. A. Spies, and J. D. Bailey. 1997. Density, ages, and growth rates in oldgrowth and young-growth forests in coastal Oregon. Canadian Journal of Forest Research 27:638-348.
- Tappeiner, J. C., and J. C. Zasada. 1993. Establishment of salmonberry, salal, vine maple, and bigleaf maple seedlings in the coastal forests of Oregon. Canadian Journal of Forest Research 23:1775-1780.
- Thomas, S. C., C. B. Halpern, D. A. Falk, D. A. Liguori, and K. A. Austin. 1999. Plant diversity in managed forests: understory responses to thinning and fertilization. Ecological Applications 9:864-879.
- Thysell, D. R., and A. B. Carey. 2000. Effects of forest management on understory and overstory vegetation: a retrospective study. USDA Forest Service General Techngical Report PNW-GTR-488. Pacific Northwest Research Station, Portland, Oregon.
- Thysell, D. R., and A. B. Carey. 2001. Manipulation of density of *Pseudotsuga menziesii* canopies: preliminary effects on understory vegetation. Canadian Journal of Forest Research 31:1513-1525.
- Tucker, G. F., and W. H. Emmingham. 1977. Morphological changes in leaves of residual western hemlock after clear and shelterwood cutting. Forest Science 23:195-203.

Alternative Thinning in Douglas-fir 13

Tucker, G. F., T. M. Hinckley, J. W. Leverenz, and S. M. Jiang. 1987. Adjustments of foliar morphology in the acclimation of understory Pacific silver fir following clearcutting. Forest Ecology and Management 21:249-268.

Received 8 August 2006 Accepted for publication 7 December 2006 Winter, L. E., L. B. Brubaker, J. F. Franklin, E. A. Miller, and D. Q. DeWitt. 2002. Initiation of an old-growth Douglas-fir stand in the Pacific Northwest: a reconstruction from tree-ring records. Canadian Journal of Forest Research 32:1039-1056.

4 Davis, Puettmann, and Tucker