

The Use of Sequence-Based SSR Mining for the Development of a Vast Collection of Microsatellites in *Aquilegia formosa*

Brandon Schlautman¹, Vera Pfeiffer², Juan Zalapa^{1,3}, Johanne Brunet^{3,4}

¹Department of Horticulture, University of Wisconsin, Madison, USA

²Department of Zoology, University of Wisconsin, Madison, USA

³USDA-ARS, VCRU, University of Wisconsin, Madison, USA

⁴Department of Entomology, University of Wisconsin, Madison, USA

Email: jbrunet@wisc.edu

Received 8 May 2014; revised 12 June 2014; accepted 7 July 2014

Copyright © 2014 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Numerous microsatellite markers were developed for *Aquilegia formosa* from sequences deposited within the Expressed Sequence Tag (EST), Genomic Survey Sequence (GSS), and Nucleotide databases in NCBI. Microsatellites (SSRs) were identified and primers were designed for 9 SSR containing sequences in the Nucleotide database, 3803 sequences in the EST database, and 2226 sequences in the GSS database. For validation purposes, 45 primer pairs were used to amplify DNA from 16 *A. formosa* individuals from the H. J. Andrews experimental forest in Oregon, a Long Term Ecological Research (LTER) site. Genetic polymorphisms were identified at 30 of the 45 microsatellite loci with an average of 13.2 alleles per locus, and the observed level of heterozygosity was greater than 0.8 for 21 of the 30 loci. The use of these polymorphic loci was sufficient to individually separate the 16 individuals using a principal coordinate analysis. This comprehensive collection of primers significantly increased the availability of microsatellite primers for *Aquilegia* spp. and provided ample material for future studies that required highly variable SSRs such as mapping and association studies and investigation of plant mating system and gene flow.

Keywords

Aquilegia formosa, Microsatellites, SSR Mining

1. Introduction

The western columbine, *Aquilegia formosa*, is a perennial herb from the family Ranunculaceae. It belongs to the

genus *Aquilegia*, which includes approximately 75 species with the similar number of species occurring in North America, Asia and Europe [1]. The phylogenetic relationships among species within the genus have been difficult to resolve, possibly due to a rapid speciation in the group [2]–[4]. Two independent radiation events have emerged, however, one leading to the North American and the other to the European clades [1] [4].

The ecology of many *Aquilegia* species has been well studied and there exists ample variation in pollinators, floral traits and mating systems among species, among populations and even within populations of the same species, with evidence of some causal relationships between pollinators, floral traits and mating systems [3] [5]–[9] [10]. Phenotypic plasticity exists for many of the floral traits with potential adaptive responses to global warming [11] [12]. Habitats can range from Alpine to Lowland, and can include desert springs, temperate forests and rocky outcrops [1] [13]. Moreover, habitat specialization may have played a greater role in the radiation of columbines in Europe while pollinator specialization has been more important in the diversification of North American species [1] [3].

The presence of unusual floral organs in *Aquilegia* species, which include sepals that have the appearance of petals (petaloid sepals), the staminodium, an additional whorl of a novel type of organ placed between carpels and stamens, and petal with a spur with nectaries found at the tip of the spurs, permits the evolutionary study of floral novelties [14] [15]. As a result of its variation in ecology, floral morphology, pollination, floral development and mating system, the genus *Aquilegia* is ascending to the status of model species for studies in ecology, evolution and plant development [16] [17]. To improve its usefulness as a model species, genomic tools are being developed for this genus [18], including the recent release of the *A. coerulea* Goldsmith whole genome sequence (<http://www.phytozome.net>). The plant *A. coerulea* Goldsmith is, however, not the wild *A. coerulea* species, but represents an unknown mixture of different *Aquilegia* species of both North American and European origins (S. A. Hodges, personal communication).

Microsatellites remain important genetic markers in many studies including genetic mapping, association mapping, investigations of plant mating systems and paternity analyses. Despite the recent development of nuclear tools for *Aquilegia*, the suitability of the primers developed for currently existing microsatellite loci [19]–[21] remains limited because some of the loci are redundant [22] and others display low levels of polymorphism [21]. In this study, we identified microsatellites (SSRs) and designed primers from the Nucleotide database, the Expressed Sequence Tag (EST) database, and the Genomic Survey Sequence (GSS) database through NCBI, and validated the screening by testing 45 of the primers using a wild *A. formosa* population. Our goal was to develop a larger number of microsatellite loci with a wide range of motif length and repeat number from this data mining process to facilitate well-needed studies of population genetic structure of *Aquilegia* species [22] [23], and to promote further investigations of the role of pollinators on plant reproductive success [24], mating system [8], floral selection and gene movement within and among populations of *Aquilegia* species.

2. Materials and Methods

2.1. Data Mining of SSRs

The SSR mining was performed on 128,082 *Aquilegia formosa* nucleotide sequences publically available online through NCBI. These sequences included 106 sequences from the nucleotide database and 12,310 sequences from the genomic survey sequences (GSS) database for *A. formosa*, and an additional 115,666 sequences in the EST database from *A. formosa* × *A. pubescens* pooled RNA. The presence of short sequence repeats (SSRs) within these sequences was detected using SSR Locator [25]. The program parameters were set to identify and localize microsatellite motifs ranging from 1 - 10 bp and only the sequences containing motifs with repeat lengths of mono ≥ 12 , di ≥ 6 , tri ≥ 4 , and tetra through nona ≥ 3 were considered to be SSRs.

We chose the program WebSat [26] to design the oligonucleotide primers from the SSR flanking sequences because it offers the unique ability to visualize the SSR containing sequences. This feature ensured that the primer sequence did not overlap with the SSR of interest and that no primer pair contained more than one SSR. The major parameters selected for primer design were a primer length of 19 to 25 bp (optimum 22 bp), a PCR product size varying between 120 and 325 bp, a GC content between 40% and 80%, and an optimum melting temperature of 55°C.

2.2. Plant Material and DNA Isolation

Leaf tissue was collected from 16 *A. formosa* individuals from a population located in the H. J. Andrews expe-

rimental forest in Oregon, a Long Term Ecological Research (LTER) site. Permission to collect was obtained from Mark Schulze, the Director of the H. J. Andrews experimental forest. The plant species, *A. formosa*, is not an endangered or protected species and the coordinates for the population were 44.12656 latitude and -122.07827 longitude. Total genomic DNA from 0.1 g leaf tissue was extracted using a Macherey-Nagel (MN) Plant II kit (Düren, Germany) following the manufacturer's instructions.

2.3. PCR, SSR Genotyping, and Diversity Analysis

The SSR forward primers were appended with the M13 sequence (5'-CACGACGTTGTAAAACGAC-3') to allow indirect fluorescent labeling of reactions [27]. The reverse primers were appended with the PIG sequence (5'-GTTTCTT-3') to promote non-templated (A) addition and facilitate genotyping [28]. The PCR reactions were performed in 8 μ L total volume using 3.5 μ L 1 \times JumpStart REDTaq ReadyMix (Sigma, St. Louis, MO, USA), 1.0 μ L of 15 ng/ μ L genomic DNA, 2.0 μ L of H₂O, 0.5 μ L of 5 μ M M13-FAM or HEX primer, 0.5 μ L of 5 μ M forward primer, 0.5 μ L of 50 μ M reverse primer. The thermocycling conditions included an initial melting step (94°C for 3 min), followed by 33 cycles of 94°C for 15 s, 55°C for 90 s, and 72°C for 2 min, a final elongation step (72°C for 30 min), and an indefinite soak at 4°C. The PCR fragment separation was performed at the University of Wisconsin Biotechnology Center DNA Sequence Facility using an ABI 3730 fluorescent sequencer (POP-6 and a 50 cm array; Applied Biosystems, Foster City, CA, USA). Individuals were genotyped using carboxyfluorescein (FAM) and hexachlorofluorescein (HEX) M13 tags and a carboxy-X-rhodamine (ROX) ladder (GeneFlo-625 ROX; CHIMERx, Milwaukee, WI, USA). The resulting sequences were visualized using GeneMarker software v1.91 (SoftGenetics, State College, PA, USA).

Forty-five SSR primer pairs were used to conduct a survey of genetic polymorphism using 16 *A. formosa* individuals. These primer pairs were selected based on the length of their motif and the number of repeats. Nine of these primer pairs came from the nucleotide sequence database, 18 from the GSS, and 18 from the EST sequences. We attempted to redesign primers for specific use with M13 sequences and PIG tails for the 32 SSR loci which were previously available for *A. formosa* [19] [20]. However, because of the high stringency of our parameters (length, GC content, melting temperatures, and annealing temperatures), the observed redundancy of loci, and the SSR position in the sequences obtained from NCBI, we were only able to redesign acceptable primer pairs for eight loci. Therefore, eight of the 9 sequences containing SSRs from the nucleotide database in this study were previously identified [19] although the current study used different parameters and protocols for primer pairs design and genotyping.

The observed number of alleles (N_a), the levels of observed (H_o) and expected heterozygosity (H_e), the Shannon Information Index (I), and the major allele frequency (M_{AF}) or frequency of the most common allele were calculated for each polymorphic locus in GenAlEx 6.4 [29]. In addition, Hardy-Weinberg Equilibrium (HWE), null allele frequency, and non-exclusion probability of identity (NE-I) were calculated for each polymorphic locus using Cervus 3.0 [30]. One-Way ANOVAs were performed to identify differences in diversity statistics among the three databases (Nucleotide, EST and GSS). Finally we performed a principal coordinate analysis (PCoA) with all polymorphic loci in GenAlEx 6.4 [29] based on the genetic distances between pairs in order to demonstrate the markers' ability to discriminate between unique individuals.

3. Results and Discussion

A screen of the 128,082 *A. formosa* sequences deposited in NCBI identified a total of 65 SSRs in the nucleotide database, 4372 SSRs in the GSS database, and 11,519 SSRs in the EST database. The most frequent motifs identified among the 15,956 ungapped SSR loci were trinucleotides (48.9%), followed by mononucleotides (19.1%), tetranucleotides (13.2%) and finally dinucleotides (9.5%) (**Figure 1**; **Table S1**). The most frequent mononucleotide motif was A (17.2% of the total), dinucleotide was AG (4.6%), trinucleotide, GAA (7.6%) and tetranucleotide, AGAA (2.0%). The 15,956 sequences containing SSR were inspected for suitable flanking sites for primer design, and primer-pairs were designed for 9 of the SSRs from the nucleotide database, 3,803 of the EST SSRs, and 2226 of the GSS SSRs using WebSat [26] (**Table S2** and **Table S3**).

Thirty of the 45 SSR primer pairs tested amplified in the expected size range (**Table 1**) and displayed polymorphism across individuals (Nucleotide: 6, GSS: 10, and EST: 14 primer pairs; (**Table 2**)). Of the remaining 15 primer pairs tested, nine did not amplify at all and 6 amplified allelic patterns inconsistent with single locus segregation in a diploid species (**Table S4**). Of the 8 SSR loci from the Nucleotide database which were rede




Figure 1. Frequency of repeat numbers within mononucleotide through nonanucleotide microsatellite (SSR) motif levels. The nonanucleotide for GSS and EST only have repeats of 3 while the octanucleotides for GSS have repeats of 3 and 4. These repeat numbers were examined within 11,519 SSRs identified in the NCBI EST database from *A. formosa* × *A. pubescens* pooled RNA and within 4,372 SSRs identified in the NCBI GSS database for *A. formosa*.

signed for this study [19], only 6 amplified in a diploid inheritance pattern (Table 2, Table S4). Therefore, of the SSR loci previously available for *Aquilegia*, 20 have low allelic diversity [21] and only 6 of the 32 SSR loci with higher allelic diversity [19] [20] met the more stringent criteria used for primer development in this study.

The 30 polymorphic loci yielded 408 alleles with an average of 13.6 alleles and a range of 5 to 25 alleles (N_a) per locus (Table 2). These values are consistent with previous primer discoveries in *Aquilegia* species from North America where, for 28 individuals and 16 primers, an average of 14.4 alleles per locus has been reported for *A. formosa*, and 13.9 alleles per locus for *A. pubescens* [19]. In addition, Gallagher *et al.* (2004) [20] reported 15.5 alleles per locus in *A. chrysantha* from 16 primers in 165 individuals. Lower number of alleles per locus has been detected in two Asian columbine species, with 2 - 4 alleles per locus in *A. flabellata* and 2 to 5 in *A. oxysepala* [21]. For most of the loci examined, the level of heterozygosity was high, a pattern that has been previously observed for another columbine species, *A. coerulea* [11].

The results of the ANOVAs revealed no statistically significant differences among the three databases, GSS, EST and Nucleotide, in the total number of alleles per locus (GSS = 14.5 ± 4.3; EST = 12.1 ± 5.5; Nucleotide = 15.5 ± 3.6), ($F_{2,29} = 1.3$, $p = 0.29$), major allele frequencies (GSS = 0.20 ± 0.08; EST = 0.28 ± 0.16; Nucleotide = 0.23 ± 0.11) ($F_{2,29} = 1.2$, $p = 0.32$), observed (H_o) (GSS = 0.83 ± 0.15; EST = 0.78 ± 0.18; Nucleotide = 0.79 ± 0.12) ($F_{2,29} = 0.29$, $p = 0.75$) and expected heterozygosities (H_e); (GSS = 0.89 ± 0.04; EST = 0.83 ± 0.12; Nucleotide = 0.88 ± 0.07) ($F_{2,29} = 1.53$, $p = 0.23$) and Shannon indices (GSS = 2.44 ± 0.31; EST = 2.15 ± 0.56; Nucleotide = 2.47 ± 0.38) ($F_{2,29} = 1.61$, $p = 0.22$) (Table 2). This lack of difference is present despite the fact that the microsatellite loci identified in the EST database are transcript-based SSRs and are therefore considered to be more evolutionarily conserved and potentially less polymorphic than SSRs developed from the genome [28]. Similar results have been observed in germplasm screens of potato and water

Table 1. Primer sequences and their relevant characteristics. These primers were designed for 30 polymorphic SSR containing loci from *A. formosa* identified in sequences retrieved from the Nucleotide, Expressed Sequence Tag (EST), and Genomic Survey Sequences (GSS) NCBI databases.

NCBI ID	Motif	Size (bp) ^d	Forward Sequence (5'-3')	Reverse Sequence (5'-3')
AY566429.1 ^a	(CT)12	205 - 263	ACCACAAATCTTATTCACGG ^e	ACAAATCTGGTTGAGAGAGATG
AY566430.1 ^a	(CT)24	305 - 350	CTGCTTGTTCTAGGGTTGTAT ^f	CGGAGATTTGAGAGAAGTAGAT
AY566431.1 ^a	(AG)37	237 - 312	CGAGATACAAGAAATCAGAGAG ^e	CAACTTTCACTTCGTCAGC
AY566433.1 ^a	(TC)23	271 - 314	CCAGTGTAAGAGGAGAAGAAAG ^e	GCTTGAAATTACCCAATACG
AY566438.1 ^a	(TC)16	252 - 279	ATGCACTTACACCACTCACTAA ^f	AGGATTCATAGTTGGGTTGA
AY566442.1 ^a	(TC)15	195 - 280	GAGACAGATACCAAGGTTTCATC ^f	TATAGAAGTGTAAAACGGTGGG
ER968006.2 ^b	(AG)16	292 - 402	GCTGCTAAAGTAGAGAGTCAGAA ^e	ATTAAGTATGAGGGTGAGAAG
ER969526.2 ^b	(AT)17	304 - 389	GATTTAAGGGTTGCTTTGGT ^e	CATCTCTGCTAAGAGTCCAAGA
ER937207.2 ^b	(AG)17	142 - 168	ACTATTCAAGAACCAGCGACT ^e	CCACGTCAGATACACATACACT
ER969057.2 ^b	(TA)18	277 - 346	CTCTGTGACTTTGTGAGCATA ^e	GGCCTACGGACTTCATTATT
ER973157.2 ^b	(GA)18	173 - 200	AGGATCATAACACTTACTTGCC ^e	ATTCTCAACAGCTTCTCATCTC
ER940655.2 ^b	(TC)19	275 - 313	GATGAGGTAGGTTTTAATGGG ^e	GCGTCTGGTTTTAGTTCTTCT
ER969333.2 ^b	(CT)21	219 - 271	CTTCCTCTTTGTCTTTTGTC ^e	CTATGAAGTGAGGCTGAAGAAC
ER973191.2 ^b	(ATT)14	286 - 359	CACGCAACAAAACACTGAAF ^e	GCTGCCTAATAATACCTACAC
ER939871.2 ^b	(AAG)15	138 - 179	GAAGTAGAGATTGGTGGTGAAG ^e	TGATCTTGAGAGAAAGTGAGGT
ER938628.2 ^b	(GAA)21	155 - 193	GAGGAGATGTGTTTTGATAGGT ^e	CGGTTGTGTGTTGAAAGAA
JZ009091.1 ^c	(CAC)6	311 - 321	TATCCTTACCACCACAAACC ^f	ATCTACTCCACCATAGTCTCCA
DR951797.1 ^c	(GA)16	310 - 380	AGCACACAAATACACCACTCTA ^f	GAGTTCAAGAAGAGATTGATCC
JZ021896.1 ^c	(AG)19	401 - 490	GTCTCCTAATAACAATAGCCCT ^f	GGTGATGGTGGATACTCATCT
JZ007530.1 ^c	(AG)19	209 - 219	ATACTCATCAGAGAACTGCACA ^f	ATGTGGCGAGAAAGATACAT
DT748959.1 ^c	(TC)19	255 - 286	CACACCCTCTTCACCTCTT ^f	GTGATATTTGTGGACTTGGTG
DR952661.1 ^c	(GA)19	275 - 321	CCTAACCAGTAACCTCTGTTCT ^f	GTTTTCGAGATGAAAGGAAG
DR924522.1 ^c	(GA)19	134 - 192	CACTCCAGTCTTGCTAATCATC ^f	GACCTTTTCCTTTCTCTGCTAT
DT755604.1 ^c	(GA)20	204 - 241	GAGTGATTCTAGCTTGGATAGTC ^f	CACAAGATTTAAGTGGAGACCT
DR912270.1 ^c	(AAG)11	211 - 228	GAGTGTTGTTAAGGCAAGAGTT ^f	AATAATGGTCTCCCTCGATT
DT763172.1 ^c	(AGA)12	311 - 362	CCTACCTTTCTTCAATTCACAC ^f	ATACTGGTTGGGACGAAGTAG
DT741616.1 ^c	(AAG)12	414 - 432	CTTGTACCATCTCACTCTCCAT ^f	CCTTCAGTTTGTAAACCATAACC
DR922072.1 ^c	(GAA)15	208 - 228	ATGCTAGTTTTAGAGGAGATGG ^f	ATAGGACTTTGAGGAGGAGAGT
DR945073.1 ^c	(CCA)16	274 - 207	CTATGCAAGTGGAACAGAAAC ^f	CCTAAGAAGAACATCCCAACT
DR916303.1 ^c	(GAA)18	208 - 228	ATGCTAGTTTTAGAGGAGATGG ^f	ATAGGACTTTGAGGAGGAGAGT

^aSequences from the NCBI Nucleotide Database; ^bSequences from the NCBI GSS Database; ^cSequences from the NCBI EST Database; ^dM13 tail and PIG tail included in the observed allele size range; ^ePCR performed with 5' FAM fluorescently labeled primers; ^fPCR performed with 5' HEX fluorescently labeled primers.

melon using EST-SSR loci [31] [32]. Perhaps the high levels of polymorphism observed in transcript-based SSRs for *A. formosa* adults in the H. J. Andrews experimental forest could be due to the fact that homozygotes

Table 2. Diversity statistics for 30 polymorphic SSR loci tested in 16 *A. formosa* individuals.

Database	NCBI Sequence ID	A	M _{AF}	H _O	H _e	I	HWE	NE-I	N (Full)
Nucleotide	AY566429.1	15	0.25	0.81	0.88	2.42	ND	0.02	0.18
	AY566430.1	17	0.13	0.88	0.92	2.70	ND	0.01	0.08
	AY566431.1	21	0.16	0.88	0.93	2.90	ND	0.01	0.01
	AY566433.1	16	0.19	0.75	0.91	2.62	ND	0.01	-0.01
	AY566438.1	14	0.20	0.87	0.88	2.39	ND	0.02	0.06
	AY566442.1	10	0.44	0.56	0.75	1.80	ND	0.08	-0.04
GSS	ER968006.2	25	0.06	0.94	0.96	3.16	ND	0.00	-0.05
	ER969526.2	16	0.34	0.88	0.84	2.32	ND	0.04	-0.06
	ER937207.2	12	0.16	0.88	0.89	2.34	ND	0.02	0.13
	ER969057.2	9	0.23	0.47	0.85	2.01	ND	0.04	0.01
	ER973157.2	13	0.13	0.87	0.91	2.50	ND	0.01	0.14
	ER940655.2	13	0.28	0.69	0.87	2.30	ND	0.03	-0.01
	ER969333.2	17	0.17	0.93	0.92	2.68	ND	0.01	0.19
	ER973191.2	15	0.22	0.81	0.89	2.46	ND	0.02	-0.01
	ER939871.2	12	0.22	0.88	0.86	2.17	ND	0.04	0.16
	ER938628.2	13	0.16	0.94	0.90	2.41	ND	0.02	0.02
EST	JZ009091.1	5	0.66	0.38	0.53	1.04	ND	0.26	0.01
	DR51797.1	22	0.09	0.81	0.95	3.02	ND	0.01	0.10
	J2021896.1	18	0.28	0.88	0.88	2.57	ND	0.02	0.04
	JZ007530.1	5	0.56	0.63	0.62	1.23	ND	0.19	0.04
	DT748959.1	13	0.23	0.80	0.89	2.38	ND	0.02	0.01
	DR952661.1	20	0.09	1.00	0.93	2.84	ND	0.01	-0.02
	DR924522.1	16	0.16	1.00	0.92	2.62	ND	0.01	0.01
	DT755604.1	15	0.31	0.94	0.84	2.26	ND	0.04	0.04
	DR912270.1	9	0.34	0.63	0.80	1.86	ND	0.06	0.06
	DT763172.1	14	0.16	0.88	0.90	2.47	ND	0.02	0.02
	DT741616.1	9	0.34	0.63	0.80	1.86	ND	0.06	0.06
	DR922072.1	8	0.22	0.88	0.85	1.99	ND	0.04	0.04
	DR945073.1	8	0.25	0.56	0.84	1.92	ND	0.05	0.05
	DR916303.1	8	0.22	0.88	0.85	1.99	ND	0.04	0.04

Note: A = Number of alleles sampled; M_{AF} = Major allele frequency; H_O = Observed heterozygosity; H_e = Expected heterozygosity; I = Shannon's diversity index; HWE = Hardy-Weinberg Equilibrium; ND = No significant deviation; NE-I = Non-exclusion probability of identity; N (Full) = Frequency of null alleles.

are eliminated in the adult population as a result of the strong inbreeding depression that exists in the species [33].

The 30 polymorphic SSRs were more than sufficient to separate all 16 *A. formosa* individuals from the H. J. Andrews experimental forest in a principal coordinate analysis (PCoA) (Figure 2). The extremely low non-ex-




Figure 2. Principal Coordinate Analysis (PCoA) showing the genetic relationships of 16 *A. formosa* individuals. Leaf tissue for these individuals was collected from the H. J. Andrews experimental forest in Oregon, USA. The PCoA used alleles from 30 polymorphic microsatellite loci identified in sequences retrieved from the NCBI Nucleotide, Expressed Sequence Tag (EST), and Genomic Survey Sequence (GSS) databases.

clusion probability of identity for the 30 markers ($9.33\text{E}-49$) indicates that these markers are suitable for genetic fingerprinting and other studies. For example, genotyping using only two primers, DR51797 and ER968006.2, discriminated all 16 individuals because the combined non-exclusion probability of identity for the two markers was $2\text{E}-5$. These results indicate the potential of these SSRs for examining population-level processes such as paternity analyses and studies where individuals must be identified [24]. Researchers can achieve the same genetic information content using one microsatellite with k alleles as $(k-1)$ biallelic markers such as SNPs [34]. Furthermore, SSRs are relatively evenly distributed across genomes which make them useful for conducting comparative genomic analyses, linkage and QTL mapping, and genome wide association studies [35]. Finally, the transcript-based SSRs for which primers were designed in this species are likely to be associated with functional genes and could be an important resource for studies searching for and identifying genes in *Aquilegia*. The 6038 primer-pairs designed for SSR loci in *A. formosa* and the 30 polymorphic SSR which were tested and validated (Tables S2 & S3) complement the molecular tools currently available for *Aquilegia* research [18].

4. Conclusion

An extensive number of microsatellite primers located in transcribed and genomic regions were developed for *A. formosa*. This collection is the most comprehensive set of microsatellite primer-pairs developed in *Aquilegia* to date, and illustrates how sequence-based SSR mining can produce many microsatellites of a diverse motif type and repeat length that adds to the existing microsatellite libraries often based on enrichment methods to provide a larger marker resource for scientists. The collection of microsatellites presented here contributes to the various molecular tools being developed in *Aquilegia*. Given the extensive knowledge of the ecology, pollination biology, floral biology and mating system of some *Aquilegia* species, these molecular tools can facilitate the mapping of functional genes and promote studies of the population genetic structure, selection and gene flow in diverse species.

Acknowledgements

Tissue samples of *A. formosa* were collected at H. J. Andrews Experimental Forest, which was funded by the National Science Foundation and the U.S. Forest Service, Pacific Northwest Research Station. Funding for this

project was provided by the USDA—Agricultural Research Service to J. B. and to J. E. Z.

References

- [1] Bastida, J.M., Alcantara, J.M., Rey, P.J., Vargas, P. and Herrera, C.M. (2010) Extended Phylogeny of *Aquilegia*: The Biogeographical and Ecological Patterns of Two Simultaneous but Contrasting Radiations. *Plant Systematics and Evolution*, **284**, 171–185. <http://dx.doi.org/10.1007/s00606-009-0243-z>
- [2] Hodges, S.A. and Arnold, M. (1994) Columbines: A Geographically Widespread Species Flock. *Proceedings of the National Academy of Sciences*, **91**, 5129–5132. <http://dx.doi.org/10.1073/pnas.91.11.5129>
- [3] Whittall, J.B. and Hodges, S.A. (2007) Pollinator Shifts Drive Increasingly Long Nectar Spurs in Columbine Flowers. *Nature*, **447**, 706–710. <http://dx.doi.org/10.1038/nature05857>
- [4] Fior, S., Li, M., Oxelman, B., Viola, R., Hodges, S.A., Ometto, L. and Varotto, C. (2013) Spatiotemporal Reconstruction of the *Aquilegia* Rapid Radiation through Next-Generation Sequencing of Rapidly Evolving cpDNA Regions. *New Phytologist*, **198**, 579–592. <http://dx.doi.org/10.1111/nph.12163>
- [5] Brunet, J. and Eckert, C.G. (1998) Effect of Floral Morphology and Display on Outcrossing in Blue Columbine, *Aquilegia caerulea* (Ranunculaceae). *Functional Ecology*, **12**, 596–606. <http://dx.doi.org/10.1046/j.1365-2435.1998.00231.x>
- [6] Fulton, M. and Hodges, S.A. (1999) Isolation between *Aquilegia formosa* and *Aquilegia pubescens*. *Proceeding of the Royal Society of London B*, **266**, 2247–2252.
- [7] Herlihy, C.R. and Eckert C.G. (2005) Evolution of Self-Fertilization at Geographical Range Margins? A Comparison of Demographic, Floral, and Mating System Variables in Central vs. Peripheral Populations of *Aquilegia canadensis* (Ranunculaceae). *American Journal of Botany*, **92**, 744–751. <http://dx.doi.org/10.3732/ajb.92.4.744>
- [8] Brunet, J. and Sweet, H.R. (2006) Impact of Insect Pollinator Group and Floral Display Size on Outcrossing Rate. *Evolution*, **60**, 183–195. <http://dx.doi.org/10.1111/j.0014-3820.2006.tb01102.x>
- [9] Herlihy, C.R. and Eckert, C.G. (2007) Evolutionary Analysis of a Key Floral Trait and Its Effect on the Mating System in *Aquilegia canadensis* (Ranunculaceae). *Evolution*, **61**, 1661–1674. <http://dx.doi.org/10.1111/j.1558-5646.2007.00137.x>
- [10] Brunet, J. (2009) Pollinators of the Rocky Mountain Columbine: Temporal Variation, Functional Groups and Associations with Floral Traits. *Annals of Botany*, **103**, 1567–1578. <http://dx.doi.org/10.1093/aob/mcp096>
- [11] Brunet, J. and Larson-Rabin, Z. (2012) The Response of Flowering Time to Global Warming in a High-Altitude Plant: The Impact of Genetics and the Environment. *Botany*, **90**, 319–326. <http://dx.doi.org/10.1139/b2012-001>
- [12] Van Etten, M.L. and Brunet, J. (2013) The Impact of Global Warming on Floral Traits That Affect the Selfing Rate in a High-Altitude Plant. *International Journal of Plant Sciences*, **174**, 1099–1108. <http://dx.doi.org/10.1086/671805>
- [13] Chase, V.C. and Raven, P.H. (1975) Evolutionary and Ecological Relationships between *Aquilegia formosa* and *A. pubescens* (Ranunculaceae), Two Perennial Plants. *Evolution*, **29**, 474–486. <http://dx.doi.org/10.2307/2407260>
- [14] Kramer, E.M., Holappa, L., BillieGould, M., Jaramillo, A., Setnikov, D. and Santiago, P.M. (2007) Elaboration of B Gene Function to Include the Identity of Novel Floral Organs in the Lower Eudicot *Aquilegia*. *The Plant Cell*, **19**, 750–766. <http://dx.doi.org/10.1105/tpc.107.050385>
- [15] Voelckel, C., Borevitz, J.O., Kramer, E.M. and Hodges, S.A. (2010) Within and between Whorls: Comparative Transcriptional Profiling of *Aquilegia* and *Arabidopsis*. *PLoS ONE*, **5**, e9735. <http://dx.doi.org/10.1371/journal.pone.0009735>
- [16] Kramer, E.M. (2009) *Aquilegia*: A New Model for Plant Development, Ecology, and Evolution. *Annual Review of Plant Biology*, **60**, 261–277.
- [17] Kramer, E.M. and Hodges, S.A. (2010) *Aquilegia* as a Model System for the Evolution and Ecology of Petals. *Philosophical Transactions of the Royal Society B*, **365**, 477–490. <http://dx.doi.org/10.1098/rstb.2009.0230>
- [18] Fang, G.C., Blackmon, B.P., Henry, D.C., Staton, M.E., Saski, C.A., Hodges, S.A., Tomkins, J.P. and Luo, H. (2010) Genomic Tools Development for *Aquilegia*: Construction of a BAC-Based Physical Map. *BMC Genomics*, **11**, 621. <http://dx.doi.org/10.1186/1471-2164-11-621>
- [19] Yang, J.Y., Counterman, B.A., Eckert, C.G. and Hodges, S.A. (2005) Cross-Species Amplification of Microsatellite loci in *Aquilegia* and *Semiaquilegia* (Ranunculaceae). *Molecular Ecology Notes*, **5**, 317–320. <http://dx.doi.org/10.1111/j.1471-8286.2005.00913.x>
- [20] Gallagher, K.G., Milligan, B.G. and White, P.S. (2004) Isolation and Characterization of Microsatellite DNA Loci in *Aquilegia* sp. *Molecular Ecology Notes*, **4**, 686–688. <http://dx.doi.org/10.1111/j.1471-8286.2004.00785.x>
- [21] Li, L.F., Pang, D., Liao, Q.L. and Xiao, H.X. (2011) Genomic and EST Microsatellite Markers for *Aquilegia flabellata* and Cross-Amplification in *A. oxysepala* (Ranunculaceae). *American Journal of Botany*, **98**, e213–e215.

- <http://dx.doi.org/10.3732/ajb.1100057>
- [22] Brunet, J., Larson-Rabin, Z. and Stewart, C.M. (2012) The Distribution of Genetic Diversity within and among Populations of the Rocky Mountain Columbine: The Impact of Gene Flow, Pollinators, and Mating System. *International Journal of Plant Sciences*, **174**, 1099-1108.
 - [23] Garrido, J.L., Fenu, G., Mattana, E. and Bacchetta, G. (2012) Spatial Genetic Structure of *Aquilegia* Taxa Endemic to the Island of Sardinia. *Annals of Botany*, **109**, 953-964. <http://dx.doi.org/10.1093/aob/mcs011>
 - [24] Brunet, J. and Holmquist, K.G.A. (2009) Influence of Distinct Pollinators on Female and Male Reproductive Success in the Rocky Mountain Columbine. *Molecular Ecology*, **18**, 3745-3758. <http://dx.doi.org/10.1111/j.1365-294X.2009.04304.x>
 - [25] da Maia, L.C., Palmieri, D.A., de Souza, V.Q., Kopp, M.M., de Carvalho, F.I. and de Oliveira, A.C. (2008) SSR Locator: Tool for Simple Sequence Repeat Discovery Integrated with Primer Design and PCR Simulation. *International Journal of Plant Genomics*, **2008**, 412696-412705.
 - [26] Martins, W.S., Lucas, D.C.S., Neves, K.F.S. and Bertioli, D.J. (2009) WebSat—A Web Software for MicroSatellite Marker Development. *Bioinformatics*, **3**, 282-283. <http://dx.doi.org/10.6026/97320630003282>
 - [27] Schuelke, M. (2000) An Economic Method for the Fluorescent Labeling of PCR Fragments. *Nature Biotechnology*, **18**, 233-234. <http://dx.doi.org/10.1038/72708>
 - [28] Zalapa, J., Cuevas, H., Zhu, H., Steffan, S., Senalik, D., Zeldin, E., McCown, B., Harbut, R. And Simon, P. (2012) Using Next-Generation Sequencing Approaches to Isolate Simple Sequence Repeat (SSR) Loci in the Plant Sciences. *American Journal of Botany*, **99**, 193-208. <http://dx.doi.org/10.3732/ajb.1100394>
 - [29] Peakall, R. and Smouse, P.E. (2006) GENALEX 6: Genetic Analysis in Excel. Population Genetic Software for Teaching and Research. *Molecular Ecology Notes*, **6**, 288-295. <http://dx.doi.org/10.1111/j.1471-8286.2005.01155.x>
 - [30] Kalinowski, S.T., Taper, M.L. and Marshall, T.C. (2007) Revising How the Computer Program CERVUS Accommodates Genotyping Error Increases Success in Paternity Assignment. *Molecular Ecology*, **16**, 1099-1006. <http://dx.doi.org/10.1111/j.1365-294X.2007.03089.x>
 - [31] Fu, Y.B., Peterson, G.W., Richards, K.W., Tarn, T.R. and Percy, J.E. (2009) Genetic Diversity of Canadian and Exotic Potato Germplasm Revealed by Simple Sequence Repeat Markers. *American Journal of Potato Research*, **86**, 38-48. <http://dx.doi.org/10.1007/s12230-008-9059-6>
 - [32] Mujaju, C., Sehic, J. and Nybom, H. (2013) Assessment of EST-SSR Markers for Evaluating Genetic Diversity in Watermelon Accessions from Zimbabwe. *American Journal of Plant Sciences*, **4**, 1448-1456.
 - [33] Yang, J.Y. and Hodges, S.A. (2010) Early Inbreeding Depression Selects for High Outcrossing Rates in *Aquilegia formosa* and *Aquilegia pubescens*. *International Journal of Plant Sciences*, **171**, 860-871. <http://dx.doi.org/10.1086/655772>
 - [34] Hamblin, M.T., Warburton, M.L. and Buckler, E.S. (2007) Empirical Comparison of Simple Sequence Repeats and Single Nucleotide Polymorphisms in Assessment of Maize Diversity and Relatedness. *PLoS ONE*, **2**, Article ID: e1367. <http://dx.doi.org/10.1371/journal.pone.0001367>
 - [35] Adam-Blondon, A.F., Roux, C., Claux, D., Butterlin, G., Merdinoglu, D. and This, P. (2004) Mapping 245 SSR Markers on the *Vitis vinifera* Genome: A Tool for Grape Genetics. *Theoretical and Applied Genetics*, **109**, 1017-1027. <http://dx.doi.org/10.1007/s00122-004-1704-y>

Appendix

Table S1. The total number of loci containing microsatellites for motif levels mono to nonanucleotide.

	Motif Level									
	Mono-	Di-	Tri-	Tetra-	Penta-	Hexa-	Hepta-	Octa-	Nona-	Total
Database										
Nucleotide	13	16	19	9	5	3	0	0	0	65
GSS	1714	505	883	797	278	142	46	2	5	4372
EST	1325	994	6902	1300	378	567	46	5	2	11,519
Combined	3052	1515	7804	2106	661	712	92	7	7	15,956

These microsatellites were identified in 106 *A. formosa* sequences from the NCBI Nucleotide database, 12,310 *A. formosa* sequences from the Genomic Survey Sequences (GSS) NCBI database, and an additional 115,666 sequences in the NCBI Expressed Sequence Tag (EST) database from *Aquilegia formosa* × *Aquilegia pubescens* pooled RNA.

Table S2. Untested primer sequences and their characteristics for 3785 microsatellite containing loci identified within *A. formosa* sequences deposited in the NCBI Expressed Sequence Tag (EST) database.

<https://www.dropbox.com/s/bg8xeu4ooj1qpqm/SupplementaryTable2.xlsx>

Table S3. Untested primer sequences and their characteristics for 2208 microsatellite containing loci identified within *A. formosa* sequences deposited in the NCBI Genomic Survey Sequence (GSS) database.

<https://www.dropbox.com/s/r935fi5q4rcer11/SupplementaryTable3.xlsx>

Table S4. Primer sequences and their relevant characteristics for 15 SSR loci from *A. formosa* which did not amplify, display polymorphism, or display the expected inheritance pattern.

Database	NCBI Sequence ID	Repeat Motif	Primer sequences (5'-3')	Fluorescence
Nucleotide	AY566435.1 ‡	(AC)11	F: GAAAGATGATATAGCGAACCC R: AGGAAGTAGTCCTCGAATAGC	FAM
	AY566441.1 ‡	(TC)25	F: TATAATGAGGGAGAAGACGAAG R: AGAAGTTATCCTGTAAAAGCCC	HEX
	GM711827.1 ‡	(TC)33	F: ATTCTCATCAGCTCTGGAAA R: CATCAGCACTCACTTCTATACG	FAM
GSS	ER936871.2 ‡	(CT)17	F: ATGACACATGGATTTCAGC R: AAACACAAGAATAAGACCCC	FAM
	ER969937.2*	(TA)19	F: GGCCTCATACATTCCAAAA R: GCATACCATCCGCTATAAATAC	FAM
	ER939203.2*	(AT)19	F: CTAGACGCTGGTAAATCTTCTC R: GAACTTAAAACAGCTCGTGACT	FAM
	ER939308.2*	(AG)21	F: GATGTGAGGAGCAAATATCATC R: TTTCTAACTGAAGAACTCTGGC	FAM
	ER972897.2*	(GA)21	F: GGGGAAGAGTAATTGCCAGA R: CCCTTATACTGGACCTTGAAT	FAM
	ER973211.2 ‡	(ATA)12	F: TCATAGAGAAAGCTGGATGAAC R: GTGTAAATTGTGAAGGAAGGAC	FAM
	ER970243.2 ‡	(CAT)15	F: ACTGCTCATTACCTCTTCTGT R: ACTTCAGACTATGGATGGGAT	FAM
	ER972134.2*	(TAT)24	F: TAAATATGCGTGGCTTTGAG R: ATAGACGACGAAAGGAAACG	FAM
EST	JZ017427.1*	(GA)16	F: CTCTTCTTACTTGTGTCATTGG R: GACTACTTTGAATTTGCCCTCT	HEX
	DR946658.1*	(CT)16	F: GTGCAACTCTCTCTCTCTCTCT R: TATACTGGAAAGACTAATGGCTG	HEX
	DR926317.1*	(GA)16	F: CTGGTACATCCAGAGCACAT R: GAAATGAACTCCTTCTCCATC	HEX
	DR928536.1*	(AG)18	F: ACTCAACTGCTACTCTCAAGT R: AGTAAAGAGAGACCCTGCTATG	HEX

Note: * = did not amplify; ‡ = amplify a non-diploid inheritance pattern.

Scientific Research Publishing (SCIRP) is one of the largest Open Access journal publishers. It is currently publishing more than 200 open access, online, peer-reviewed journals covering a wide range of academic disciplines. SCIRP serves the worldwide academic communities and contributes to the progress and application of science with its publication.

Other selected journals from SCIRP are listed as below. Submit your manuscript to us via either submit@scirp.org or [Online Submission Portal](#).

