U.S. FOREST SERVICE Research Paper PNW-3 169

CT HERE

NATURAL REGENERATION OF DOUGLAS-FIR AND ASSOCIATED SPECIES

USING MODIFIED CLEAR-CUTTING SYSTEMS

IN THE OREGON CASCADES

by

Jerry F. Franklin

September 1963

PACIFIC NORTHWEST FOREST AND RANGE EXPERIMENT STATION Philip A. Briegleb, Director Portland, Oregon

FOREST SERVICE U.S. DEPARTMENT OF AGRICULTURE

CONTENTS

and a construction will be the same

																								Page
INTROD	DUCTION				•	•	•		•				•	•		•	•	•		•			,	1
STUDY	AREAS	• •	•••				•	• •	•															1
CONE C	ROPS .						•		•			•												3
FIELD N	MEASUR	EME	NTS				• •						•							•				3
RESULI	FANT ST	OCKI	NG				•												•				r	4
REGENI	ERATION	N ANI	d Gr	OSS	C	HA	RA	CT	ΈF	RIS	TI	CS	6 C	F	C	LE	A	R	CL	JT	S			ó
REGENI	ERATION	N ANI	O CH	IAR	AC	ΤE	RIS	STI	CS	0	FI	MI	LA	7C	RI	ΕI	ΡL	0	ГS					6
S	Stand Sha	de.						•																6
I	Distance	from	See	d So	ur	ce		•				•				•		•						8
DISCUSS	SION OF	THE	MO	DIFI	ED	C C	UT	TII	NG.	S			·	·	•		·							9
Γ	North-So	uth St	rip	Clea	r (Cut	ts.							•						•				9
F	East-Wes	st Str	ip C	lear	Cı	uts			•		•	•		•				•		•				9
F	Patch Cle	ear C	uts		۰.																			11
S	Seed-Tre	e Cut	ting								•	•										•		11
CONCLU	JSIONS	•••							•			•								•			•	12
LITERA	TURE C	ITED																						13



INTRODUCTION

Clear cutting is the standard harvesting system in old-growth Douglasfir (Pseudotsuga menziesii) forests in the Pacific Northwest. Usually these clear cuts are in "staggered settings" of 15 to 80 acres with the surrounding stand left uncut to provide seed and serve as a firebreak. However, satisfactory natural regeneration of Douglas-fir does not always develop on such cuttings. The erratic and infrequent occurrence of abundant Douglas-fir seed crops and high surface soil temperatures appear to be major deterrents in many areas. Consequently, some modified clear cuts were tried in old-growth Douglas-fir in the western Oregon Cascade Range in 1954 and 1955 to determine their effect on natural regenerations. One trial of the seed-tree method was included. Cuttings were designed to leave residual trees in patterns that would reduce surface soil temperatures by providing shade and also improve distribution of natural seed fall. Resultant regeneration was examined in the summer of 1959.

STUDY AREAS

The study areas are located in the H. J. Andrews Experimental Forest on Lookout Creek, a tributary of McKenzie River in the western Oregon Cascades. Timber type before cutting was 400-year-old Douglas-fir with considerable western hemlock (Tsuga heterophylla) and some western redcedar (Thuja plicata). Merchantable timber volumes averaged 50,000 to 100,000 board feet per acre. Precipitation is heavy, in excess of 80 inches annually, and temperatures are mild, averaging 65° F. during the growing season (Berntsen and Rothacher, 1959).

Five general types of cuttings were tried: (1) Clear cutting in strips oriented north and south, (2) clear cutting in strips oriented east and west, (3) clear cutting in small patches ranging from 1/4 acre to 4 acres in size, (4) a seed-tree cutting, and (5) a staggered-setting clear cutting (table 1). Areas were cut and slash burned in 1954 and 1955. Considerable mineral soil seedbed was exposed in all treatments. Surface soils are clay loam in texture on all cuttings. Ground cover at time of the 1959 examination was relatively uniform with no outstanding differences among cutting units. $\frac{1}{2}$

 $\frac{1}{}$ For a discussion of vegetative composition and coverage on clear cuts in this area, the reader is referred to Yerkes (1960).

Table 1. -- Physical and historical data on modified cutting units,

	:	:	Dimension	is			Year		
Shape and number of unit	: Size :	: Width	: : Length :	: Radius :	Average slope	Average elevation	: Logged	Burned	
	Acres		<u>Feet</u>		Percent	Feet			
North-south oriented strip clear cuts:									
S1 S2 S3 S4	0.7 1.2 3.6 8.4	50 100 200 350	600 550 800 1,050		60 60 50 40	2,025 2,100 2,125 2,175	1954 1954 1954 1954	1954 1954 1955 1955	
East-west oriented strip clear cuts:									
\$5 \$6 \$7	2.3 3.1 4.0	100 200 300	1,000 650 575		10 10 20	2,550 2,600 2,650	1954 1954 1954	1954 1954 1955	
Rectangular patch clear cuts:									
G1 G2 G3	4.0 2.0 1.0	43 5 250 200	400 350 220		20 10 0	3,100 3,125 3,075	1955 1955 1955	1955 1955 1955	
Circular patch clear cuts:									
G4 G5 G6	.75 .50 .25			102 83 49	0 10 20	3,025 3,025 3,050	1955 1955 1955	1955 1955 1955	
Staggered-setting clear cut:									
5в	48.4				20	3,200	1955	1955	
Seed-tree cutting:									
50	20.4				10	3,300	1955	1955	

H. J. Andrews Experimental Forest

¢

Exposure and topography within individual clear cuts are fairly uniform except units S4, 5B, and 5C (table 1), where exposure varies due to presence of minor stream drainages. The north-south strip clear cuts are on a steep, southfacing slope with a relatively shallow soil at an elevation of 1,850 to 2,375 feet. The other cuttings are on a gentle, south-facing slope with deeper soil at elevations ranging from 2,550 to 3,500 feet. Thus, the north-south strip clear cuts occupy a hotter and drier habitat than the other cuttings.

CONE CROPS

Continuous records of Douglas-fir and western hemlock cone crops have been kept on the experimental forest. During the study period, average ratings²/ were:

Years	Douglas-fir	Western hemlock
1954	Medium	Abundant
1955	Light	Light
1956	Abundant	Medium
1957	Failure	Medium
1958	Failure	Light

FIELD MEASUREMENTS

Species, amount, and distribution of natural regeneration were used as the main criteria for comparing the different types of cuttings. Tree seedlings were counted on milacre plots located along transects. Transects (usually 10 per cutting) were surveyed across cuttings with their direction parallel to the narrow side of the cutting. Number of milacre plots varied from 90 to 225 per cutting, depending in part on size and shape of unit.

A Douglas-fir seedling in its second growing season was considered established; two l-year-old seedlings were considered its equivalent. A western hemlock or western redcedar seedling in its third season was considered established; three seedlings either l or 2 years old were considered its equivalent. For strip and patch clear cuts, hours during which the center of the plot received

 $[\]frac{2}{}$ Data based on averages of general observations of number of cones at seven locations. Ratings provide a relative measure only.

shade from adjacent timber between 6 a.m. and 6 p.m. were estimated, using a technique described by Silen. $\frac{3}{2}$

RESULTANT STOCKING

If stocking of any species on 30 percent of the milacre plots is considered adequate, all cuttings except the staggered-setting were adequately stocked with tree seedlings (table 2). $\frac{4}{}$ Considering Douglas-fir alone, all cuttings except the staggered-setting and 2-acre patch clear cut were stocked. As groups, the east-west strip clear cuts had the best stocking, north-south strip clear cuts and staggered-setting were poorest, and patch clear cuts and the seed-tree cutting were intermediate.

The staggered-setting clear cut had only 407 Douglas-fir seedlings per acre and 28 percent of the plots stocked with any species.

Poor stocking is typical of many large clear cuts in this general area. For example, twelve 3- to 5-year-old clear cuts on the experimental forest surveyed in 1956 had an average of only 301 (range 81 to 525) Douglas-fir seedlings per acre. Total plots stocked with any species averaged 28 (range 11 to 48) percent and plots stocked with Douglas-fir averaged only 24 (range 9 to 36) percent, even though both 1- and 2-year-old seedlings were included in the tally.

 $\frac{3}{}$ Silen, Roy Ragnar. Lethal surface temperatures and their interpretation for Douglas-fir. Unpublished Ph.D. thesis on file Oreg. State Univ. 170 pp., illus. 1960.

This technique involves use of a protractor placed with the index mark at center of plot and oriented with the 90° mark toward south. Protractor is tilted at a 45° angle toward north to aline its surface with celestial equator. Path of the sun for six spring and summer months lies somewhere in a band 23° wide north of celestial equator marked by protractor edge. Stand edge producing shade must lie within the skyward projection of this imaginary band. Since the sun moves along the band at 15° per hour, hours of shade on each plot can be estimated with the protractor in terms of degrees and then converted to hours. A pencil, with the point held at the center of the plot on the protractor index mark, was swung on the sun's apparent path, and the hours the plot was receiving stand shade were estimated. The principles for such approximations are found in any general surveying textbook.

 $\frac{4}{}$ Thirty percent is the usual 1-milacre stocking criterion used in the Northwest. Percent of milacre plots stocked is not completely satisfactory as a measure of overall adequacy of regeneration on a clear cut, although it is better than number of seedlings per acre. A map showing distribution of stocked plots within a cutting is probably best. In this study, stocked plots were generally well distributed on small cuttings but confined to the outer area of the staggered-setting cutting.

Table 2.--Stocking established seedlings on standard and modified clear cuts,

:			Stocking					
Unit :	Kind of cutting unit	Milacre plots	Plots stocked, all species	: Plots stocked, : Douglas-fir :	: Douglas-fir seedlings per acre			
		Number	Percent	Percent	Number			
S1 S2 S3 S4	North-south 50-foot strip North-south 100-foot strip North-south 200-foot strip North-south 350-foot strip	102 118 96 141	61 35 40 60	57 31 34 45	1,466 597 573 908			
S5 S6 S7	East-west 100-foot strip East-west 200-foot strip East-west 300-foot strip	90 99 174	81 89 69	6 8 81 59	1,733 4,561 1,899			
G1 G2 G3	4-acre square patch 2-acre square patch 1-acre square patch	204 116 102	58 60 81	42 29 45	824 474 814			
04 G5 G6	<pre>2/4-acre circular patch 1/2-acre circular patch 1/4-acre circular patch</pre>	114 107 130	75 72 72	58 64 44	1,658 2,276 888			
5B	Staggered-setting	225	28	21	407			
50	Seed-tree	125	61	50	1,448			

H. J. Andrews Experimental Forest, 1959

The wide variation in level of stocking among units essentially homogeneous in character is noteworthy. For example, the three east-west strip clear cuts occupy similar topography, but percent of plots stocked with Douglasfir varied from 59 to 81 and number of Douglas-fir seedlings varied from 1,733 to 4,561 per acre.

REGENERATION AND GROSS CHARACTERISTICS OF CLEAR CUTS

The first analyses were made to determine if stocking levels were related to measurable characteristics of the 13 small clear cuts. Three different measures of stocking were used: (1) percent of plots stocked with all species, (2) percent of plots stocked with Douglas-fir, and (3) number of Douglas-fir seedlings per acre. Characteristics of clear cuts were expressed as: (1) size of clear cut, (2) average hours of shade on clear cut, (3) width of clear cut in narrowest dimension, (4) width of clear cut in north-south dimension, and (5) average distance of plots from stand edge. In the regression analyses, all relationships were tested, using each clear cut as a single observation.

Only one relationship was revealed to be $\operatorname{significant}_{5/}^{5/}$ -the relation of percent of milacre plots stocked with any species and the north-south width of the clear cut. For the 13 small clear cuts sampled, apparently no relationship exists between stocking of either Douglas-fir or all species and size of clear cut, average hours of shade on clear cut, width of clear cut in narrowest dimension, or average distance of plots from stand edge.

REGENERATION AND CHARACTERISTICS OF MILACRE PLOTS

The failure of the initial analysis to reveal relationships between stocking levels and gross characteristics of clear cuts prompted a closer look at milacre-plot data on individual clear-cut units. Additional regression analyses were carried out to determine if number of Douglas-fir seedlings per plot was related to (1) hours of shade on plot or (2) distance of plot from seed source. These two analyses were conducted for each of the 13 patch and strip clear cuts. In addition, plot data from each of the major groups of cuttings--east-west strips, north-south strips, and patch clear cuts--were pooled and analyzed in the same manner.

Stand Shade

Stand shade favored establishment of natural regeneration. On some cuttings, this was apparent simply from examination of plotted data (fig. 1). The statistical analyses revealed that a significant relationship existed between

 $[\]frac{5}{1}$ In this paper, "significant" is used in a statistical sense and means that a relationship was shown to exist at the 5-percent level of probability. "Highly significant" means a relationship was shown to exist at the 1-percent level of probability.

hours of shade and number of Douglas-fir seedlings per plot on 8 of 13 units. The pooled data from all three major types of cuttings showed a highly significant correlation between hours of shade and number of Douglas-fir seedlings per plot.



Figure 1. --Relation between stocking and distance from the south edge of timber on two east-west oriented strip clear cuts. Unit S5 is 100 feet wide and unit S6 is 200 feet wide.

In general, the relation between hours of shade and regeneration was more marked on east-west strip clear cuts and patch clear cuts than on northsouth strip clear cuts. A partial explanation of differences between units is provided by Silen.⁶ He found seedlings must be exposed to a high soil surface temperature for a continuous period of time before heat damage occurs to the seedling. Apparently, intermittent sunlight is less damaging than an equal duration of continuous sunlight. Only part of the variation between units can be explained in this way; other factors, such as time of day plot receives stand shade, pattern of ground shade, and microexposure, doubtless play a major role.

How much shade can be expected from a particular stand edge? The theoretical relationship of tree height, percent slope, angle of sun, and stand shade is shown in figure 2. In practice, this relationship is modified by the continuity and crown condition of the stand providing shade. Plotted results from two units (fig. 1) indicate that a timber edge on the south affects a smaller

 $\frac{6}{-}$ See footnote 3, p. 4.

-7-



area than indicated by the diagram. The differences between the obviously shade-influenced areas on these units (approximately 35 feet and 70 feet) reflect differences in the shade effects of the timber edge. $\frac{7}{2}$

Distance from Seed Source

The number of Douglas-fir seedlings per plot was generally not related to distance from seed source. All portions of strip and patch clear cuts were within 200 feet (and usually much less) of a timber edge. Since Douglas-fir seed is readily carried much greater distances (Isaac, 1943), a relation between stocking and distance from seed source would not be expected on these units. On a few cuttings, there were significant relations between regeneration and distance

 $\frac{7}{}$ These units were chosen because the relationship between stand shade and stocking is least influenced by extraneous factors such as roads and cutting boundary irregularities.

-8-

from stand edge, but these probably resulted from the partial relationship which exists between hours of shade and distance from stand edge. However, on the staggered-setting clear cut, the relation between stocking and distance from stand edge was highly significant:

	Total	Plots	Percent
Distance from stand edge	plots	stocked	stocked
Under 200 feet	106	52	49
Over 200 feet	119	10	8

Worthington (1953) reported similar results after studying regeneration on a large clear cut and some small patch clear cuts 1.2 to 4.0 acres in size in western Washington. Distance from stand edge affected stocking only beyond 500 feet on the large clear cut. None of the small patch cuts he studied had sample points over 500 feet from stand edge, and no relation between stocking and distance from stand edge was therefore found on these units.

This also explains the lack of relationship between clear cut size and stocking mentioned earlier. The patch and strip clear cuts were simply too small to show such a relation. Lavender et al. (1956) studied much larger clear cuts and concluded there was no relation between the size of clear cut and regeneration.

DISCUSSION OF THE MODIFIED CUTTINGS

North-South Strip Clear Cuts

Although north-south strip clear cuts had the poorest stocking of the modified cuttings, they were surprisingly well stocked considering the severity of the site. Analysis of the individual north-south strip clear cuts generally indicated no significant relation between number of Douglas-fir seedlings per plot and hours of shade or distance from seed source. Lack of a relation between shade hours and regeneration can be explained, at least in part, by the continuous rather than intermittent character of the shade (fig. 3). There does not appear to be any significant relation between stocking and width of north-south strip.

East-West Strip Clear Cuts

The long timber edge available for shading regeneration on the east-west strip clear cuts was probably responsible for superior stocking on these units. Furthermore, much of the shading was of the intermittent type. Analysis of data from individual units and of the pooled data indicated the number of Douglasfir seedlings per plot was significantly related to hours of shade.

No star Indentification de la constante



Figure 3.--Pattern of shading on north-south oriented strip clear cuts. Top, in the morning; middle, atmidday; bottom, in the afternoon.





Patch Clear Cuts

Stocking of patch clear cuts was generally intermediate between the eastwest and north-south strip clear cuts. As a group, the patch clear cuts showed a consistent relationship between shade and number of Douglas-fir seedlings per plot. Two exceptions were units G2 and G5. Part of unit G2 was a low swampy area which offered little mineral seedbed. More than half of the stocked plots contained western hemlock and/or western redcedar; and 29 percent of the plots were stocked with Douglas-fir. Unit G5 had a road as its southern boundary and the effective environment of this clear cut was larger than its half-acre size would indicate. In addition, a second unit just across the road from G5 was cut in the spring of 1959. This disturbance made it impossible to obtain a valid estimate of amount of shading during the previous 4 years.

Most references to strip and patch clear cutting list the main advantage as a provision for an effective seed source (Haig et al., 1941; Hawley and Smith, 1954; Heinselman, 1959; Lexen, 1949). Hawley and Smith (1954) mention in addition that "...this kind of modification [strips or patches] may also be employed...to create an environmental condition intermediate between the protection of a shelterwood and the drastic exposure of a large clear cut area." Troup (1952) emphasizes the protective function of such cuttings. A system known as Wagner's "Blendersaumschlag" involves narrow strip cuttings oriented east and west which progress from north to south with the specific objective of protecting seedlings from the sun on the strip being regenerated. In European clear-strip systems, cuttings are often restricted to widths no more than the height of adjacent forest in order to secure side protection from the sun for either artificial or natural regeneration. Results of the current study suggest that patch and strip clear cuttings for Douglas-fir are primarily of value in aiding seedling establishment.

Seed-Tree Cutting

The seed-tree cutting was a small staggered-setting on which three dozen Douglas-firs were left on two-thirds of the area to supply seed and shade (fig. 4). Trees were selected for vigor and at a spacing of approximately 120 feet; slash was pulled away from the seed trees during logging to prevent damage during slash burning.

Figure 4. --General view of seed-tree cutting.



The seed-tree portion of unit 5C was well stocked 4 years after logging (table 2). An important factor in prompt seedling establishment was probably the shade cast by seed trees. Garman (1955) found that a full-crowned, mature Douglas-fir seed tree gives temporary protection to an area of one-third acre during critical times of the year. Mineral seedbed resulting from yarding and slash-disposal operations and additional seed produced by seed trees were other important factors. Seed trees have been found to be heavier cone producers and more efficient distributors of seed than marginal timber (Garman, 1951; Isaac, 1940). Poorest stocking was on the portion of the unit without seed trees, but a statistical comparison was not possible because of variations in slope, exposure, steepness, and edge-shade effects.

Careful selection of sound, vigorous, and apparently windfirm seed trees and prevention of damage to them during logging-slash burning resulted in almost 100-percent seed-tree survival. None of the Douglas-fir trees left in 1955 had blown down and only two had died from exposure as of September 1961. Most were full crowned and had been producing seed.

Several studies of Douglas-fir regeneration have indicated the value of seed trees for regeneration. Garman (1955) found that shade definitely improved seedling survival in British Columbia. He felt that shade from two to eight seed trees per acre was indicated on severe sites and that cutting of Douglas-fir would eventually evolve to harvesting in 300-foot-wide strips (on good sites) and a shelterwood system on more severe sites. He concluded, "Douglas-fir should be cut by a method that leaves the shade of scattered seed trees." Bever (1954), following a study of factors affecting natural regeneration in central western Oregon, concluded the seed-source value of seed trees was higher than marginal timber or seed blocks. In contrast, early studies of seed trees indicated mortality, wind, logging, and slash burning quickly destroyed most of the trees (Isaac, 1940).

CONCLUSIONS

Strip and patch clear cuttings and a seed-tree cutting were adequately stocked with seedlings of Douglas-fir and associated species within 4 years after logging and slash burning. There was, however, a great deal of variability in level of stocking between and within the experimental cuttings. A staggeredsetting clear cut of comparable age was.not adequately stocked. Quantity of Douglas-fir regeneration was not related to gross characteristics of the cutting areas such as size or width. However, stocking of seedlings of all species was related to the north-south width of the clear cut. Analyses of the plot data from individual clear cuts showed that, of several factors studied, number of hours of stand shade on a plot was most closely related to number of Douglas-fir seedlings. Regeneration was not related to distance from seed source on the relatively small patch and strip clear cuts.

The chief value of modified cutting systems appeared to be in seedling establishment, particularly through reduction of insolation losses. Intermittent

shade may have been more effective than a similar amount of shade received in a single period of time, although this was not demonstrated conclusively.

It is recommended that consideration be given to the use of modified cutting methods such as east-west strip clear cuts, small patch clear cuts, and seed-tree cuttings to improve natural restocking of Douglas-fir on sites difficult to regenerate. Such cuttings should be laid out primarily to provide shade and secondarily to favor seed dispersal.

LITERATURE CITED

Berntsen, Carl M., and Rothacher, Jack.

Bever, Dale N.

1954. Evaluation of factors affecting natural reproduction of forest trees in central western Oregon. Oreg. State Bd. Forestry Res. Bul.
3, 49 pp., illus.

Garman, E. H.

- 1951. Seed production by conifers in the coastal region of British Columbia related to dissemination and regeneration. Brit. Columbia Forest Serv. Tech. Pub. T. 35, 47 pp., illus.
- 1955. Regeneration problems and their silvicultural significance in the coastal forests of British Columbia. Brit. Columbia Forest Serv. Tech. Pub. T. 41, 67 pp., illus.

Haig, Irvine T., Davis, Kenneth P., and Weidman, Robert H.

1941. Natural regeneration in the western white pine type. U.S. Dept. Agr. Tech. Bul. 767, 99 pp., illus.

Hawley, Ralph C., and Smith, David M.

1954. The practice of silviculture. Ed. 6, 525 pp., illus. New York: John Wiley & Sons.

Heinselman, M. L.

- 1959. Natural regeneration of swamp black spruce in Minnesota under various cutting systems. U.S. Dept. Agr. Prod. Res. Rpt. 32, 22 pp., illus.
- Isaac, Leo A. 1940. Mortality of Douglas-fir seed trees on cutover lands. In U.S. Forest Serv. Pac. NW. Forest Expt. Sta. Res. Note 31.pp. 11-12.

^{1959.} A guide to the H. J. Andrews Experimental Forest. U.S. Forest Serv. Pac. NW. Forest & Range Expt. Sta., 21 pp., illus.

Isaac, Leo A.

1943. Reproductive habits of Douglas-fir. 107 pp., illus. Washington D.C.: Charles Lathrop Pack Forestry Foundation.

Lavender, Denis P., Bergman, Morris H., and Calvin, Lyle D. 1956. Natural regeneration on staggered settings. Oreg. State Bd. Forestry Res. Bul. 10, 36 pp., illus.

Lexen, Bert.

1949. Alternate clear-strip cutting in the lodgepole pine type. U.S. Forest Serv. Rocky Mountain Forest & Range Expt. Sta. Sta. Paper 1, 20 pp., illus.

Troup, R. S.

1952. Silvicultural systems. Ed. 2, 216 pp., illus. London: Oxford Univ. Press.

Worthington, Norman P.

1953. Reproduction following small group cuttings in virgin Douglas-fir. U.S. Forest Serv. Pac. NW. Forest & Range Expt. Sta. Res. Note 84, 5 pp.

Yerkes, Vern P.

1960. Occurrence of shrubs and herbaceous vegetation after clear cutting old-growth Douglas-fir in the Oregon Cascades. U.S. Forest Serv. Pac. NW. Forest & Range Expt. Sta. Res. Paper 34, 12 pp.