

## SALMON POPULATIONS AND LARGE SCALE ATMOSPHERIC EVENTS

David Greenland  
Department of Geography, University of Oregon  
Eugene, Oregon

### ABSTRACT

Catches of coho salmon off the coast of Washington and Oregon since 1925 appear to be related to large scale events in the atmosphere which in turn affect ocean currents and coastal upwelling intensities in the northeastern Pacific ocean. At least two time scales of variation can be identified. The first is that of the El Niño/Southern Oscillation phenomenon giving rise to an irregular cycle of between 3 to 7 years. The occurrence of El Niños is associated with warm water anomalies off the coast and often with depressed salmon catches. The second time scale of variation seems to have a periodicity of about 20 years, although this is based on a limited data set. This variation has been postulated to be related to the manner in which the Subarctic ocean current operates; sometimes directing more nutrient-rich water into the Alaskan current which moves it northwards into the Alaskan gyre, while at other times guiding more nutrient-rich water southwards into the Californian current off the shores of Washington and Oregon. During the former times there tends to be low coho salmon catches off the Washington and Oregon coasts and the opposite also seems to hold true. The difference in these modes of the Subarctic current may be related to long-term differences in the mode of operation of the atmospheric currents. The atmospheric currents may be quantitatively described by the use of teleconnective indices which include: the Southern Oscillation Index (SOI) and the Central North Pacific (CNP) and Pacific North American (PNA). The values of these indices also have some explanatory power, at least in winter, over temperatures and precipitation values in the Pacific Northwest. This paper endeavors to describe how, if real, these atmospheric-oceanic effects are integrated and might affect the salmon catch. The possibility must also be considered that the atmospheric events are symbiotically related to the oceanic events and further that both may be enmeshed in even longer term variability of climate.

### INTRODUCTION

The story of salmon in the terrestrial-based freshwater aquatic ecosystems of the north west of the North American continent and the marine ecosystems of the north Pacific ocean is a fascinating and multi-faceted one. Humans suffer from a lack of knowledge of many of the facets of the tale. One of the aspects which, until recently, has received little attention is that of the long chain of events which starts, at one end with variations in the atmosphere and ends at the other with variability of population sizes of certain kinds of salmon. Climate is one factor which effects salmon population sizes either directly or indirectly in both their freshwater and marine phases.

This paper will review some of the aspects of the effect of climate on salmon. These aspects occur on a variety of time scales. It will then outline a statistical relationship which appears to exist on an inter-decadal time scale between the population size of coho salmon as indicated by the catch of these fish off the coast of Washington and Oregon and the air temperature in parts of Oregon. An attempt to explain this relationship requires an examination of the large scale variability and flow of air currents in the atmosphere of the western part of the Northern hemisphere and the possibly related variation of ocean currents in the north eastern Pacific ocean. Suggestions will be made for further research which could provide new insights into these longer term relationships.

### **Climate-Salmon Interactions in General**

A cursory look at the literature reveals several important links between salmon population sizes and climatic variables. These links occur both directly and indirectly, and at a variety of time and space scales.

An example of a small time scale link is the small time window of a matter of a few days during which juvenile salmon embark on their journey from the estuary to the sea. Adverse weather conditions, such as severe storms with high onshore winds, during this period could negatively impact survival. Spence and Hall (1994) have shown that the size of this window increases from high to low latitudes. This fact makes storms and other potentially adverse climatic conditions in the higher latitude estuaries and river mouths particularly critical.

On a monthly time scale it has been shown that while salmon are young, colder river water temperature slows down growth and warmer water accelerates growth (Netboy, 1980). However, a temperature increase in rivers generally reduces survival because it may be associated with an increase in disease and fungal attacks in adults.

On a seasonal and annual time scale there appear to be many climatic effects on salmon populations. The amount of water in rivers as controlled by periods of high and low flow conditions is critical for salmon survival. Sharp (1992) claims that local populations of salmon throughout the western continental United States have suffered from drought during the period 1976 to 1991. The relationship of salmon population size to El Niño conditions, which also occurs on this time scale, will be treated more specifically below. With the exception of the one discussed later, not many studies have been performed concerning the relationship between climate and salmon on decadal and century time scales. On a millennial time scale there is limited and somewhat tenuous evidence to suggest that the inverse relationship of warmer climate and decreased salmonids, found on the El Niño type time scale, may persist. Neitzel et al., (1991) believe that during the Hypsithermal period of the Holocene, salmon were less plentiful in the Pacific Northwest (PNW).

One very important point, which is often neglected, is that the fossil salmon record extends back to the Eocene and Miocene (Percy, 1992, p 5). The implication of this is at least

some species of salmon have survived just about every conceivable extreme climate and type of climate variability which the planet can offer, or is likely to offer in the next century.

### **El Niños and Salmon**

El Niño occurrences can impact salmon populations in a number of ways. During El Niños anomalously warmer water is found off the coast of North America suppressing coastal upwelling and its associated nutrients. There is evidence that the 1957-58 El Niño negatively affected salmon giving rise, in 1960, to the lowest ocean landings off the continental U.S. coast since 1917 (Pearcy, 1992). In addition, the lingering 1991-92 El Niño also negatively affected salmon catch (Finley, 1993). It has been established that years just following El Niño events generally tend to be years of low coho salmon catches (Miller and Fluharty, 1992).

One of the more spectacular and intense El Niño events occurred in 1982-83. During this event salmon in the PNW appeared to be stalled in estuaries because of the warmer water in the ocean. The salmon in the estuaries were subject to disease and predation which decreased the populations. In addition, reduced primary productivity off the coast impacted both juvenile and adult coho salmon (Pearcy, 1992). During this El Niño event 58% of the adults predicted to return actually died in the ocean. The same El Niño markedly decreased growth and fecundity of the salmon of the year. Warm ocean temperatures off the Oregon coast shifted the center of distribution of juvenile coho salmon northwards to the coast of Washington.

Another feature of El Niños which does not seem to have received much attention is that they are often accompanied in the PNW by warm, dry winters (Redmond and Koch, 1991, Greenland, 1994). Whether the lack of precipitation leads to stream discharge values low enough to affect salmon has not been investigated explicitly.

### **Interdecadal Relations Between PNW Air Temperatures and Coho Salmon Catch off the Coast of Washington and Oregon**

When temporally smoothed data are used for the period 1925 to 1985 there is a close inverse statistical relationship between the PNW Air Temperatures and coho salmon catch off the coast of Washington and Oregon. This relationship was discovered by using the following steps. Frances and Sibley (1991) had previously reported a close relationship between winter (November to March) air temperatures at Sitka, AK and surface water temperature at Langara Spit, Queen Charlotte Island, B.C. and the catch of pink salmon in the Gulf of Alaska. They had also reported an inverse relationship between the catch of pink salmon in the Gulf of Alaska and the catch of coho salmon off the coast of Washington and Oregon. Both relationships covered the period 1925 to 1985 and were found when the data were normalized and subjected to a 7 year weighted filter (Frances, pers. comm. 1994). Given these relationships I reasoned that it would be likely that there would be an inverse relationship between air temperatures in the PNW and the catch of coho salmon catch off the coast of Washington and Oregon. I used a 5 year unweighted filter of the annual mean air temperatures at the H.J. Andrews Long-Term Ecological Research site. The filter was applied to values of

Andrews' temperatures which were normalized to the long-term mean for the 1925-1985 period. It has been shown elsewhere that the Andrews' temperatures are well related to those of western Oregon in the general (Greenland, 1994). Coho salmon catch data were extracted from the graphs of Frances and Sibley and a close inverse relationship was found (Figure 1). This relationship and time series suggest an approximate 20 year cycle in both air temperatures in the PNW and the catch of coho salmon catch off the coast of Washington and Oregon. In order to attempt to explain this inverse relationship we must explore the large spatial and long temporal aspects of atmospheric and ocean currents as well as further thinking of Frances and Sibley and their colleagues.

### **Large Spatial and Long Temporal Aspects of Atmospheric and Ocean Currents**

It is well known that the currents of the atmosphere and ocean are related. Airflow over the surface of the ocean is one of the major driving forces of the ocean currents. Ocean flow in the eastern North Pacific is dominated by the Subarctic current and West Wind Drift, to its south, which together transport water eastward approximately between latitudes 40 and 50° N. This flow is driven by the westerly atmospheric flow at these latitudes. Towards the east of the ocean the flow of these currents bifurcates. Part of it flows northward, as the Alaskan current, along the coast of British Columbia and then westward along the southern coast of Alaska to form the Alaskan Gyre. Another part flows southward forming the Californian current, part of which flows eastward, under the influence of the Trade Winds when it reaches about 20° N. The West Wind Drift, California Current and return westward flow of the North Equatorial current forms part of the Central Pacific Gyre.

In two parts of these flows nutrient-rich water can be brought from lower levels of the ocean by upwelling. Along the coast of the conterminous U.S. this is achieved by an application of the Coriolis force to coastal airflow (with a northerly component in summer) and coastal ocean current flow. The Coriolis force tends to deflect objects to the right of their path of motion in the Northern Hemisphere. This leads to offshore and divergent surface ocean flow which, in turn, draws the nutrient rich water up from below. There is also upwelling on the northern side of the sub-arctic current in southern part of the Alaskan gyre. This upwelling is due to a process called Ekman pumping which is caused by the change of direction of flow of the faster velocity water near the surface compared to the slower velocity water at increasing depths in the ocean.

The winds in the atmosphere above the ocean are responsible for both the velocity of the ocean currents and their precise location from year to year and from decade to decade. They may also be responsible for the precise location of the bifurcation of the subarctic current and the way in which quantities of nutrient-rich water are partitioned into the Alaskan and California currents.

The atmospheric currents of importance here are the winter flow of the mid-latitude westerly winds and to a lesser extent the flow of the easterly Trade winds. At the surface of the Earth the westerlies in winter are manifested by geostrophic flow of air at the southern part of the Aleutian Low pressure system. In the upper air at about 18-20,000 feet above the

Earth's surface the flow takes the form of long waves (parallel to the earth's surface). The position and strength of the air flow in the waves is variable but on the average is characterized by a ridge of high pressure over the eastern Pacific sandwiched between a trough of low pressure in the western Pacific, off the coast of Asia, and another trough over eastern North America and the western Atlantic ocean (Barry and Chorley, 1987. p136). The actual position and strength of the Aleutian Low and the position, degree of sinuosity (or amplitude of the waves), and strength of the flow in the upper air westerlies from year to year and from decade to decade is critical to controlling the flow and location of the ocean currents.

Climatologists have developed some simple indices to collapse a large amount of this atmospheric flow into a few numbers. The indices are sometimes called teleconnective indices since they help explain how pressure variations in one part of the hemisphere are related (teleconnected) to those in another part. The indices of use in this discussion are the Central North Pacific index (CNP), the Pacific North American index (PNA) and the Southern Oscillation Index (SOI). It should also be remembered that the strength of the teleconnective patterns is not necessarily stable over time.

The SOI is commonly used to measure the strength of the El Niño and, the opposite, La Niña, phenomena. The SOI is measured as the mean sea level pressure (MSLP) difference between Tahiti and Darwin, Australia. Due partly to difficulties in terminology it is becoming increasingly common to refer to "warm events" (which include El Ninos) and "cold events" (which include La Nina). In the newer terminology "warm" and "cold" refer to the sea surface temperatures (SSTs) of the central Pacific Ocean.

Wallace and Gutzler (1981) were the first to introduce the PNA index. They characterized a strong winter PNA pattern as one associated with higher than normal temperatures in the PNW resulting from a strong ridge of high pressure in the 700 mb height field extending from the PNW to Canada and Alaska. Wallace and Gutzler use a PNA index comprised of the pressure at four points on the latitude, longitude grid. The PNA index designed by Leathers et al. (1991) and Leathers and Palecki (1992) following Yarnal and Diaz (1986) is the one used in this study. It uses a linear combination of standardized 700 mb geopotential height anomalies ( $Z^*$ ) at the grid points nearest the anomaly field centers. It is constructed as follows:

$$PNA = 1/3\{-Z^*(47.9^\circ N, 170.0^\circ W) + Z^*(49.0^\circ N, 111.0^\circ W) - Z^*(29.7^\circ N, 86.3^\circ W)\}.$$

The PNA describes the amplitude of the 700 mb flow pattern over the U.S. which has a basic pattern of troughs of low pressure in the eastern Pacific and the eastern U.S, and a ridge of high pressure over the Rocky Mountain cordillera (see Leathers et al. 1991. Fig.1). The meridional extreme of the pattern produces positive PNA values (and potentially more SW winds over the PNW) while the zonal extreme produces negative PNA values (and potentially more W winds over the PNW). Yarnal and Diaz (1986) demonstrated how strongly positive PNA and negative (reverse) PNA patterns are associated respectively with warm and cold El

Niño/Southern Oscillation (ENSO) events and, in turn, with precipitation and temperature anomalies on the west coast of North America.

Cayan and Peterson (1989) designed the CNP index as being the MSLP over the region 35-55° N and 170° E to 150° W. They show that streamflow in the West have correlations in the range 0.3 to 0.6 SLP with anomalies in the North Pacific. During times of a weak CNP streamflows are high in Washington and Oregon. During times of a strong CNP the polar front jet stream flows north of the PNW and times of below average streamflow are observed. This is also often observed during El Niño events (See Cayan and Peterson, 1989, Fig. 9).

It has also been shown that there are interrelationships between the values of the SOI, PNA and the CNP indices (Cayan and Peterson, 1989). Significant correlations appear between the SOI and PNA during winter and spring, the SOI and the CNP during winter, and the PNA and the CNP during winter spring and fall. All of the synoptic studies indicate quite clearly the linkages which exist between SSTs and particular pressure and teleconnective patterns in the Pacific Ocean and various aspects of climate on the West Coast and the in the PNW.

### **The HWFS Model and its Relationship to Atmospheric Flow**

In order to explain the relations between the catches of Alaskan pink salmon and northern sea and air temperatures and the inverse relationship between the catches of Gulf of Alaskan pink salmon and Washington/Oregon coho salmon Frances and Sibley (1991) and Frances (1993) make use of a model suggested by Hollowed and Wooster (1991). For convenience I will call this model the HWFS after the initial letters of all four investigators. According to this model the inverse relationship in catch in the two ocean areas might be explained by the north or south movement of the of the divergence, or bifurcation, zone between the Alaskan and California current and the greater or lesser effect of the currents related to that north-south movement. The model is bimodal and postulates two states or modes of operation of the ocean currents. When the bifurcation zone is more to the north (HWFS Type A - West Coast) more cold Sub-arctic current water is taken into the Californian current and upwelling of nutrient-rich water off the Oregon and Washington coasts is enhanced. When the bifurcation zone is further to the south (HWFS Type B - Alaska) more cold Sub-arctic current water is taken into the Alaskan current and water off the Oregon and Washington coasts will be warmer.

### **The Synthesis of the Teleconnective Indices with the HWFS Model**

The HWFS model may be placed in a larger atmospheric context by noting its relationship to the teleconnective index values discussed earlier (Figure 2). HWFS Type A West Coast flow is associated with a high CNP value. This describes a weak Aleutian low pressure with winds coming more directly from the west across the Pacific at the latitudes of Washington and Oregon. This is consistent with a negative value of the PNA. The more northerly bifurcation of the Sub-arctic current pushes more water into the Californian current and gives rise too negative SST anomalies. These conditions are not generally consistent with

intense El Niño conditions. HWFS Type B Alaska flow is associated with a low CNP value. This describes a strong Aleutian low pressure and enhanced south westerly winds in the northeast Pacific. It is associated with meridional upper air flow and a positive PNA value. The more southerly Subarctic current bifurcation enhances northward ocean flow into the Alaskan current giving rise to positive SST anomalies in the eastern part of the NE Pacific. These conditions are consistent with the results of El Niño events.

Hollowed and Wooster (1991) and Frances (1993) have suggested that El Niños are not unrelated to the change of state between Type A and Type B flow. They suggest that over the last 60 years the switch from Type A mode to Type B mode has always occurred at the time of significant El Niño events (e.g. 1925/26, 1940/41). This relationship, if true, is worthy of further investigation.

### FURTHER RESEARCH AND CONCLUDING COMMENTS

There is a large amount of research to be completed before the ideas discussed above can be finalized and confirmed by observational data. Some of the more important research tasks include:

- 1) Examine individual El Niños and salmon catch off Washington and Oregon with respect to the entire available salmon catch record and the intensity values of individual El Niños.
- 2) Examine in more detail what are the exact biophysical processes leading to the decrease of salmon population size during El Niño years.
- 3) Further identify the variability of the bifurcation of ocean water flow from the central north Pacific into the Alaska and Californian Currents. Possible tools for doing this include: a) an examination of ocean temperature and salinity and other flow-identifying data from ocean weather ship P, and b) the use of remote sensing imagery.
- 4) The PNW air temperature/coho salmon catch should be recalculated using the more representative Climate Divisional air temperature records of the National Weather Service and a temporally coherent salmon catch data series updated into the mid 1990s.
- 5) These analyses should be extended to look for relations between salmon catch and the teleconnective indices. Work on this topic has been started by Jamir et al. (this volume).
- 6) It can be hypothesized that intense El Niño conditions more often occur under HWFS Type B mode conditions than Type A mode conditions. This hypothesis should be tested.
- 7) More research should be carried out concerning the flip-over between the HWFS Type A to Type B mode and vice versa. In particular we should investigate the role, if any, of El Niño events in this as well as the relation of the mode change to known major step functions in atmospheric flow such as that which occurred in 1976 (Trenberth, 1990).

8) If the HWFS model is confirmed it then needs to be asked a) how is it effected by long term climate change at scales longer than the interdecadal scale, and b) what feedbacks are there from the ocean to the atmosphere and vice versa and how do they operate? With respect to b) Namias, has pointed out, in a whole series of papers (e.g. Namias 1969), the importance of the two way interaction between the Pacific ocean and the atmosphere above it. This needs to be taken into account in a more time specific framework with respect to the HWFS model.

If the results of these tasks and other investigations confirm the existence of interdecadal cycles related to the atmospheric and ocean current operation such confirmation will have important management implications for the salmon resource. Not the least, it will be important to cast any management plans within a framework of interdecadal natural variability which may also have some predictability. Possibly more important is to further examine the historic and paleo record to further establish the strength of the variability and possibility of forecasting it in the future.

#### ACKNOWLEDGEMENTS

This preparation of this paper was supported in part by Cooperative Agreement No. PNW 92-0221 from the Pacific Northwest Research Station of the US Forest Service and the Long-Term Ecological Research Grant number BSR-9011663 from the National Science Foundation. I am grateful to Robert Frances of the School of Fisheries at the University of Washington for sharing some of his original data and unpublished notes on the HWFS model.

#### REFERENCES

- Barry, R.G. and Chorley, R.J. 1987. Atmosphere, Weather, and Climate. 5th Ed. Routledge. London. 460 pp.
- Cayan, D.R., and Peterson, D.H. 1989. The Influence of North Pacific Circulation on Streamflow in the West. in Aspects of climate variability in the Pacific and western Americas. Ed. D.H. Peterson. American Geophysical Union. 445 pp. 375-397.
- Francis, R.C. 1993. Climate change and salmonid production in the North Pacific Ocean. pp. 33-42 in Proceedings of the Ninth Annual Pacific Climate (PACLIM) Workshop. California Department of Water Resources.
- Francis, R.C. and Sibley, T.H. 1991. Climate Change and Fisheries: What are the real issues?. The Northwest Environmental Journal. 7:295-307.
- Finley, C. 1993. Low coho return may halt '94 ocean season. The Oregonian Nov. 18.

- Greenland, D. 1994. The Pacific Northwest Regional Context of the Climate of the H.J. Andrews Experimental Forest Long-Term ecological research Site. Northwest Science. in press.
- Hollowed, A.B. and Wooster, W.S. 1991. Year class variability of marine fishes and winter ocean conditions in the Northeast Pacific Ocean. ICES Variability Symposium. Paper No. 33. Session 3.
- Leathers, D.J., Yarnal, B., and Palecki, M.A. 1991. The Pacific/North American teleconnection pattern and United States climate. Part I: Regional temperature and precipitation associations. *Journal of Climate*. 4(5)517-528.
- Leathers, D.J. and Palecki, M.A. 1992. The Pacific/North American teleconnection pattern and the United States climate. Part ii: Temporal Characteristics and index specification. *Journal of Climate*. 5:707-716.
- Miller, K.A. and Fluharty, D.L. 1992. El Niño and Variability in the Northeast Pacific Salmon industry: implications for coping with climate change. pp. 49-88 in M.H. Glantz. (Ed.). *Climate variability, climate change and fisheries*. Cambridge University Press. New York. 450 pp.
- Namias, J., 1969. Seasonal interactions between the North Pacific Ocean and the atmosphere during the 1960s. *Monthly Weather Review* 97(3)173-192.
- Netboy, A. 1980. *The Columbia River salmon and steelhead trout: their fight for survival*. University of Washington Press. Seattle. WA.
- Neitzel, D.A., Scott, M.J., Shankle, S.A., and Chatters, J.C. 1991. The effect of climatic change on stream environments: The salmonid resource of the Columbia river basin. *The Northwest Environmental Journal*. 7:271-293.
- Pearcy, W.G. 1992. *Ocean ecology of North Pacific salmonids*. University of Washington Press, Seattle. WA. 179 pp.
- Redmond, K.T. and Koch, R.W. 1991. Surface climate and streamflow variability in the western United States and their relationship to large scale circulation indices. *Water Resources Research*. 27:2381-2399.
- Sharp, G. 1992. Fishery catch records, El Niño / Southern Oscillation, and longer term climate change as inferred from fish remains in marine sediments. pp 379 - 417. H.F. Diaz and V. Markgraf (Eds.). Cambridge University Press. Cambridge. 476 pp.
- Spence, B.C. and Hall, J.D. 1994. Analysis of spatio-temporal variation in the migration timing of Coho Salmon (*Oncorhynchus kisutch*) smolts and implications of climate

change. Abstract. Oregon Chapter. American Fisheries Society. 30th Annual Meeting. Sunriver, OR. 16-18 February. p.19.

Trenberth, K.E. 1990. Recent observed interdecadal climate changes in the northern hemisphere. *Bulletin of the American Meteorological Society*. 71(7)988-993.

Wallace, J.M. and Gutzler, D.S. 1981. Teleconnections in the geopotential height field during the Northern Hemisphere winter. *Monthly Weather Review*. 109:784-812.

Yarnal, B. and Diaz, H.F. 1986. Relationships between extremes of the Southern Oscillation and the winter climate of the Anglo-American Pacific Coast. *Journal of Climatology*. 6:197-219.




Figure 1. Smoothed values (salmon catch 7 yr. filter, Temperature 5 yr. filter) of annual mean temperatures at the H.J Andrews Experimental Forest and the catch of coho salmon off the coast of Washington and Oregon between 1927 and 1983.





Figure 2. Two types of flow hypothesized in the HWFS model and its associated atmospheric and oceanic events.



For copies, send \$15 U.S. (includes postage) to:  
Oregon Chapter, American Fisheries Society  
P.O. Box 722  
Corvallis, OR 97339

Contents of this publication do not necessarily reflect the view and policies of the American Fisheries Society, Oregon Department of Fish and Wildlife, National Marine Fisheries Service, Bureau of Land Management, U. S. Forest Service, or the Bonneville Power Administration, nor does mention of trade names or commercial products constitute their endorsement by any above-named groups.

**Salmon Ecosystem Restoration:  
MYTH AND REALITY**

**Proceedings of the 1994 Northeast Pacific  
Chinook and Coho Salmon Workshop**

**Eugene, Oregon  
November 7-10, 1994**

**Edited by  
MaryLouise Keefe**

**Sponsored by  
Oregon Department of Fish and Wildlife  
National Marine Fisheries Service, Northwest Region  
Bureau of Land Management  
U. S. Forest Service  
Bonneville Power Administration**

**This document has not been peer-reviewed.**